

**COPPERTONE SPORT SUNCREEN SPF 50- avobenzone 3%, homosalate 10%, octisalate 5%, octocrylene 5% aerosol, spray
Beiersdorf Inc**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Coppertone Sport Spray SPF 50

Drug Facts

Active ingredients

Avobenzone 3%, Homosalate 10%, Octisalate 5%, Octocrylene 5%

Purpose

Sunscreen

Use

■ helps prevent sunburn

Warnings

For external use only

Flammable: Do not use near heat, flame, or while smoking

Do not use on damaged or broken skin

When using this product

■ keep away from face to avoid breathing it

■ keep out of eyes. Rinse with water to remove.

■ use only as directed. Intentional misuse by deliberately concentrating and inhaling the contents can be harmful or fatal.

■ do not puncture or incinerate. Contents under pressure. Do not store at temperatures above 120°F.

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Directions

- shake well before each use
- spray liberally and spread evenly by hand 15 minutes before sun exposure
- hold container 4 to 6 inches from the skin to apply
- do not spray directly into face. Spray on hands then apply to face.
- do not apply in windy conditions
- use in a well-ventilated area
- reapply:
 - after 80 minutes of swimming or sweating
 - immediately after towel drying
 - at least every 2 hours
- Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including:
 - limit time in the sun, especially from 10 a.m. – 2 p.m.
 - wear long-sleeve shirts, pants, hats, and sunglasses
 - children under 6 months: Ask a doctor

Other information

- protect this product from excessive heat and direct sun
- store between 20° to 25°C (68° to 77°F)
- may stain or damage some fabrics or surfaces

Inactive ingredients

SD alcohol 40-B (73.9% v/v), isobutane, neopentyl glycol dipheptanoate, polyester-27, acrylates/octylacrylamide copolymer, dimethicone, fragrance, ascorbyl palmitate

Questions

1-866-288-3330

Coppertone® Sunscreen Spray

Sport

50

4-in-1 Performance

Stops up to 98% UV Rays

+ Resists Sweat + Heat + Water

Free of Oxybenzone, Octinoxate, PABA & Dyes

Water Resistant (80 Minutes)

Broad Spectrum SPF 50

NET WT 1.6 OZ (45 g)



50

4-IN-1 PERFORMANCE

Stops up to 98% UV Rays

+RESISTS

Sweat + Heat + Water

Free of Oxybenzone, Octinoxate, PABA, Phthalates & Dyes

Water Resistant (80 Minutes)
Broad Spectrum SPF 50

NET WT 1.6 OZ (45 g)

Beiersdorf Beiersdorf Inc. Stamford, CT 06901 ©=reg. TM. of Beiersdorf AG, Germany

www.coppertone.com

Drug Facts

Active ingredients	Purpose
(To Deliver) Avobenzone 3%, Homosalate 10%, Octisalate 5%, Octocrylene 5%	Sunscreen

Use ■ helps prevent sunburn

Warnings

For external use only

Flammable: Do not use near heat, flame, or while smoking

Do not use on damaged or broken skin

When using this product ■ keep away from face to avoid breathing it ■ keep out of eyes. Rinse with water to remove. ■ use only as directed. Intentional misuse by deliberately concentrating and inhaling the contents can be harmful or fatal. ■ do not puncture or incinerate. Contents under pressure. Do not store at temperatures above 120°F.

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Directions ■ shake well before each use ■ spray liberally and spread evenly by hand 15 minutes before sun exposure ■ hold container 4 to 6 inches from the skin to apply ■ do not spray directly into face. Spray on hands then apply to face. ■ do not apply in windy conditions ■ use in a well-ventilated area ■ reapply: ■ after 80 minutes of swimming or sweating ■ immediately after towel drying ■ at least every 2 hours ■ **Sun Protection Measures.** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including: ■ limit time in the sun, especially from 10 a.m. – 2 p.m. ■ wear long-sleeve shirts, pants, hats, and sunglasses ■ children under 6 months: Ask a doctor

Other information ■ protect this product from excessive heat and direct sun ■ store between 20° to 25°C (68° to 77°F) ■ may stain or damage some fabrics or surfaces

Inactive ingredients SD alcohol 40-B (73.9% v/v), isobutane, neopentyl glycol dipheptanoate, polyester-27, acrylates/octylacrylamide copolymer, dimethicone, fragrance, ascorbyl palmitate

Questions? 1-866-288-3330



20006615
DOT-2P M5655



COPPERTONE SPORT SUNSCREEN SPF 50

avobenzone 3%, homosalate 10%, octisalate 5%, octocrylene 5% aerosol, spray

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:66800-4356
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
HOMOSALATE (UNII: V06SV4M95S) (HOMOSALATE - UNII:V06SV4M95S)	HOMOSALATE	10 g in 100 g
OCTOCRYLENE (UNII: 5A68WGF6WM) (OCTOCRYLENE - UNII:5A68WGF6WM)	OCTOCRYLENE	5 g in 100 g
OCTISALATE (UNII: 4X49Y0596W) (OCTISALATE - UNII:4X49Y0596W)	OCTISALATE	5 g in 100 g
AVOBENZONE (UNII: G63QQF2NOX) (AVOBENZONE - UNII:G63QQF2NOX)	AVOBENZONE	3 g in 100 g

Inactive Ingredients

Ingredient Name	Strength
ALCOHOL (UNII: 3K9958V90M)	
NEOPENTYL GLYCOL DIHEPTANOATE (UNII: 5LKW3C543X)	
ASCORBYL PALMITATE (UNII: QN83US2B0N)	
ISOBUTANE (UNII: BXR49TP611)	
POLYESTER-7 (UNII: 0841698D2F)	
ACRYLATE/ISOBUTYL METHACRYLATE/N-TERT-OCTYLACRYLAMIDE COPOLYMER (75000 MW) (UNII: JU3XHR8VWK)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
FRAGRANCE FLORAL ORC0902236 (UNII: R66Z4YW3X0)	

Product Characteristics

Color	yellow (yellow to pale yellow)	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66800-4356-1	45 g in 1 CAN; Type 0: Not a Combination Product	11/03/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part352	11/03/2020	

Labeler - Beiersdorf Inc (001177906)

