

**DOXYLAMINE SUCCINATE AND PYRIDOXINE HYDROCHLORIDE- doxylamine succinate and pyridoxine hydrochloride tablet, delayed release
Bryant Ranch Prepack**

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use DOXYLAMINE SUCCINATE AND PYRIDOXINE HYDROCHLORIDE DELAYED-RELEASE TABLETS safely and effectively. See full prescribing information for DOXYLAMINE SUCCINATE AND PYRIDOXINE HYDROCHLORIDE DELAYED-RELEASE TABLETS.

Doxylamine succinate and pyridoxine hydrochloride delayed-release tablets, for oral use.

Initial U.S. Approval: 1976

----- **RECENT MAJOR CHANGES** -----

Warnings and Precautions, Interference with Urine Screen for Methadone, Opiates and Phencyclidine Phosphate (PCP) (5.3) 05/2018

----- **INDICATIONS AND USAGE** -----

Doxylamine succinate and pyridoxine hydrochloride delayed-release tablets are a fixed dose combination drug product of doxylamine succinate, an antihistamine, and pyridoxine hydrochloride, a Vitamin B6 analog, indicated for the treatment of nausea and vomiting of pregnancy in women who do not respond to conservative management. (1)

----- **DOSAGE AND ADMINISTRATION** -----

Take two tablets daily at bedtime. If symptoms are not adequately controlled, the dose can be increased to a maximum recommended dose of four tablets daily (one in the morning, one mid-afternoon and two at bedtime) as described in the full prescribing information. (2)

----- **DOSAGE FORMS AND STRENGTHS** -----

Delayed-release tablets containing 10 mg doxylamine succinate and 10 mg pyridoxine hydrochloride. (3)

----- **CONTRAINDICATIONS** -----

- Known hypersensitivity to doxylamine succinate, other ethanolamine derivative antihistamines, pyridoxine hydrochloride or any inactive ingredient in the formulation (4)
- Monoamine oxidase (MAO) inhibitors (4, 7)

----- **WARNINGS AND PRECAUTIONS** -----

- Activities requiring mental alertness: Avoid engaging in activities requiring complete mental alertness, such as driving or operating heavy machinery, while using doxylamine succinate and pyridoxine hydrochloride delayed-release tablets until cleared to do so by a healthcare provider (5.1)
- Central nervous system (CNS) depressants: Concurrent use with alcohol or other CNS depressants is not recommended (5.1)
- Anticholinergic actions: Use with caution in patients with asthma, increased intraocular pressure, narrow angle glaucoma, stenosing peptic ulcer, pyloroduodenal obstruction and urinary bladder-neck obstruction (5.2)
- Interference with urine drug screen: Doxylamine succinate and pyridoxine hydrochloride delayed-release tablets may interfere with urine screening for methadone, opiates and PCP (5.3)

----- **ADVERSE REACTIONS** -----

The most common adverse reaction with doxylamine succinate and pyridoxine hydrochloride delayed-release tablets (≥ 5 percent and exceeding the rate in placebo) is somnolence. (6)

To report SUSPECTED ADVERSE REACTIONS, contact Analog Pharma Inc. at 1-844-884-5505 or medicalinfo@analogpharma.com or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

----- **DRUG INTERACTIONS** -----

- Severe drowsiness can occur when used in combination with alcohol or other sedating medications. (7)

----- **USE IN SPECIFIC POPULATIONS** -----

Doxylamine succinate and pyridoxine hydrochloride delayed-release tablets are intended for use in pregnant women. (8.1)

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 12/2020

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

2.1 Dosage Information

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

5.1 Activities Requiring Mental Alertness

5.2 Concomitant Medical Conditions

5.3 Interference with Urine Screen for Methadone, Opiates and Phencyclidine Phosphate (PCP)

6 ADVERSE REACTIONS

6.1 Clinical Trial Experience

6.2 Postmarketing Experience

7 DRUG INTERACTIONS

7.1 Drug Interactions

7.2 Drug-Food Interactions

7.3 False Positive Urine Tests for Methadone, Opiates and PCP

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

8.2 Lactation

8.4 Pediatric Use

10 OVERDOSAGE

10.1 Signs and Symptoms of Overdose

10.2 Management of Overdose

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis and Impairment of Fertility

14 CLINICAL STUDIES

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

Somnolence and Severe Drowsiness

Interference with urine drug screening

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

Doxylamine succinate and pyridoxine hydrochloride delayed-release tablets are indicated for the treatment of nausea and vomiting of pregnancy in women who do not respond to conservative management.

Limitations of Use

Doxylamine succinate and pyridoxine hydrochloride delayed-release tablets have not been studied in women with hyperemesis gravidarum.

2 DOSAGE AND ADMINISTRATION

2.1 Dosage Information

Initially, take two doxylamine succinate and pyridoxine hydrochloride delayed-release tablets orally at bedtime (Day 1). If this dose adequately controls symptoms the next day, continue taking two tablets daily at bedtime. However, if symptoms persist into the afternoon of Day 2, take the usual dose of two tablets at bedtime that night then take three tablets starting on Day 3 (one tablet in the morning and two tablets at bedtime). If these three tablets adequately control symptoms on Day 4, continue taking three tablets daily. Otherwise take four tablets starting on Day 4 (one tablet in the morning, one tablet mid-afternoon and two tablets at bedtime).

The maximum recommended dose is four tablets (one in the morning, one in the mid-afternoon and two at bedtime) daily.

Take on an empty stomach with a glass of water [see *Clinical Pharmacology (12.3)*]. Swallow tablets whole. Do not crush, chew, or split doxylamine succinate and pyridoxine hydrochloride delayed-release tablets.

Take as a daily prescription and not on an as needed basis. Reassess the woman for continued need for doxylamine succinate and pyridoxine hydrochloride delayed-release tablets as her pregnancy progresses.

3 DOSAGE FORMS AND STRENGTHS

Doxylamine succinate and pyridoxine hydrochloride delayed-release tablets are white, round, film coated tablets containing 10 mg doxylamine succinate and 10 mg pyridoxine hydrochloride. The tablets are imprinted with the pink image of a pregnant woman on one side.

4 CONTRAINDICATIONS

Doxylamine succinate and pyridoxine hydrochloride delayed-release tablets are contraindicated in women with any of the following conditions:

- Known hypersensitivity to doxylamine succinate, other ethanolamine derivative antihistamines, pyridoxine hydrochloride or any inactive ingredient in the formulation
- Monoamine oxidase (MAO) inhibitors intensify and prolong the adverse central nervous system effects of doxylamine succinate and pyridoxine hydrochloride delayed-release tablets [see *Drug Interactions (7.1)*].

5 WARNINGS AND PRECAUTIONS

5.1 Activities Requiring Mental Alertness

Doxylamine succinate and pyridoxine hydrochloride delayed-release tablets may cause somnolence due to the anticholinergic properties of doxylamine succinate, an antihistamine. Women should avoid engaging in activities requiring complete mental alertness, such as driving or operating heavy machinery, while using doxylamine succinate and pyridoxine hydrochloride delayed-release tablets until cleared to do so by their healthcare provider.

Doxylamine succinate and pyridoxine hydrochloride delayed-release tablets use is not recommended if a woman is concurrently using central nervous system (CNS) depressants including alcohol. The combination may result in severe drowsiness leading to falls or accidents [see *Drug Interactions (7.1)*].

5.2 Concomitant Medical Conditions

Doxylamine succinate and pyridoxine hydrochloride delayed-release tablets have anticholinergic properties and, therefore, should be used with caution in women with: asthma, increased intraocular

pressure, narrow angle glaucoma, stenosing peptic ulcer, pyloroduodenal obstruction and urinary bladder-neck obstruction.

5.3 Interference with Urine Screen for Methadone, Opiates and Phencyclidine Phosphate (PCP)

There have been reports of false positive urine screening tests for methadone, opiates, and PCP with doxylamine succinate/pyridoxine hydrochloride use [see *Drug Interactions (7.3)*].

6 ADVERSE REACTIONS

The following adverse reactions are discussed elsewhere in the labeling:

- Somnolence [see *Warnings and Precautions (5.1)*]
- Falls or other accidents resulting from the effect of the combined use of doxylamine succinate and pyridoxine hydrochloride delayed-release tablets with CNS depressants including alcohol [see *Warnings and Precautions (5.1)*]

6.1 Clinical Trial Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

The safety and efficacy of doxylamine succinate and pyridoxine hydrochloride delayed-release tablets were compared to placebo in a double-blind, randomized, multi-center trial in 261 women with nausea and vomiting of pregnancy. The mean gestational age at enrollment was 9.3 weeks, range 7 to 14 weeks gestation [see *Clinical Studies (14)*]. Adverse reactions for doxylamine succinate and pyridoxine hydrochloride delayed-release tablets that occurred at an incidence \geq 5 percent and exceeded the incidence for placebo are summarized in Table 1.

Table 1: Number (Percent) of Subjects with \geq 5 Percent Adverse Reactions in a 15-Day Placebo-Controlled Study of Doxylamine Succinate and Pyridoxine Hydrochloride Delayed-Release Tablets (Only Those Adverse Reactions Occurring at an Incidence \geq 5 Percent and at a Higher Incidence with Doxylamine Succinate and Pyridoxine Hydrochloride Delayed-Release Tablets than Placebo are Shown)

	Doxylamine succinate and pyridoxine hydrochloride delayed-release tablets (N = 133)	Placebo (n = 128)
Somnolence	19 (14.3%)	15 (11.7%)

6.2 Postmarketing Experience

The following adverse events, listed alphabetically, have been identified during post-approval use of the combination of 10 mg doxylamine succinate and 10 mg pyridoxine hydrochloride. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Cardiac disorders: dyspnea, palpitation, tachycardia

Ear and labyrinth disorders: vertigo

Eye disorders: vision blurred, visual disturbances

Gastrointestinal disorders: abdominal distension, abdominal pain, constipation, diarrhea

General disorders and administration site conditions: chest discomfort, fatigue, irritability, malaise

Immune system disorders: hypersensitivity

Nervous system disorders: dizziness, headache, migraines, paresthesia, psychomotor hyperactivity

Psychiatric disorders: anxiety, disorientation, insomnia, nightmares

Renal and urinary disorders: dysuria, urinary retention

Skin and subcutaneous tissue disorders: hyperhidrosis, pruritus, rash, rash maculo-papular

7 DRUG INTERACTIONS

7.1 Drug Interactions

Use of doxylamine succinate and pyridoxine hydrochloride delayed-release tablets is contraindicated in women who are taking monoamine oxidase inhibitors (MAOIs), which prolong and intensify the anticholinergic (drying) effects of antihistamines. Concurrent use of alcohol and other CNS depressants (such as hypnotic sedatives and tranquilizers) with doxylamine succinate and pyridoxine hydrochloride delayed-release tablets is not recommended.

7.2 Drug-Food Interactions

A food-effect study demonstrated that the delay in the onset of action of doxylamine succinate and pyridoxine hydrochloride delayed-release tablets may be further delayed, and a reduction in absorption may occur when tablets are taken with food [*see Dosage and Administration (2), Clinical Pharmacology (12.3)*]. Therefore, doxylamine succinate and pyridoxine hydrochloride delayed-release tablets should be taken on an empty stomach with a glass of water [*see Dosage and Administration (2)*].

7.3 False Positive Urine Tests for Methadone, Opiates and PCP

False positive drug screens for methadone, opiates, and PCP can occur with doxylamine succinate/pyridoxine hydrochloride use. Confirmatory tests, such as Gas Chromatography Mass Spectrometry (GC-MS), should be used to confirm the identity of the substance in the event of a positive immunoassay result.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

Doxylamine succinate and pyridoxine hydrochloride delayed-release tablets are intended for the treatment of nausea and vomiting of pregnancy in women who do not respond to conservative management. Maternal risks are discussed throughout the labeling. No increased risk for congenital malformations has been reported in epidemiologic studies in pregnant women.

In the U.S. general population, the estimated background risks for major birth defects and miscarriage in clinically recognized pregnancies are 2-4% and 15-20%, respectively.

Data

Human Data

The combination of doxylamine succinate and pyridoxine hydrochloride has been the subject of many epidemiological studies (cohort, case control and meta-analyses) designed to detect possible teratogenicity. A meta-analysis of 16 cohort and 11 case-control studies published between 1963 and

1991 reported no increased risk for malformations from first trimester exposures to doxylamine succinate and pyridoxine hydrochloride, with or without dicyclomine hydrochloride. A second meta-analysis of 12 cohort and 5 case-control studies published between 1963 and 1985 reported no statistically significant relationships between fetal abnormalities and the first trimester use of the combination doxylamine succinate and pyridoxine hydrochloride with or without dicyclomine hydrochloride.

8.2 Lactation

□Risk Summary

Women should not breastfeed while using doxylamine succinate and pyridoxine hydrochloride delayed-release tablets.

The molecular weight of doxylamine succinate is low enough that passage into breast milk can be expected. Excitement, irritability and sedation have been reported in nursing infants presumably exposed to doxylamine succinate through breast milk. Infants with apnea or other respiratory syndromes may be particularly vulnerable to the sedative effects of doxylamine succinate and pyridoxine hydrochloride delayed-release tablets resulting in worsening of their apnea or respiratory conditions.

Pyridoxine hydrochloride is excreted into breast milk. There have been no reports of adverse events in infants presumably exposed to pyridoxine hydrochloride through breast milk.

8.4 Pediatric Use

The safety and effectiveness of doxylamine succinate and pyridoxine hydrochloride delayed-release tablets in children under 18 years of age have not been established.

Fatalities have been reported from doxylamine overdose in children. The overdose cases have been characterized by coma, grand mal seizures and cardiorespiratory arrest. Children appear to be at a high risk for cardiorespiratory arrest. A toxic dose for children of more than 1.8 mg/kg has been reported. A 3 year old child died 18 hours after ingesting 1,000 mg doxylamine succinate. However, there is no correlation between the amount of doxylamine ingested, the doxylamine plasma level and clinical symptomatology.

10 OVERDOSAGE

10.1 Signs and Symptoms of Overdose

Doxylamine succinate and pyridoxine hydrochloride delayed-release tablets are a delayed-release formulation, therefore, signs and symptoms of intoxication may not be apparent immediately.

Signs and symptoms of overdose may include restlessness, dryness of mouth, dilated pupils, sleepiness, vertigo, mental confusion and tachycardia.

At toxic doses, doxylamine exhibits anticholinergic effects, including seizures, rhabdomyolysis, acute renal failure and death.

10.2 Management of Overdose

If treatment is needed, it consists of gastric lavage or activated charcoal, whole bowel irrigation and symptomatic treatment. For additional information about overdose treatment, call a poison control center (1-800-222-1222).

11 DESCRIPTION


Doxylamine succinate and pyridoxine hydrochloride delayed-release tablets are round, white, film-coated, delayed-release tablets containing 10 mg of doxylamine succinate and 10 mg of pyridoxine

hydrochloride. Tablets are imprinted on one side with the pink image of a pregnant woman.

Inactive ingredients are as follows: ammonium hydroxide, n-butanol, carnauba wax powder, colloidal silicon dioxide, croscarmellose sodium, D&C Red#27, denatured alcohol, FD&C Blue#2, hypromellose, isopropyl alcohol, magnesium stearate, magnesium trisilicate, methacrylic acid copolymer, microcrystalline cellulose 102, PEG 400, PEG 8000, polysorbate 80, propylene glycol, shellac glaze, simethicone, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triethyl citrate.

Doxylamine Succinate


Doxylamine succinate is classified as an antihistamine. The chemical name for doxylamine succinate is ethanamine, N,N-dimethyl-2-[1-phenyl-1-(2-pyridinyl)ethoxy]-, butanedioate (1:1). The empirical formula is $C_{17}H_{22}N_2O \cdot C_4H_6O_4$ and the molecular mass is 388.46. The structural formula is:



Doxylamine succinate is a white to creamy white powder that is very soluble in water and alcohol, freely soluble in chloroform and very slightly soluble in ether and benzene.

Pyridoxine Hydrochloride

Pyridoxine hydrochloride is a vitamin B6 analog. The chemical name for pyridoxine hydrochloride is 3,4-pyridinedimethanol, 5-hydroxy-6-methyl-, hydrochloride. The empirical formula is $C_8H_{11}NO_3 \cdot HCl$ and the molecular mass is 205.64. The structural formula is:



Pyridoxine hydrochloride is a white or practically white crystalline powder that is freely soluble in water, slightly soluble in alcohol and insoluble in ether.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

The mechanism of action of doxylamine succinate and pyridoxine hydrochloride delayed-release tablets is unknown.

12.3 Pharmacokinetics

The pharmacokinetics of doxylamine succinate and pyridoxine hydrochloride delayed-release tablets has been characterized in healthy non-pregnant adult women. Pharmacokinetic results for doxylamine and pyridoxine, including its vitamin B6 metabolites, pyridoxal, pyridoxal 5'-phosphate, pyridoxamine and pyridoxamine 5'-phosphate, are summarized in Tables 2 to 5.

Absorption

A single-dose (two tablets) and multiple-dose (four tablets daily), open-label study was conducted to assess the safety and pharmacokinetic profile of doxylamine succinate and pyridoxine hydrochloride delayed-release tablets administered in healthy non-pregnant adult women. Single-doses (two tablets at bedtime) were administered on Days 1 and 2. Multiple-doses (one tablet in the morning, one tablet in the afternoon and two tablets at bedtime) were administered on Days 3-18.

Blood samples for pharmacokinetic analysis were collected pre-and post-dose on Days 2 and 18 as well as pre-dose prior to bedtime dose only (trough) on Days 9, 10, 11, 16, 17, and 18.

Doxylamine and pyridoxine are absorbed in the gastrointestinal tract, mainly in the jejunum.

The C_{max} of doxylamine and pyridoxine are achieved within 7.5 and 5.5 hours, respectively (see Table 2).

Table 2 – Single-Dose and Multiple-Dose Pharmacokinetics of Doxylamine Succinate and Pyridoxine Hydrochloride Delayed-Release Tablets in Healthy Non-Pregnant Adult Women

	Single Dose			Multiple Dose		
	AUC _{0-inf} (ng•h/mL)	C _{max} (ng/mL)	T _{max} (h)	AUC _{0-inf} (ng•h/mL)	C _{max} (ng/mL)	T _{max} (h)
Doxylamine	1280.9 ± 369.3	83.3 ± 20.6	7.2 ± 1.9	3721.5 ± 1318.5	168.6 ± 38.5	7.8 ± 1.6
Pyridoxine	43.4 ± 16.5	32.6 ± 15.0	5.7 ± 1.5	64.5 ± 36.4	46.1 ± 28.3	5.6 ± 1.3
Pyridoxal	211.6 ± 46.1	74.3 ± 21.8	6.5 ± 1.4	1587.2 ± 550.0	210.0 ± 54.4	6.8 ± 1.2
Pyridoxal 5`Phosphate	1536.4 ± 721.5	30.0 ± 10.0	11.7 ± 5.3	6099.7 ± 1383.7	84.9 ± 16.9	6.3 ± 6.6
Pyridoxamine	4.1 ± 2.7	0.5 ± 0.7	5.9 ± 2.1	2.6 ± 0.8	0.5 ± 0.2	6.6 ± 1.4
Pyridoxamine 5'-phosphate	5.2 ± 3.8	0.7 ± 0.5	14.8 ± 6.6	94.5 ± 58.0	2.3 ± 1.7	12.4 ± 11.2

Multiple-dose administration of doxylamine succinate and pyridoxine hydrochloride delayed-release tablets results in increased concentrations of doxylamine as well as increases in doxylamine C_{max} and AUC_{0-last} of absorption. The time to reach the maximum concentration is not affected by multiple doses. The mean accumulation index is more than 1.0 suggesting that doxylamine accumulates following multiple dosing (see Table 3).

Although no accumulation was observed for pyridoxine, the mean accumulation index for each metabolite (pyridoxal, pyridoxal 5'-phosphate, and pyridoxamine 5'-phosphate) is more than 1.0 following multiple-dose administration of doxylamine succinate and pyridoxine hydrochloride delayed-release tablets. The time to reach the maximum concentration is not affected by multiple doses (see Table 2).

Table 3 – Pharmacokinetics of Doxylamine and Pyridoxine Following Single Dose and Multiple Dose Administration of Doxylamine Succinate and Pyridoxine Hydrochloride Delayed-Release Tablets to Healthy Non-Pregnant Adult Women

		AUC_{0-last} (ng•h/mL)	AUC_{0-inf} (ng•h/mL)	C_{max} (ng/mL)	T_{max} (h)	T_{1/2el} (h)
Doxylamine Mean±SD N=18	Single	911.4 ± 205.6	1280.9 ± 369.3	83.3 ± 20.6	7.2 ± 1.9	10.1 ± 2.1
	Multiple	3661.3 ± 1279.2	3721.5 ± 1318.5	168.6 ± 38.5	7.8 ± 1.6	11.9 ± 3.3
Pyridoxine Mean±SD N=18	Single	39.3 ± 16.5	43.4 ± 16.5	32.6 ± 15.0	5.7 ± 1.5	0.5 ± 0.2
	Multiple	59.3 ± 33.9	64.5 ± 36.4	46.1 ± 28.3	5.6 ± 1.3	0.5 ± 0.1

Food Effect

The administration of food delays the absorption of both doxylamine and pyridoxine. This delay is associated with a lower peak concentration of doxylamine, but the extent of absorption is not affected (see Table 4).

The effect of food on the peak concentration and the extent of absorption of the pyridoxine component is more complex because the pyridoxal, pyridoxamine, pyridoxal 5'-phosphate and pyridoxamine 5'-phosphate metabolites also contribute to the biological activity. Food significantly reduces the bioavailability of pyridoxine, lowering its C_{max} and AUC by approximately 50% compared to fasting conditions. Similarly, food significantly reduces pyridoxal AUC and reduces its C_{max} by 50% compared to fasting conditions. In contrast, food slightly increases pyridoxal 5'-phosphate C_{max} and extent of absorption. As for pyridoxamine and pyridoxamine 5'-phosphate, the rate and extent of absorption seem to decrease under fed conditions.

Table 4 – Pharmacokinetics of Doxylamine and Pyridoxine Following Administration of Doxylamine Succinate and Pyridoxine Hydrochloride Delayed-Release Tablets Under Fed and Fasted Conditions in Healthy Non-Pregnant Adult Women

		AUC_{0-t} (ng•h/mL)	AUC_{0-inf} (ng•h/mL)	C_{max} (ng/mL)	T_{max} (h)	T_{1/2el} (h)
Doxylamine Mean±SD N=42	Fasted	1407.2 ± 336.9	1447.9 ± 332.2	94.9 ± 18.4	5.1 ± 3.4	12.6 ± 3.4
	Fed	1488.0 ± 463.2	1579.0 ± 422.7*	75.7 ± 16.6	14.9 ± 7.4	12.5 ± 2.9*
Pyridoxine Mean±SD N=42	Fasted	33.8 ± 13.7	39.5 ± 12.9†	35.5 ± 21.4	2.5 ± 0.9	0.4 ± 0.2†
	Fed	18.3 ± 14.5	24.2 ±14.0‡	13.7 ± 10.8	9.3 ± 4.0	0.5 ± 0.2‡

* N=37

† N=31

‡ N=18

Distribution

Pyridoxine is highly protein bound, primarily to albumin. Its main active metabolite, pyridoxal 5'-phosphate (PLP) accounts for at least 60% of circulating vitamin B6 concentrations.

Metabolism

Doxylamine is biotransformed in the liver by N-dealkylation to its principle metabolites N-desmethyl-doxylamine and N, N-didesmethyldoxylamine.

Pyridoxine is a prodrug primarily metabolized in the liver.

Excretion

The principle metabolites of doxylamine, N-desmethyl-doxylamine and N, N-didesmethyldoxylamine, are excreted by the kidney.

The terminal elimination half-life of doxylamine and pyridoxine are 12.5 hours and 0.5 hours, respectively (see Table 5).

Table 5 – Terminal Elimination Half-Life ($T_{1/2el}$) for Doxylamine Succinate and Pyridoxine Hydrochloride Delayed-Release Tablets Administered as a Single Dose of Two Tablets under Fasting Conditions in Healthy Non-Pregnant Adult Women

	$T_{1/2el}$ (h)
Doxylamine	12.6 ± 3.4
Pyridoxine	0.4 ± 0.2
Pyridoxal	2.1 ± 2.2
Pyridoxal 5'-Phosphate	81.6 ± 42.2
Pyridoxamine	3.1 ± 2.5
Pyridoxamine 5'-Phosphate	66.5 ± 51.3

Use in Specific Populations

Race: No pharmacokinetic studies have been conducted related to race.

Hepatic Impairment: No pharmacokinetic studies have been conducted in hepatic impaired patients.

Renal Impairment: No pharmacokinetic studies have been conducted in renal impaired patients.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis and Impairment of Fertility

Carcinogenicity

Two-year carcinogenicity studies in rats and mice have been conducted with doxylamine succinate. Doxylamine succinate is not likely to have human carcinogenic potential. The carcinogenic potential of pyridoxine hydrochloride has not been evaluated.

14 CLINICAL STUDIES

A double-blind, randomized, multi-center, placebo-controlled study was conducted to support the safety and efficacy of doxylamine succinate and pyridoxine hydrochloride delayed-release tablets in the treatment of nausea and vomiting of pregnancy. Adult women 18 years of age or older and 7 to 14 weeks gestation (median 9 weeks of gestation) with nausea and vomiting of pregnancy were randomized to 14 days of doxylamine succinate and pyridoxine hydrochloride delayed-release tablets or placebo. Two tablets of doxylamine succinate and pyridoxine hydrochloride delayed-release tablets were

administered at bedtime on Day 1. If symptoms of nausea and vomiting persisted into the afternoon hours of Day 2, the woman was directed to take her usual dose of two tablets at bedtime that night and, beginning on Day 3, to take one tablet in the morning and two tablets at bedtime. Based upon assessment of remaining symptoms at her clinic visit on Day 4 (\pm 1 day), the woman may have been directed to take an additional tablet mid-afternoon. A maximum of four tablets (one in the morning, one in the mid-afternoon and two at bedtime) were taken daily.

Over the treatment period, 19% of doxylamine succinate and pyridoxine hydrochloride delayed-release tablets -treated patients remained on 2 tablets daily, 21% received 3 tablets daily, and 60% received 4 tablets daily.

The primary efficacy endpoint was the change from baseline at Day 15 in the Pregnancy Unique-Quantification of Emesis (PUQE) score. The PUQE score incorporates the number of daily vomiting episodes, number of daily heaves, and length of daily nausea in hours, for an overall score of symptoms rated from 3 (no symptoms) to 15 (most severe).

At baseline, the mean PUQE score was 9.0 in the doxylamine succinate and pyridoxine hydrochloride delayed-release tablets arm and 8.8 in the placebo arm. There was a 0.7 (95% confidence interval 0.2 to 1.2 with p-value 0.006) mean decrease (improvement in nausea and vomiting symptoms) from baseline in PUQE score at Day 15 with doxylamine succinate and pyridoxine hydrochloride delayed-release tablets compared to placebo (see Table 6).

Table 6 – Change from Baseline in the Primary Endpoint, Pregnancy Unique-Quantification of Emesis (PUQE) Score at Day 15. (Intent-to-Treat Population with Last-Observation Carried Forward)

PUQE Score*	Doxylamine Succinate + Pyridoxine Hydrochloride	Placebo	Treatment Difference [95% Confidence Interval]
Baseline	9.0 \pm 2.1	8.8 \pm 2.1	
Change from baseline at Day 15	-4.8 \pm 2.7	-3.9 \pm 2.6	-0.7 [-1.2, -0.2]

* The Pregnancy-Unique Quantification of Emesis and Nausea (PUQE) score incorporated the number of daily vomiting episodes, number of daily heaves, and length of daily nausea in hours, for an overall score of symptoms rated from 3 (no symptoms) to 15 (most severe). Baseline was defined as the PUQE score completed at the enrollment visit.

16 HOW SUPPLIED/STORAGE AND HANDLING

NDC: 63629-8358-1 30 Tablets in a BOTTLE

17 PATIENT COUNSELING INFORMATION

See FDA-approved patient labeling (Patient Information)

Somnolence and Severe Drowsiness

Inform women to avoid engaging in activities requiring complete mental alertness, such as driving or operating heavy machinery, while using doxylamine succinate and pyridoxine hydrochloride delayed-

release tablets until cleared to do so.

Inform women of the importance of not taking doxylamine succinate and pyridoxine hydrochloride delayed-release tablets with alcohol or sedating medications, including other antihistamines (present in some cough and cold medications), opiates and sleep aids because somnolence could worsen leading to falls or other accidents.

Interference with urine drug screening

Inform women that use of doxylamine succinate and pyridoxine hydrochloride delayed-release tablets may result in false positive urine drug screening for methadone, opiates and PCP.

Distributed by:

Analog Pharma Inc.

Bryn Mawr, PA, 19010

www.analogpharma.com

Patient Package Insert

Patient Information
<p style="text-align: center;">Doxylamine succinate and pyridoxine hydrochloride delayed-release tablets</p> <p>What are doxylamine succinate and pyridoxine hydrochloride delayed-release tablets?</p> <ul style="list-style-type: none">• Doxylamine succinate and pyridoxine hydrochloride delayed-release tablets are a prescription medicine used to treat nausea and vomiting of pregnancy in women who have not improved with change in diet or other non-medicine treatments.• It is not known if doxylamine succinate and pyridoxine hydrochloride delayed-release tablets are safe and effective in women with severe nausea and vomiting of pregnancy, a condition called hyperemesis gravidarum. Women with this condition may need to be hospitalized.• It is not known if doxylamine succinate and pyridoxine hydrochloride delayed-release tablets are safe and effective in children under 18 years of age.
<p>Who should not take doxylamine succinate and pyridoxine hydrochloride delayed-release tablets?</p> <p>Do not take doxylamine succinate and pyridoxine hydrochloride delayed-release tablets if you:</p> <ul style="list-style-type: none">• are allergic to doxylamine succinate, other ethanolamine derivative antihistamines, pyridoxine hydrochloride or any of the ingredients in doxylamine succinate and pyridoxine hydrochloride delayed-release tablets. See the end of this leaflet for a complete list of ingredients in doxylamine succinate and pyridoxine hydrochloride delayed-release tablets.• take monoamine oxidase inhibitors (MAOIs). Ask your healthcare provider or pharmacist if you are not sure if you take an MAOI, including Marplan, Nardil, Emsam, Eldepryl, Zelapar, and Parnate.
<p>Before taking doxylamine succinate and pyridoxine hydrochloride delayed-release tablets, tell your healthcare provider about all of your medical conditions, including if you:</p> <ul style="list-style-type: none">• have asthma.• have eye problems called increased intraocular pressure or narrow angle glaucoma.• have a stomach problem called stenosing peptic ulcer or pyloroduodenal obstruction.• have a bladder problem called urinary bladder-neck obstruction.• are breastfeeding or plan to breastfeed. Doxylamine succinate and pyridoxine hydrochloride delayed-release tablets can pass into your breast milk and may harm your baby. You should not breastfeed while using doxylamine succinate and pyridoxine hydrochloride delayed-release tablets.
<p>Tell your healthcare provider about all the medicines you take, including prescription and over-the-</p>

counter medicines, vitamins, and herbal supplements.

How should I take doxylamine succinate and pyridoxine hydrochloride delayed-release tablets?

- Talk to your healthcare provider about how much doxylamine succinate and pyridoxine hydrochloride delayed-release tablets to take and when to take it.
- Take doxylamine succinate and pyridoxine hydrochloride delayed-release tablets everyday as prescribed by your healthcare provider. Do not stop taking doxylamine succinate and pyridoxine hydrochloride delayed-release tablets without talking to your healthcare provider first.
- **See the following schedule for the right way you should start taking doxylamine succinate and pyridoxine hydrochloride delayed-release tablets:**
 - Day 1- Take 2 tablets, by mouth at bedtime.
 - Day 2- Take 2 tablets at bedtime. If your nausea and vomiting is better or controlled on Day 2, continue to take 2 tablets every night at bedtime. This will be your usual dose unless your healthcare provider tells you otherwise.
 - Day 3- If you still had nausea and vomiting on Day 2, take 3 tablets on Day 3 (1 tablet in the morning and 2 tablets at bedtime).
 - Day 4- If your nausea and vomiting was better or controlled on Day 3, continue to take 3 tablets each day (1 tablet in the morning and 2 tablets at bedtime). If you still had nausea and vomiting on Day 3, start taking 4 tablets each day (1 tablet in the morning, 1 tablet in the afternoon, and 2 tablets at bedtime).
- Do not take more than 4 tablets (1 in the morning, 1 in the mid-afternoon, and 2 at bedtime) each day.
- Take doxylamine succinate and pyridoxine hydrochloride delayed-release tablets on an empty stomach with a glass of water.
- Take doxylamine succinate and pyridoxine hydrochloride delayed-release tablets whole. Do not crush, chew, or break doxylamine succinate and pyridoxine hydrochloride delayed-release tablets before swallowing. If you cannot swallow doxylamine succinate and pyridoxine hydrochloride delayed-release tablets whole, tell your healthcare provider.
- If you take too many doxylamine succinate and pyridoxine hydrochloride delayed-release tablets (overdose), you may have the following symptoms: restlessness, dry mouth, the pupils of your eyes become larger (dilated), sleepiness, dizziness, confusion, fast heart rate, seizures, muscle pain or weakness, and sudden and severe kidney problems. If you have these symptoms and they are severe, they may lead to death. Stop taking doxylamine succinate and pyridoxine hydrochloride delayed-release tablets, call your healthcare provider or go to the nearest hospital emergency room right away. For more information about overdose treatment, call your poison control center at 1-800-222-1222.

What are the possible side effects of doxylamine succinate and pyridoxine hydrochloride delayed-release tablets?

Doxylamine succinate and pyridoxine hydrochloride delayed-release tablets may cause serious side effects, including drowsiness.

Drowsiness is a common side effect when taking doxylamine succinate and pyridoxine hydrochloride delayed-release tablets, but can also be severe:

- **Do not** drive, operate heavy machinery, or other activities that need your full attention unless your healthcare provider says that you may do so.
- **Do not** drink alcohol, or take other central nervous system depressants such as cough and cold medicines, certain pain medicines, and medicines that help you sleep while you take doxylamine succinate and pyridoxine hydrochloride delayed-release tablets. Severe drowsiness can happen or become worse causing falls or accidents.

Doxylamine succinate and pyridoxine hydrochloride delayed-release tablets may cause false positive urine drug screening test for methadone, opiates and PCP.

These are not all the possible side effects of doxylamine succinate and pyridoxine hydrochloride delayed-release tablets.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store doxylamine succinate and pyridoxine hydrochloride delayed-release tablets?

- Store doxylamine succinate and pyridoxine hydrochloride delayed-release tablets between 68°F to 77°F (20°C to 25°C).
- Keep doxylamine succinate and pyridoxine hydrochloride delayed-release tablets dry, in a tightly closed container, and out of the light.
- Safely throw away medicine that is out of date or no longer needed.

Keep doxylamine succinate and pyridoxine hydrochloride delayed-release tablets and all medicines out of the reach of children.

General information about the safe and effective use of doxylamine succinate and pyridoxine hydrochloride delayed-release tablets.

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. You can ask your pharmacist or healthcare provider for information about doxylamine succinate and pyridoxine hydrochloride delayed-release tablets that is written for health professionals. Do not use doxylamine succinate and pyridoxine hydrochloride delayed-release tablets for a condition for which it was not prescribed. Do not give doxylamine succinate and pyridoxine hydrochloride delayed-release tablets to other people, even if they have the same symptoms that you have. It may harm them.

What are the ingredients in doxylamine succinate and pyridoxine hydrochloride delayed-release tablets?

Active ingredient: doxylamine succinate (an antihistamine) and pyridoxine hydrochloride (vitamin B₆).

Inactive ingredients: ammonium hydroxide, n-butanol, carnauba wax powder, colloidal silicon dioxide, croscarmellose sodium, D&C Red#27, denatured alcohol, FD&C Blue #2, hypromellose, isopropyl alcohol, magnesium stearate, magnesium trisilicate, methacrylic acid copolymer, microcrystalline cellulose 102, PEG 400, PEG 8000, polysorbate 80, propylene glycol, shellac glaze, simethicone, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triethyl citrate.

Distributed by: Analog Pharma Inc., Bryn Mawr, PA, 19010

www.analogpharma.com

This Patient Information has been approved by the U.S. Food and Drug Administration

Issued: October 2018

Doxylamine Succ. Pyridoxine Hcl 10/10m

Packaged by Bryant Ranch Prepack

Burbank, CA 91504

Doxylamine Succ. Pyridoxine Hcl 10/10m

WHITE ROUND LOGO

LOT 1523487

Store at room temp of
20°-25°C (68°-77°F)

Compare To

Diclegis 10/10mg Tablet

Analog Pharma

Keep all drugs out of
reach of children.

RX Only

30

EXP MM/YY

NDC

6362983581



08358301523487

DOXYLAMINE SUCCINATE AND PYRIDOXINE HYDROCHLORIDE

doxylamine succinate and pyridoxine hydrochloride tablet, delayed release

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:63629-8358(NDC:70505-100)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	10 mg
PYRIDOXINE HYDROCHLORIDE (UNII: 68Y4CF58BV) (PYRIDOXINE - UNII:KV2JZ1BI6Z)	PYRIDOXINE HYDROCHLORIDE	10 mg

Inactive Ingredients

Ingredient Name	Strength
AMMONIA (UNII: 5138Q19F1X)	
BUTYL ALCOHOL (UNII: 8PJ61P6TS3)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	

CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)
D&C RED NO. 27 (UNII: 2LRS185U6K)
ALCOHOL (UNII: 3K9958V90M)
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)
ISOPROPYL ALCOHOL (UNII: ND2M416302)
MAGNESIUM STEARATE (UNII: 70097M6I30)
MAGNESIUM TRISILICATE (UNII: C2E1CI501T)
METHACRYLIC ACID - ETHYL ACRYLATE COPOLYMER (1:1) TYPE A (UNII: NX76LV5T8J)
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)
POLYETHYLENE GLYCOL 8000 (UNII: Q662QK8M3B)
POLYSORBATE 80 (UNII: 6OZP39ZG8H)
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)
SHELLAC (UNII: 46N107B71O)
DIMETHICONE (UNII: 92RU3N3Y1O)
SODIUM BICARBONATE (UNII: 8MDF5V39QO)
SODIUM LAURYL SULFATE (UNII: 368GB5141J)
TALC (UNII: 7SEV7J4R1U)
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	8mm
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63629-8358-1	30 in 1 BOTTLE; Type 0: Not a Combination Product	12/17/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA authorized generic	NDA021876	11/15/2018	

Labeler - Bryant Ranch Prepack (171714327)

Registrant - Bryant Ranch Prepack (171714327)

Establishment

Name	Address	ID/FEI	Business Operations
Bryant Ranch Prepack		171714327	REPACK(63629-8358) , RELABEL(63629-8358)

