

POLYETHYLENE GLYCOL 3350- polyethylene glycol 3350 powder, for solution
Preferred Pharmaceuticals Inc.

Polyethylene Glycol 3350 Powder for Oral Solution, 17 g

ACTIVE INGREDIENT (IN EACH DOSE)

(Bottle Only)

Polyethylene Glycol 3350, 17 g (cap filled to line)

(Packet Only)

Polyethylene Glycol 3350, 17 g

PURPOSE

Osmotic Laxative

USE

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

WARNINGS

Allergy alert: Do not use if you are allergic to polyethylene glycol

DO NOT USE

if you have kidney disease, except under the advice and supervision of a doctor

ASK A DOCTOR BEFORE USE IF YOU HAVE

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

ASK A DOCTOR OR PHARMACIST BEFORE USE IF YOU ARE

taking a prescription drug

WHEN USING THIS PRODUCT

you may have loose, watery, more frequent stools

STOP USE AND ASK A DOCTOR IF

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

IF PREGNANT OR BREAST-FEEDING,

ask a health professional before use.

KEEP OUT OF REACH OF CHILDREN

In case of overdose, get medical help or contact a Poison Control Center right away.

DIRECTIONS

(Bottle Only)

- **do not take more than directed unless advised by your doctor**
- the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap)
- adults and children 17 years of age and older:
 - use once a day
 - fill to top of white section in cap which is marked to indicate the correct dose (17 g)
 - stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
- do not combine with starch-based thickeners used for difficult swallowing
- ensure that the powder is fully dissolved before drinking
- do not drink if there are any clumps
- do not use more than 7 days
- children 16 years of age or under: ask a doctor

(Packet Only)

- **do not take more than directed unless advised by your doctor**
- adults and children 17 years of age and older:
 - use once a day
 - stir and dissolve one packet of powder (17 g) in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
- do not combine with starch-based thickeners used for difficult swallowing
- ensure that the powder is fully dissolved before drinking
- do not drink if there are any clumps

Active Ingredient/Active Moiety

Ingredient Name		Basis of Strength	Strength	
POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P) (POLYETHYLENE GLYCOL 3350 - UNII:G2M7P15E5P)		POLYETHYLENE GLYCOL 3350	17 g in 17 g	
Product Characteristics				
Color	white	Score		
Shape		Size		
Flavor		Imprint Code		
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68788-8601-2	238 g in 1 BOTTLE; Type 0: Not a Combination Product	03/08/2024	
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA		ANDA214990	03/08/2024	

Labeler - Preferred Pharmaceuticals Inc. (791119022)

Registrant - Preferred Pharmaceuticals Inc. (791119022)

Establishment

Name	Address	ID/FEI	Business Operations
Preferred Pharmaceuticals Inc.		791119022	RELABEL(68788-8601)