

ACETAMINOPHEN- acetaminophen tablet extended release tablet, extended release
MUSCLE ACHES AND PAINS ACETAMINOPHEN EXTENDED RELEASE-
acetaminophen tablet, extended release
TIME CAP LABORATORIES, INC.

Timely 699R 704R Acetaminophen Extended-Release Tablets USP, 650 mg

Active ingredient (in each caplet)

Acetaminophen 650 mg

Purpose

Pain reliever/fever reducer

Uses

For Arthritis Pain

temporarily relieves minor aches and pains due to:

- minor pain of arthritis
- muscular aches
- backache
- premenstrual and menstrual cramps
- the common cold
- headache
- toothache

temporarily reduces fever

For Muscle Aches & Pains

temporarily relieves minor aches and pains due to:

- muscular aches
- backache
- minor pain of arthritis
- toothache
- premenstrual and menstrual cramps
- headache
- the common cold

temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product.

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may

include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if you have liver disease.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Stop using and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning:

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

For Arthritis Pain

Do not take more than directed. See overdose warning

adults:

- take 2 caplets every 8 hours with water
- swallow whole; do not crush, chew, split or dissolve
- do not take more than 6 caplets in 24 hours
- do not use for more than 10 days unless directed by a doctor

under 18 years of age: ask a doctor

For Muscle Aches & Pains

Do not take more than directed. See overdose warning

adults and children 12 years and over:

- take 2 caplets every 8 hours with water
- swallow whole; do not crush, chew, split or dissolve

- do not take more than 6 caplets in 24 hours
- do not use for more than 10 days unless directed by a doctor

children under 12 years: do not use

Other information

- store between 20-25°C (68-77°F)
- The FDA approved Dissolution methods differ from USP

Inactive ingredients carnauba wax, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, povidone, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin

Questions or comments?

Call **1-877-290-4008**

<p>Drug Facts (continued)</p> <p>Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ hives ■ rash. If a skin reaction occurs, stop use and seek medical help right away.</p> <p>Do not use</p> <ul style="list-style-type: none"> ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product. <p>Ask a doctor before use if you have liver disease</p> <p>Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present <p>These could be signs of a serious condition.</p> <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children.</p> <p>Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p>Directions Do not take more than directed. See overdose warning.</p> <table border="1"> <tr> <td>adults</td> <td> <ul style="list-style-type: none"> ■ take 2 caplets every 8 hours with water or dissolve ■ swallow whole; do not crush, chew, split, or do not take more than 6 caplets in 24 hours ■ do not use for more than 10 days unless directed by a doctor </td> </tr> <tr> <td>under 18 years of age</td> <td> <ul style="list-style-type: none"> ■ ask a doctor </td> </tr> </table>		adults	<ul style="list-style-type: none"> ■ take 2 caplets every 8 hours with water or dissolve ■ swallow whole; do not crush, chew, split, or do not take more than 6 caplets in 24 hours ■ do not use for more than 10 days unless directed by a doctor 	under 18 years of age	<ul style="list-style-type: none"> ■ ask a doctor
adults	<ul style="list-style-type: none"> ■ take 2 caplets every 8 hours with water or dissolve ■ swallow whole; do not crush, chew, split, or do not take more than 6 caplets in 24 hours ■ do not use for more than 10 days unless directed by a doctor 				
under 18 years of age	<ul style="list-style-type: none"> ■ ask a doctor 				
<p>Other information</p> <ul style="list-style-type: none"> ■ store between 20-25°C (68-77°F) ■ The FDA approved dissolution methods differ from USP <p>Inactive ingredients carnauba wax, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, povidone, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin</p> <p>Questions or comments? Call 1-877-290-4008</p> <p>Contains No Aspirin</p>					

Inside

<p>NDC 49483-699-01</p> <p>*Compare to Tylenol® 8hr Arthritis Pain active ingredient</p> <p>timely™</p> <p>Arthritis Pain Relief</p> <p>Acetaminophen</p> <p>Extended-Release Tablets USP, 650 mg</p> <p>Pain Reliever/Fever Reducer</p> <p>Lasts up to 8 hours For the temporary relief of minor arthritis pain</p> <p>DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN</p> <p>*Capsule-Shaped Bi-Layer Tablets</p> <p>ACTUAL SIZE</p> <p>100 Caplets* 650 mg EACH</p>		<p>TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING</p>	<p>Drug Facts</p> <p>Active ingredient (in each caplet) Purpose Acetaminophen 650 mg.....Pain reliever/fever reducer</p> <p>Uses</p> <ul style="list-style-type: none"> ■ temporarily relieves minor aches and pains due to: <ul style="list-style-type: none"> ■ minor pain of arthritis ■ muscular aches ■ backache ■ premenstrual and menstrual cramps ■ the common cold ■ headache ■ toothache ■ temporarily reduces fever <p>Warnings</p> <p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 6 caplets in 24 hours, which is the maximum daily amount</p> <ul style="list-style-type: none"> ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product 	<p>Drug Facts (continued under label)</p> <p>This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., McNeil Consumer Healthcare division, owner of the registered trademark Tylenol® for Arthritis Pain</p> <p>Distributed by: Time-Can Labs, Inc. 7 Michael Avenue, Farmingdale, NY 11735 659R 0723 Made in India</p> <p>Lot No.: 3 Exp Date: 01/2023</p> <p>Varnish Omit Area</p> <p>PEEL HERE</p>
--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--	-----------------------------------------------------------------------------------------------------------------------------------------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

NDC 49483-699-40



¹Compare to Tylenol® 8hr Arthritis Pain active ingredient

Arthritis Pain Relief

Acetaminophen

Extended-Release Tablets USP, 650 mg
Pain Reliever/Fever Reducer

Lasts up to 8 hours
For the temporary relief of
minor arthritis pain



ACTUAL SIZE

400 Caplets*
650 mg EACH

DO NOT USE WITH OTHER MEDICINES
CONTAINING ACETAMINOPHEN

*Capsule-Shaped Bi-Layer Tablets

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts

Active ingredient (in each caplet) Purpose
Acetaminophen 650 mg.....Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches
 - backache
 - premenstrual and menstrual cramps
 - the common cold
 - headache
 - toothache
- temporarily reduces fever

Warnings

- Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take
 - more than 6 caplets in 24 hours, which is the maximum daily amount
 - with other drugs containing acetaminophen
 - 3 or more alcoholic drinks every day while using this product
- Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:
 - skin reddening
 - blisters
 - rash
 If a skin reaction occurs, stop use and seek medical help right away.
- Do not use**
 - with any other drug containing acetaminophen

Drug Facts (continued)

(prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - new symptoms occur
 - redness or swelling is present
- These could be signs of a serious condition.

if pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

Do not take more than directed. See overdose warning.

- adults
- take 2 caplets every 8 hours with water

Drug Facts (continued)

- swallow whole; do not crush, chew, split, or dissolve
- do not take more than 6 caplets in 24 hours
- do not use for more than 10 days unless directed by a doctor

under 18 years of age

- ask a doctor

Other information

- store between 20-25°C (68-77°F)
- The FDA approved dissolution methods differ from USP

Inactive ingredients: carnauba wax, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, povidone, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin

Questions or comments? Call 1-877-290-4008

Contains No Aspirin

This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., McKell Consumer Healthcare division, owner of the registered trademark Tylenol® & Arthritis Pain

Distributed by: Time-Cap Labs, Inc.
7 Michael Avenue, Farmingdale, NY 11735
699R 0723 Made in India



Lot No.: Varnish Omit Area
Exp. Date: 4948316940

NDC 49483-704-05



Muscle Aches & Pain

Acetaminophen

Extended-Release Tablets USP, 650 mg
Pain Reliever/Fever Reducer

For up to 8 hours relief of Minor Muscle Aches & Pain

DO NOT USE WITH OTHER MEDICINES
CONTAINING ACETAMINOPHEN

*Capsule-Shaped Bi-Layer Tablets

50 Caplets*
650 mg EACH

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

- Uses** temporarily relieves minor aches and pains due to:
 - muscular aches
 - backache
 - minor pain of arthritis
 - toothache
 - premenstrual and menstrual cramps
 - headache
 - the common cold
 - temporarily reduces fever

Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

 If a skin reaction occurs, stop use and seek medical help right away.

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

 These could be signs of a serious condition.

if pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** **Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions Do not take more than directed. See overdose warning adults and children 12 years and over:

- take 2 caplets every 8 hours with water
- swallow whole; do not crush, chew, split or dissolve
- do not take more than 6 caplets in 24 hours
- do not use for more than 10 days unless directed by a doctor

 children under 12 years: do not use

Other information

- store between 20-25°C (68-77°F)
- The FDA approved Dissolution methods differ from USP

Questions or comments? Call 1-877-290-4008

Distributed by: Time-Cap Labs, Inc. 7 Michael Avenue, Farmingdale, NY 11735 704R 1022 Made in India
40146401 Code No.: GO/DRUGS/515

Varnish Omit Area

Lot No.:
Exp. Date:

Print & Varnish
Omit

Lot No.:
Exp. Date:



TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

5155500300300 6029

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Drug Facts

Active ingredient **Purpose**
(In each caplet) **Acetaminophen 650 mg.....Pain reliever/fever reducer**

Uses
temporarily relieves minor aches and pains due to:
muscular aches ■ headache
minor pain of arthritis ■ toothache
premenstrual and menstrual cramps
headache ■ the common cold
temporarily reduces fever

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
more than 6 caplets in 24 hours, which is the maximum daily amount
with other drugs containing acetaminophen
3 or more alcoholic drinks every day while using this product
Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:
skin reddening ■ blisters ■ rash
if a skin reaction occurs, stop use and seek medical help right away.
Do not use
with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
if you are allergic to acetaminophen or any of the inactive ingredients in this product

Drug Facts (continued)

Ask a doctor before use if you have liver disease
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if
pain gets worse or lasts more than 10 days
fever gets worse or lasts more than 3 days
new symptoms occur
redness or swelling is present
These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.
Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions
Do not take more than directed. See overdose warning

adults and children 12 years and over
take 2 caplets every 8 hours with water
swallow whole; do not crush, chew, split or dissolve
do not take more than 6 caplets in 24 hours
do not use for more than 10 days unless directed by a doctor

children under 12 years
do not use

Other information
store between 20-25°C (68-77°F)
The FDA approved Dissolution methods differ from USP

Inactive ingredients caranuba wax, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, polydioxane, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin
Questions or comments? Call 1-877-296-4008

100146001

Contains No Aspirin

This product is not manufactured or distributed by McNeil Consumer Healthcare, owner of the registered trademark Tylenol® 8hr

Distributed by:
Time-Cap Labs, Inc.
7 Mitchell Avenue
Farmingdale, NY 11735
704K 102Z
Made in India

NDC 49483-704-05
*Compare to Tylenol® 8hr in active ingredient



Muscle Aches & Pain
Acetaminophen

Extended-Release Tablets USP, 650 mg
Pain Reliever/Fever Reducer
For up to 8 hours relief of Minor Muscle Aches & Pain



50 Caplets*
650 mg EACH

DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN
*Capsule-Shaped Bi-Layer Tablets

Varnish Omit

E1384

timely
Muscle Aches & Pain
Extended-Release Tablets USP, 650 mg
Pain Reliever/Fever Reducer
For up to 8 hours relief of Minor Muscle Aches & Pain
50 Caplets*
650 mg EACH
*Capsule-Shaped Bi-Layer Tablets





NDC 49483-699-26
 *Compare to Tylenol®
 8^{hr} Arthritis Pain
 active ingredient

Arthritis Pain Relief Acetaminophen

Extended-Release Tablets USP, 650 mg
Pain Reliever/Fever Reducer

Lasts up to 8 hours
For the temporary relief of
minor arthritis pain



DO NOT USE WITH OTHER MEDICINES
CONTAINING ACETAMINOPHEN

ACTUAL SIZE
225 Caplets*
650 mg EACH

*Capsule-Shaped Bi-Layer Tablets

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts

Active ingredient (in each caplet) Purpose
Acetaminophen 650 mg.....Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches
 - backache
 - premenstrual and menstrual cramps
 - the common cold
 - headache
 - toothache
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
 more than 6 caplets in 24 hours, which is the maximum daily amount
 with other drugs containing acetaminophen
 3 or more alcoholic drinks every day while using this product
Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:
 skin reddening blisters rash
 If a skin reaction occurs, stop use and seek medical help right away.

Do not use

with any other drug containing acetaminophen (prescription)

Drug Facts (continued under label)

This product is not manufactured or distributed by McNeil Consumer Healthcare, owner of the registered trademark Extra Strength Tylenol®.

Distributed by:

Time-Cap Labs, Inc.

7 Michael Avenue

Farmingdale, NY 11735

999R 1022 Made in India

Code No.: 60DRUGS515

40146301



Lot No.:

Exp. Date:

Varnish Omit Area



NDC 49483-699-42

Arthritis Pain Relief Acetaminophen

Extended-Release Tablets USP, 650 mg
Pain Reliever/Fever Reducer

Lasts up to 8 hours
For the temporary relief of
minor arthritis pain

DO NOT USE WITH OTHER MEDICINES
CONTAINING ACETAMINOPHEN

24 Caplets*
650 mg EACH

*Capsule-Shaped Bi-Layer Tablets

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Uses temporarily relieves minor aches and pains due to:

- minor pain of arthritis
- muscular aches
- backache
- premenstrual and menstrual cramps
- the common cold
- headache
- toothache

 temporarily reduces fever

Warnings **Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take more than 6 caplets in 24 hours, which is the maximum daily amount with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product. **Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening blisters rash If a skin reaction occurs, stop use and seek medical help right away. **Stop use and ask a doctor if:** pain gets worse or lasts more than 10 days fever gets worse or lasts more than 3 days new symptoms occur redness or swelling is present. These could be signs of a serious condition. **If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** **Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions Do not take more than directed. See overdose warning **adults:** take 2 caplets every 8 hours with water swallow whole; do not crush, chew, split or dissolve do not take more than 6 caplets in 24 hours do not use for more than 10 days unless directed by a doctor **under 18 years of age:** ask a doctor

Other information store between 20-25°C (68-77°F) The FDA approved Dissolution methods differ from USP

Questions or comments? Call 1-877-290-4008 Distributed by: Time-Cap Labs, Inc. 7 Michael Avenue, Farmingdale, NY 11735 999R 1022 Made in India 40146301 Code No.: 60DRUGS515

Lot No.:

Exp. Date:

Varnish Omit Area

Drug Facts (continued)

or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop using and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

Do not take more than directed.

See overdose warning

adults

- take 2 caplets every 8 hours with water
- swallow whole; do not crush, chew, split or dissolve
- do not take more than 6 caplets in 24 hours
- do not use for more than 10 days unless directed by a doctor

under 18 years of age

- ask a doctor

Other information

- store between 20-25°C (68-77°F)
- The FDA approved Dissolution methods differ from USP

Inactive ingredients

carmauba wax, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, povidone, pregelatinized starch, sodium starch glycolate, titanium dioxide, Tracelrin

Questions or comments? Call 1-877-290-4008

Contains No Aspirin

Adhesive Area

Lot No.: Varnish Omit
Exp Date:



3 16948316994215
6998 1022 Made in India
Distributed by: Time-Cap Labs, Inc.
7 Michael Avenue, Farmingdale, NY 11735
Registered trademark "Timely"
Johnson & Johnson Consumer Inc., owner of the
This product is not manufactured or distributed by
THE IMPRINTED FOIL SEAL OVER THE MOUTH OF
THE BOTTLE IS CUT, TORN, BROKEN OR MISSING
TAMPER EVIDENT. DO NOT USE THIS PRODUCT IF

0066 No. 60DRUS515

40146101

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Drug Facts

Active ingredient (in each caplet) **Purpose**
Acetaminophen 650 mg.....Pain reliever/
fever reducer

Uses ■ temporarily relieves minor aches and pains due to: ■ minor pain of arthritis
■ muscular aches ■ backache
■ premenstrual and menstrual cramps
■ the common cold ■ headache ■ toothache
■ temporarily reduces fever

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
■ more than 6 caplets in 24 hours, which is the maximum daily amount
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product
Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:
■ skin reddening ■ blisters ■ rash

Drug Facts (continued)

If a skin reaction occurs, stop use and seek medical help right away.

Do not use
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if
■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days
■ new symptoms occur
■ redness or swelling is present
These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions Do not take more than directed. See overdose warning

adults
■ take 2 caplets every 8 hours with water
■ swallow whole; do not crush, chew, split or dissolve
■ do not take more than 6 caplets in 24 hours
■ do not use for more than 10 days unless directed by a doctor

under 18 years of age ■ ask a doctor

Other information

■ store between 20-25°C (68-77°F)
■ The FDA approved Dissolution methods differ from USP

Inactive ingredients carnauba wax, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, povidone, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin

Questions or comments?
Call 1-877-290-4008

Contains No Aspirin

NDC 49483-699-42
Compare to Tylenol® & Arthritis Pain active ingredient



ACTUAL SIZE
24 Caplets*
650 mg EACH

timely™
Arthritis Pain Relief

Acetaminophen
Extended-Release Tablets USP, 650 mg
Pain Reliever/Fever Reducer

Lasts up to 8 hours
For the temporary relief of minor arthritis pain

DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN
*Caplets Shaped Like Tylenol® Tablets

Arthritis Pain Relief
timely™
Acetaminophen
Extended-Release Tablets USP, 650 mg
Pain Reliever/Fever Reducer
Lasts up to 8 hours
For the temporary relief of minor arthritis pain
24 Caplets* 650 mg EACH
*Caplets Shaped Like Tylenol® Tablets



Read the important information about this medicine carefully before you take it, and then each time you take it. This information does not take the place of that from your doctor about your condition or treatment. This information does not tell you everything you need to know about this medicine. You should always follow the directions on the product labeling and ask your doctor or pharmacist if you have any questions about the medicine.

Drug Facts
Active ingredient: Acetaminophen
Other ingredients: ...
Contains: ...
Warnings: ...
Directions: ...
Other information: ...
Questions or comments? ...

Top of Package
 Contains 50 caplets
 Do not use if you have had a seizure in the last 2 weeks
 Do not use if you are taking other medicines that contain acetaminophen
 Do not use if you are taking other medicines that contain salicylates
 Do not use if you are taking other medicines that contain alcohol
 Do not use if you are taking other medicines that contain ...

Side of Package
 Print & Varnish Omit
 Lot No.:
 Exp. Date:
 2437

Drug Facts (Continued)
Directions: ...
Warnings: ...
Directions: ...
Other information: ...
Questions or comments? ...

Bottom of Package
 Contains 50 caplets
 Do not use if you have had a seizure in the last 2 weeks
 Do not use if you are taking other medicines that contain acetaminophen
 Do not use if you are taking other medicines that contain salicylates
 Do not use if you are taking other medicines that contain alcohol
 Do not use if you are taking other medicines that contain ...

Side of Package
 Print & Varnish Omit
 Lot No.:
 Exp. Date:
 2437

Other information: ...
Questions or comments? ...

Bottom of Package
 Contains 50 caplets
 Do not use if you have had a seizure in the last 2 weeks
 Do not use if you are taking other medicines that contain acetaminophen
 Do not use if you are taking other medicines that contain salicylates
 Do not use if you are taking other medicines that contain alcohol
 Do not use if you are taking other medicines that contain ...

Side of Package
 Print & Varnish Omit
 Lot No.:
 Exp. Date:
 2437

NDC 49483-699-06
 "Compare to Tylenol" &
 Arthritis Pain active ingredient

timely™
 Arthritis Pain Relief
Acetaminophen
 Extended-Release Tablets USP, 650 mg
Pain Reliever/Fever Reducer

Lasts up to 8 hours
 For the temporary relief of minor arthritis pain

DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN
 *Capsule-Shaped Bi-Layer Tablets

50 Caplets*
 650 mg EACH

Actual Size

Varnish Omit E1384

NDC 49483-699-05

"Compare to Tylenol" &
 Arthritis Pain active ingredient

timely™
 Arthritis Pain Relief
Acetaminophen
 Extended-Release Tablets USP, 650 mg
Pain Reliever/Fever Reducer

Lasts up to 8 hours
 For the temporary relief of minor arthritis pain

DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN
 *Capsule-Shaped Bi-Layer Tablets

Actual Size
 50 Caplets*
 650 mg EACH

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Uses temporarily relieves minor aches pains due to:
 • minor of arthritis • muscular aches • backache
 • menstrual and menstrual cramps • the common cold
 • headache • toothache • temporarily reduces fever

Warnings: Liver warning: This product contains acetaminophen. Severe liver damage may occur if you: **take** more than 6 caplets in 24 hours, which is the maximum daily amount • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product. **Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:
 • skin reddening • blisters • rash • If a skin reaction occurs, stop use and seek medical help right away. **Stop use and ask a doctor if:** • pain gets worse or lasts more than 10 days • fever gets worse or lasts more than 3 days • new symptoms occur • redness or swelling is present These could be signs of a serious condition. If pregnant or breastfeeding, a health professional should be consulted before use. **Keep out of reach of children.** Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions do not take more than directed (see overdose warning) adults: take 2 caplets every 8 hours with water swallow whole; do not crush, chew, split or dissolve do not take more than 6 caplets in 24 hours do not use for more than 10 days unless directed by a doctor under 18 years of age: ask a doctor

Other information store between 20-25°C (68-77°F)
 The FDA approved Dissolution methods differ from USP
Questions or comments? Call 1-877-290-4008

This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., owner of the registered trademark "Tylenol" Distributed by: Time-Cap Labs, Inc.
 7 Michael Avenue, Farmingdale, NY 11735
 689R 0921 Made in India
 99IL437 Code No.: 60DRUGS/515

Lot No.:
Exp. Date:
 Varnish Omit Area

ACETAMINOPHEN

acetaminophen tablet extended release tablet, extended release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49483-699
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
POVIDONE K30 (UNII: U725QWY32X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
TRIACETIN (UNII: XHX3C3X673)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)	
HYDROXYETHYL CELLULOSE (140 MPA.S AT 5%) (UNII: 8136Y38GY5)	

Product Characteristics

Color	white (White to off white)	Score	no score
Shape	CAPSULE	Size	19mm
Flavor		Imprint Code	71
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49483-699-40	400 in 1 BOTTLE; Type 0: Not a Combination Product	10/04/2021	
2	NDC:49483-699-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	10/04/2021	
3	NDC:49483-699-05	50 in 1 CARTON	10/04/2021	
3		1 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:49483-699-26	225 in 1 BOTTLE; Type 0: Not a Combination Product	11/07/2022	
5	NDC:49483-699-42	1 in 1 CARTON	11/07/2022	

5	24 in 1 BOTTLE; Type 0: Not a Combination Product		
---	---------------------------------------------------	--	--

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA215486	10/04/2021	

MUSCLE ACHES AND PAINS ACETAMINOPHEN EXTENDED RELEASE

acetaminophen tablet, extended release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49483-704
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
POVIDONE K30 (UNII: U725QWY32X)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
TRIACETIN (UNII: XHX3C3X673)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)	
HYDROXYETHYL CELLULOSE (140 MPa.S AT 5%) (UNII: 8136Y38GY5)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	

Product Characteristics

Color	white (White to off white)	Score	no score
Shape	CAPSULE (Capsule-shaped tablet)	Size	19mm
Flavor		Imprint Code	71
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	NDC:49483-704			

1	NDC:49483-704-05	1 in 1 CARTON	11/07/2022	
1		50 in 1 BOTTLE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
ANDA	ANDA215486		11/07/2022	

Labeler - TIME CAP LABORATORIES, INC. (037052099)

Registrant - TIME CAP LABORATORIES, INC. (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LIMITED		925822975	manufacture(49483-699, 49483-704)

Revised: 3/2024

TIME CAP LABORATORIES, INC.