

**BERKLEY AND JENSEN IBUPROFEN- ibuprofen tablet, film coated
BJWC**

BJWC Ibuprofen Tablets 200 mg Drug Facts

Active ingredient (in each tablet)

Ibuprofen 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- headache
- muscular aches
- minor pain of arthritis
- toothache
- backache
- the common cold
- menstrual cramps
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chances are higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen,

or others]

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **do not take more than directed**
- the smallest effective dose should be used
- Adults and children 12 years and older:
 - take 1 tablet every 4 to 6 hours while symptoms persist
 - if pain or fever does not respond to 1 tablet, 2 tablets may be used
 - do not exceed 6 tablets in 24 hours, unless directed by a doctor
- Children under 12 years: ask a doctor

Other information

- **read all warnings and directions before use**
- store at 20-25°C (68-77°F)
- avoid high humidity and excessive heat above 40°C (104°F)

Inactive ingredients

colloidal silicon dioxide, corn starch, croscarmellose sodium, hypromellose, iron oxide red, iron oxide yellow, microcrystalline cellulose, polyethylene glycol, polysorbate 80, stearic acid, titanium dioxide

Questions or comments?

1-800-934-1204

Principal Display Panel

Compare to the active ingredient in Advil®

SEE NEW WARNINGS

IBUPROFEN TABLETS, 200mg

PAIN RELIEVER/FEVER REDUCER (NSAID)

GLUTEN FREE

gf

500 TABLETS

200 mg EACH

ACTUAL SIZE

Compare to the active ingredient in Advil®**

SEE NEW WARNINGS

berkley jensen

IBUPROFEN

TABLETS, 200mg

PAIN RELIEVER/FEVER REDUCER (NSAID)

GLUTEN FREE

gf

500 TABLETS | 200 mg EACH

ACTUAL SIZE

2

Drug Facts

Active ingredient (in each tablet)
Ibuprofen 200 mg (NSAID)* Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Purposes
Pain reliever/fever reducer

Uses
temporarily relieves minor aches and pains due to:
headache muscular aches minor pain of arthritis toothache
backache the common cold menstrual cramps temporarily reduces fever

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives facial swelling
asthma (wheezing) shock skin reddening rash blisters
If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chances are higher if you are age 60 or older
have had stomach ulcers or bleeding problems take a blood thinning (anticoagulant) or steroid drug take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) have 3 or more alcoholic drinks every day while using this product take more or for a longer time than directed
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.
Do not use if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer night before or after heart surgery
Ask a doctor before use if you have problems or serious side effects from taking pain relievers or fever reducers the stomach bleeding warning applies to you you have a history of stomach problems, such as heartburn you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke you are taking a diuretic
Ask a doctor or pharmacist before use if you are taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin under a doctor's care for any serious condition taking any other drug
When using this product take with food or milk if stomach upset occurs

Drug Facts (continued)
Stop use and ask a doctor if you experience any of the following signs of stomach bleeding:
feel faint vomit blood have bloody or black stools have stomach pain that does not get better you have symptoms of heart problems or stroke: chest pain trouble breathing
weakness in one part or side of body slurred speech leg swelling pain gets worse or lasts more than 10 days fever gets worse or lasts more than 3 days redness or swelling is present in the painful area any new symptoms appear
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions
do not take more than directed the smallest effective dose should be used
Adults and children 12 years and older:
take 1 tablet every 4 to 6 hours while symptoms persist
if pain or fever does not respond to 1 tablet, 2 tablets may be used
do not exceed 6 tablets in 24 hours, unless directed by a doctor
Children under 12 years: ask a doctor

Other information
read all warnings and directions before use
store at 20-25°C (68-77°F)
avoid high humidity and excessive heat above 40°C (104°F)

Inactive ingredients colloidal silicon dioxide, corn starch, croscarmellose sodium, hypromellose, iron oxide red, iron oxide yellow, microcrystalline cellulose, polyethylene glycol, polysorbate 80, stearic acid, titanium dioxide

Questions or comments? 1-800-934-1204

**This product is not manufactured or distributed by Pfizer, distributor of Advil®.

Every Berkley Jensen product comes with our 100% MONEY-BACK GUARANTEE.

Distributed by: BJs Wholesale Club
25 Research Drive
Westborough, MA 01581

8 88670 01802 6

Man. # 075798058

EU # 00000

CODE AREA

: L0490 D3 F8

BERKLEY AND JENSEN IBUPROFEN

ibuprofen tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68391-604
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XY110QM) (IBUPROFEN - UNII:WK2XY110QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	

Product Characteristics

Color	BROWN	Score	no score
Shape	ROUND	Size	10 mm

Flavor		Imprint Code	I2	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68391-604-62	1 in 1 CARTON	07/27/2010	
1		24 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:68391-604-90	500 in 1 BOTTLE; Type 0: Not a Combination Product	06/20/2007	
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA072096	06/20/2007		

Labeler - BJWC (159082692)

Revised: 12/2019

BJWC