

PAIN RELIEF FAST RELIEF- aspirin, citric acid monohydrate, and sodium bicarbonate tablet, effervescent
Select Brand

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Effervescent Pain Relief
Fast Relief

Drug Facts

Active ingredients (in each effervescent tablet)	Purpose
Aspirin (NSAID)* 325 mg	Pain reliever
Citric acid 1000 mg	Antacid
Sodium bicarbonate (heat-treated) 1916 mg	Antacid

* nonsteroidal anti-inflammatory drug

Uses

For the temporary relief of: heartburn, acid indigestion, and sour stomach when accompanied with headache or body aches and pains

- upset stomach with headache from overindulgence in food or drink
- headache, body aches and pain alone

Warnings

Reye's syndrome

Children and teenagers should not use this medicine for chicken pox or flu symptoms before a doctor is consulted about Reye's syndrome, a rare but serious illness reported to be associated with aspirin.

Allergy alert

Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning

This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take other drugs containing prescription or nonprescription NSAID's (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for longer time than directed

Alcohol warning

If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take aspirin or other pain relievers/fever reducers.

Aspirin may cause stomach bleeding.

Do not use: if you have ever had an allergic reaction to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have a sodium-restricted diet

Ask a doctor or pharmacist before use if

- you are presently taking a prescription drug. Antacids may interact with certain prescription drugs.
- you are taking a prescription drug for anticoagulation (thinning the blood), diabetes, gout or arthritis.

When using this product do not exceed the recommended dosage

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better
- symptoms get worse or last more than 10 days
- new symptoms occur
- redness or swelling is present
- ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- Dissolve tablets in 4 ounces of water before taking

Adults and Children 12 Years and over	<ul style="list-style-type: none">▪ 2 tablets every 4 hours, or as directed by a doctor▪ do not exceed 8 tablets in 24 hours
60 Years and over	<ul style="list-style-type: none">▪ 2 tablets every 4 hours, or as directed by a doctor▪ do not exceed 4 tablets in

24 hours

Other information

- **Each tablet contains:** sodium 540 mg
- store at room temperature. Avoid excessive heat
- in water, this tablet contains principally the antacid sodium citrate and the analgesic sodium acetylsalicylate

Inactive Ingredients

None

Distributed by
select brand[®] /SAJ Distributors
3017 N. Midland Drive
Pine Bluff, AR 71603

PRINCIPAL DISPLAY PANEL - 36 Tablet Carton

NDC 15127-968-36

select brand[®]
the lower price name brand

FAST RELIEF

EFFERVESCENT
Pain Relief

Aspirin (NSAID) / Pain reliever,
Citric acid / Antacid, Sodium bicarbonate / Antacid.

Relieves:
Heartburn • Acid Indigestion
• Sour Stomach with
Headache
• Body Aches
and Pain

Compare to Active
Ingredients in
Alka-Seltzer[®] *

36 EFFERVESCENT
TABLETS

select brand
the lower price name brand

EFFERVESCENT

Pain Relief

Tablets sealed in printed packets for your protection.
Do not use if torn or punctured.

NDC 15127-968-36

select brand
the lower price name brand

FAST RELIEF

EFFERVESCENT

Pain Relief

Aspirin (NSAID) / Pain reliever,
Citric acid / Antacid, Sodium bicarbonate / Antacid.

Relieves:

- Heartburn • Acid Indigestion
- Sour Stomach with Headache
- Body Aches and Pain

Compare to Active Ingredients in Alka-Seltzer.*

36 EFFERVESCENT TABLETS

Place two (2) tablets in 4 oz. of water. Dissolve tablets completely.



You do not have to drink any of the residue that may be on the bottom of the glass. The medicines are in the water.

Take this product any time - morning, noon or night - when you need relief from heartburn, acid indigestion, sour stomach with headache or body aches and pains.



Distributed by
select brand® /SAJ Distributors
3017 N. Midland Drive
Pine Bluff, AR 71603

BOXD-020 REV-4



Drug Facts

Active ingredients (in each effervescent tablet) Purpose

Aspirin (NSAID)* 325 mg	Pain reliever
Citric acid 1000 mg	Antacid
Sodium bicarbonate (heat-treated) 1916 mg	Antacid

*nonsteroidal anti-inflammatory drug

Uses

For the temporary relief of: heartburn, acid indigestion, and sour stomach when accompanied with headache or body aches and pains ■ upset stomach with headache from overindulgence in food or drink ■ headache, body aches and pain alone

Warnings

Reye's syndrome: Children and teenagers should not use this medicine for chicken pox or flu symptoms before a doctor is consulted about Reye's syndrome, a rare but serious illness reported to be associated with aspirin.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take other drugs containing prescription or nonprescription NSAID's (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for longer time than directed

Alcohol warning: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take aspirin or other pain relievers/fever reducers. Aspirin may cause stomach bleeding.

Do not use: if you have ever had an allergic reaction to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if: ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma ■ you have a sodium-restricted diet

Ask a doctor or pharmacist before use if: ■ you are presently taking a prescription drug. Antacids may interact with certain prescription drugs. ■ you are taking a prescription drug for anticoagulation (thinning the blood), diabetes, gout or arthritis.

When using this product do not exceed the recommended dosage ►

Drug Facts (continued)

Stop use and ask a doctor if: ■ an allergic reaction occurs. Seek medical help right away. ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ symptoms get worse or last more than 10 days ■ new symptoms occur ■ redness or swelling is present ■ ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

■ Dissolve tablets in 4 ounces of water before taking

Adults and Children 12 Years and over	■ 2 tablets every 4 hours, or as directed by a doctor ■ do not exceed 8 tablets in 24 hours
60 Years and over	■ 2 tablets every 4 hours, or as directed by a doctor ■ do not exceed 4 tablets in 24 hours

Other information

■ Each tablet contains: sodium 540 mg
■ store at room temperature. Avoid excessive heat
■ in water, this tablet contains principally the antacid sodium citrate and the analgesic sodium acetylsalicylate

Inactive Ingredients None

†Alka-Seltzer® is a registered trademark of Bayer HealthCare LLC, Morristown, NJ 07962.
This product is not manufactured or distributed by Bayer HealthCare LLC. DL1 DFrev3-210

PAIN RELIEF FAST RELIEF

aspirin, citric acid monohydrate, and sodium bicarbonate tablet, effervescent

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:15127-968
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Aspirin (UNII: R16CO5Y76E) (Aspirin - UNII:R16CO5Y76E)	Aspirin	325 mg
Citric Acid Monohydrate (UNII: 2968PHW8QP) (Anhydrous Citric Acid - UNII:XF417D3PSL)	Anhydrous Citric Acid	1000 mg
Sodium Bicarbonate (UNII: 8MDF5V39QO) (Bicarbonate Ion - UNII:HN1ZRA3Q20)	Sodium Bicarbonate	1916 mg

Product Characteristics

Color	WHITE	Score	2 pieces
Shape	ROUND	Size	25mm
Flavor		Imprint Code	C11
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:15127-968-36	18 in 1 CARTON		
1		2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part343	10/23/2006	

Labeler - Select Brand (043562370)

Revised: 9/2014

Select Brand