ARNICARE- arnica montana gel Laboratoires Boiron

Disclaimer: This homeopathic product has not been evaluated by the Food and Drug Administration for safety or efficacy. FDA is not aware of scientific evidence to support homeopathy as effective.

Arnicare Gel

Active ingredient**

Arnica montana 1X HPUS 7%

The letters "HPUS" indicate that the component in this product is officially monographed in the Homeopathic Pharmacopoeia of the United States.

Purpose*

Arnica montana 1X HPUS 7% Relieves muscle pain & stiffness, swelling from injuries, discoloration from bruises

Uses*

temporarily relieves muscle pain and stiffness due to:

- minor injuries
- overexertion
- falls

reduces symptoms of bruising such as:

- pain
- swelling
- discoloration

Warnings

For external use only.

Do not use if you are allergic to Arnica montana or to any of this product's inactive ingredients.

When using this product

avoid contact with eyes, mucous membranes, wounds, damaged or irritated skin

use only as directed

dryness or irritation may occur

do not tightly wrap or bandage the treated area

do not apply heat or ice to treated area immediately before or after use

Stop use and ask a doctor if

- condition persists for more than 3 days or worsens
- symptoms clear up and occur again within a few days

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

Apply a thin layer of Arnicare Gel to the affected area as soon as possible after minor injury. Repeat 3 times a day or as needed. If heat or ice is applied, wait 5 minutes before applying Arnicare Gel.

Arnicare Gel Roll-On (2019)- Shake well before each use. With the applicator facing down, lightly squeeze the tube then roll ball against skin to apply a thin layer of the product to affected area. Massage gently into skin 3 times a day or as needed. If heat or ice is applied, wait 5 minutes before applying Arnicare Roll-on.

Arnicare Gel Roll-On (2023)- Twist roller to the "ON" position and gently squeeze tube to apply a thin layer of product to affected area. Twist roller to the "OFF" position and continue massaging as needed. Repeat 3 times a day. If heat or ice is applied, wait 5 minutes before applying Arnicare Roll-on Gel.

Alcohol, carbomer, purified water, sodium hydroxide

Other information

do not use if glued carton end flaps are open or if the tube seal is broken

do not use if pouch is torn or open

Arnicare Gel 0.18 oz (5g)- dispose 1 day after opening

Arnicare Gel Roll-On (2019)- dispose of 6 months after opening

Arnicare Gel Roll-On (2023)- dispose of 1 year after opening

0.18 oz (5g)

0.5 oz (14g)

- 1.5 oz (45g)
- 2.6 oz (75g)
- 4.2 oz (120g)

3 oz (90g)

Pain Relief*

Muscle Pain & Stiffness Swelling from Injuries Bruises*

*CLAIMS BASED ON TRADITIONAL HOMEOPATHIC PRACTICE, NOT ACCEPTED MEDICAL EVIDENCE. NOT FDA EVALUATED.

*C,K,CK, and X are homeopathic dilutions: see BoironUSA.com/info for details.

Questions, Comments?

Distributed by Boiron, Inc., Newtown Square, PA 19073

1-800-BOIRON-1 (1-800-264-7661)

BoironUSA.com

Arnicare.com Info@Boiron.com Made in France





Quickly Absorbed

NET WT.

1.5 oz (45 a

NDC 0220-9000-84

PAIN RELIEF

Bruises

- Fragrance-Free
- Plant-Powered Relief

The state of the s





Twist roller to the "OFF" position and continue massaging as needed. Repeat 3 times a day. applied, wait 5 minutes before applying Amicare Rol-On GeL Other information do not use if glued carton end flaps are open 🛛 🔳 dispose of 1 year after opening Inactive ingredients acohol, carbomer, purified water, sodium hydroxide Made in France Distributed by Boiron Inc., Newtown Square, PA 19073 1-800-80 RON-1 (1-800-264-7661) Amicare.com Drug Facts Active ingredient Purpose Relieves muscle pain & Arnica montana ... stiffness, swelling from injuries, 1X HPUS 7% discoloration from bruises Uses temporarily relieves muscle pain and stiffness due to: minor injuries overexertion fa s reduces symptoms of bruising such as: pain sweing discoloration Warnings For external use only. Do not use if you are allergic to Arnica montana or to any of this product's inactive ingredients. When using this product avoid contact with eyes, mucous membranes, wounds, damaged or irritated skin 🛛 🔳 use only as directed dryness or irritation may occur do not tightly wrap or bandage the treated area. do not apply heat or ice to the treated area immediately

Stop use and ask a doctor if Condition persists for more than 3 days or worsens

help or contact a Poison Control Center right away. Directions Twist roler to the "ON" position and gently squeeze tube to apply a thin layer of product to affected area.

symptoms clear up and occur again within a few days Keep out of reach of children. If swallowed, get medical

before or after use Stop use and ask a doctor if

condition persists for more than 3 days or worsens symptoms clear up and occur again within a few days Keep out of reach of children, I swallowed, get medical help or contact a Poison Control Center right away. Directions Twist roler to the "ON" position and gently squeeze tube to apply a thin layer of product to affected area. Twist roller to the "OFF" position and continue massaging as needed. Repeat 3 times a day. applied, wait 5 minutes before applying Amicare Roll-On Gel. Other information do not use if glued carton end flaps are open dispose of 1 year after opening Inactive ingredients acohol, carbomer, purified water, socium hydroxide Made in France Distributed by Boiron Inc., Newtown Square, PA 19073

1-800-B0IRON-1 (1-800-264-7661) | Amicare.com

4-05/73 UPILO

TUP 1JC4 05/23

Purpose

I faits





Drug Facts Active ingredient Amica montana 1X HPUS 7% Purpose Relieves muscle pain & stiffness, swelling from injuries, discoloration from bruises Uses temporarily relieves muscle pain and stiffness due to: minor injuries moverexertion mitals reduces symptoms of bruising such as: sweling discoloration a pain Warmings For external use only. Do not use if you are allergic to Arrica montana or to any of this product's inactive ingredients. When using this product and contact with eyes, mucous membranes, wounds, damaged or initiated skin and use only as directed and infrintes or initiation may occur and on ottightly wrap or bandage the treated area. and not apply heat or los to the treated area immediately before area and on rot apply heat or los to the treated area immediately before area and the second and a dealer if the conditioned area. or after use. Stop use and ask a doctor if m condition persists for more than 3 days or worsets musymptoms dear up and occur again within a few days. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. Directions: Apply a thin layer of Arricare Gel to affected area as soon as possible after minor injury, Repeat 3 times a day. If heat or ice is applied, wait 5 minutes before applying Arricare Gel. Other Information mdo not use if pouch is torn or open dispose of 1 day after opening Inactive ingredients alcohol, carborner, purified water, socium hydroxide -0 Questions or comments? Amicare.com • 1-800-264-7661 0 **Nade in France** Distributed by Boiron Inc. ALU.1197 "CLAIMS BASED ON TRADITIONAL HOMEOPATHIC PRACTICE, NOT ACCEPTED MEDICAL EVIDENCE NOT FOA EVALUATED, ""C, K, CK, and X are homeopathic dilutions: see BoironUSA.com/into for details. Newtown Square, PA 19073 ò 0 M









Drug Facts (continued)

Other information do not use if glued carton end flaps are open or if the tube seal is broken

Inactive ingredients a coho, carbomer, purified water, sodium hydroxide

Questions or comments? Amicare.com | BoironUSA.com nfo@Boiron.com 1-800-80 RON-1 (1-800-264-7661)

The Boiron Promise

Your health deserves the greatest respect. That's why we're committed to your satisfaction. Please visit BoironUSA.com/promise for details.

> Made in France Distributed by Boiron Inc. Newtown Square, PA 19073

(B) please recycle





Purpose





Drug Facts Active ingredient"

Purpose

Uses Temporarily relieves muscle pain and stiffness due to: Initian injuries overexertion fails reduces symptoms of bruising such as: I pain swelling discoloration Warnings For external use only. Do not use if you are allergic to Arnica montana or to any of this product in active ingredients. When using this product avoid contact with eyes, nuccus membranes, wounds, damaged or initiated skie use only as directed the treated area I do not apply heat or izerts the treated area do not apply heat or izerts the treated area do not apply heat or izerts the treated area Stop use and the tadoctor if n condition parsists for apple than 3 days or worsens symptoms clear up apployceur again within a few days. Keep out of reach of oplighter. If swellowed, get medical help or contact a Power Directions

Directions Apply a thin layer of Arnicare Gel to affected area as soon as possible after minor njury. Repeat 3 times a day or as needed. If heat or ice is applied, wait 5 minutes before applying Amicare Gel.

Other information do not use if glued carlon end flaps are open or if the tube seal is broken

Inactive ingredients alcohol, carbomer, purified water, sodium hydroxide

Questions or comments? Annicare.com I BoironUSA.com Info@Boiron.com 1-500-50BR0N-1 (1-800-264-7661)

Made in France Distributed by Boiron Inc., Newtown Square, PA 19073

*CLAIMS BASED ON TRADITIONAL HOMEOPATHIC PRACTICE, NOT ACCEPTED MEDICAL EVIDENCE, NOT FDA EVALUATED, **C, K, CK, and X are homeopathic diutions: see BoironUSA.com/info for details. TUA 1FPT-04-2020





























Drug Facts

Purpose'

Trauma, muscle

pain & stiffness, swelling from injuries, discoloration from bruising The letters HPUS indicate that this incredient

is officially included in the Homeopathic Pharmacopoeia of the United States.

temporarily relieves muscle pain and stiffness due to minor injuries, overexertion and falls reduces pain, swelling and discoloration from bruises

For external use only. Avoid contact with eyes, mucous membranes, damaged skin or wounds. Do not use if you are allergic to Arnica montana or to any of this product's

When using this product use only as directed do not bandage tightly or use a heating pad.

Stop use and ask a doctor if condition persists for more than 3 days

Keep out of reach of children. f swallowed, get medical help or contact a Poison Control Center

Apply a thin layer of Arnicare Ge to affected area and massage gently as soon as possible after minor injury. Repeat 3 times a day or as needed.

Other information

do not use if glued carton end flaps are open or if the tube seal is broken store at 68-77°F (20-25°C)

"These "Uses" have not been evaluated by the Food and Drug Administration. "C, K, CK, and X are homeopathic dilutions: see www.boironusa.com for details.



Drug Facts

(continued)

Other information do not use if the tube seal is broken

Inactive ingredients alcohol, carborner, purified water, sodium hydroxide

Made in France 🛞 please recycle

Questions or comments? Arnicare.com BoironUSA.com 1-800-BOIRON-1 (1-800-264-7661) Distributed by Boiron Inc. 6 Campus Boulevard Newtown Square, PA 19073-3267

*Homeopathic mother tincture made from Amica montana fresh whole plant. "These "Uses" have not been evaluated by the Food and Drug Administration, "C, K, CK, and X are homeopathic dilutions: see BoironUSA.com/info for details.




FEEL BETTER FASTER

Neck, Back, Shoulder, Leg and Foot Muscle Pain & Stiffness Swelling from Injuries • Bruising*



Arnicare Gel is made from Arnica montana (a mountain daisy), which has been used for centuries to naturally relieve pain. It has a cooling effect for fast pain relief. Arnicare is recommended by doctors, plastic surgeons and pharmacists and used by professional athletes and savvy moms.





ARNICARE					
arnica montana gel					
Product Information					
Product Type	HUMAN OTC DRUG	Item Code (So	ource)	NDC:0)220-9000
Route of Administration	TOPICAL				
a					
Active Ingredient/Active Moiety					
Ingredient Name			Basis of Strengt		Strength
ARNICA MONTANA (UNII: 080TY20 UNII:080TY208ZW)		ARNICA MONTANA		1 [hp_X] in 1 g	
Inactive Ingredients					
	Ingredient Name				Strength

WATER	(UNII:	059QF0KO0R)
-------	--------	-------------

SODIUM HYDROXIDE (UNII: 55X04QC32I)

CARBOMER HOMOPOLYMER TYPE C (UNII: 4Q93RCW27E)

Packaging

#	ltem Code	Package Description		Marketing Start Date	Marketing End Date	
1	NDC:0220-9000- 66	1 in 1 PACKAGE	01	/09/2007		
1		120 g in 1 TUBE; Type 0: Not a Combination Product				
2	NDC:0220-9000- 59	1 in 1 PACKAGE	01	/09/2007		
2		75 g in 1 TUBE; Type 0: Not a Combination Product				
3	NDC:0220-9000- 54	1 in 1 PACKAGE	01	/09/2007		
3		45 g in 1 TUBE; Type 0: Not a Combination Product				
4	NDC:0220-9000- 65	1 in 1 PACKAGE	01/09/2007			
4		14 g in 1 TUBE; Type 0: Not a Combination Product				
5	NDC:0220-9000- 84	1 in 1 PACKAGE	06/01/2018			
5		45 g in 1 TUBE; Type 0: Not a Combination Product				
6	NDC:0220-9000- 86	2 in 1 PACKAGE	06	/01/2018		
6		45 g in 1 TUBE; Type 0: Not a Combination Product				
7	NDC:0220-9000- 94	5 g in 1 POUCH; Type 0: Not a Combination Product	06/09/2023			
Marketing Information						
	Marketing Category	Application Number or Monograp Citation			Marketing End Date	
unapproved homeopathic			01/09/2007			

Labeler - Laboratoires Boiron (282560473)

Registrant - Boiron Inc. (014892269)

Establishment					
Name	Address	ID/FEI	Business Operations		
Boiron		282560473	manufacture(0220-9000)		