

POLYETHYLENE GLYCOL 3350 NF- polyethylene glycol 3350 powder, for solution
SUNRISE PHARMACEUTICAL, INC

Polyethylene Glycol 3350 powder, for solution

Drug Facts

Active ingredient (in each dose)

Polyethylene Glycol 3350, 17 g (cap filled to line)

Purpose

Osmotic Laxative

Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

Warnings

Allergy alert:

Do not use if you are allergic to polyethylene glycol

Do not use if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

Ask doctor or pharmacist before use if you are

taking a prescription drug

When using this product

you may loose, watery, more frequent stools

Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed unless advised by your doctor**
- the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (clear section in cap)
- adults and children 17 years of age and older:
 - fill to top of clear section in cap which is marked to indicate the correct dose (17 g)
 - stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
 - use once a day
 - use no more than 7 days
- children 16 years of age or under: ask a doctor

Other information

- store at 20° - 25°C (68° - 77°F)
- tamper-evident: do not use if foil seal under cap, printed with “SEALED for YOUR PROTECTION” is missing, open or broken

Inactive ingredient

none

Questions or comments?

1-800-FDA-1088

Manufactured & Distributed by:

Sunrise Pharmaceutical, Inc.

Rahway, NJ 07065

PRINCIPAL DISPLAY PANEL

NDC 11534-**180**-19

Polyethylene Glycol

3350

Powder for Solution, Osmotic Laxative

- **Relieves Occasional Constipation/Irregularity**
- **Softens Stool**
- **Dissolves in Any Beverage**
Unflavored Powder Grit Free
Polyethylene Glycol 3350 increases frequency of bowel movements and softens the stool.

NDC 11534-180-19
compare to MiraLAX® active ingredient*

Original Prescription Strength

Polyethylene Glycol

3350

Unflavored Powder
Grit Free

Powder for Solution, Osmotic Laxative

- **Relieves Occasional Constipation (Irregularity)**
- **Softens Stool**
- **Dissolves in Any Beverage**

Polyethylene Glycol 3350 increases frequency of bowel movements and softens the stool.

7 ONCE-DAILY DOSES
NET WT 4.1 OZ (119 g)

SUNRISE PHARMA

Drug Facts

Active ingredient (in each dose)
Polyethylene Glycol, 3350, 17 g (cap filled to line)Osmotic Laxative

Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

Warnings

Allergy alert: Do not use if you are allergic to polyethylene glycol

Do not use if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome


Ask a doctor or pharmacist before use if you are taking a prescription drug

*This product is not manufactured or distributed by Bayer Healthcare LLC, distributor of MiraLAX® **MADE IN USA**

Manufactured & Distributed by:
Sunrise Pharmaceutical, Inc.
Rahway, NJ 07065

Rev. 05/2017
5151/02

● **Sugar Free**



11534-180-19

No Varnish

Peel here

TEMPORARY GLUE

Drug Facts (continued)

When using this product you may have loose, watery, more frequent stools

Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

if pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed unless advised by your doctor
- the bottle top is measuring cap marked to contain 17 grams of powder when filled to the indicated line (clear section in cap)
- adults and children 17 years of age and older:
 - fill to top of clear section in cap which is marked to indicate the correct dose (17g)
- stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
 - use once a day
 - use no more than 7 days
- children 16 years of age or under: ask a doctor

Other information

- store at 20° - 25°C (68° - 77°F)
- tamper-evident: do not use if foil seal under cap, printed with "SEALED FOR YOUR PROTECTION" is missing, open or broken

Inactive ingredients none

Questions or comments?
1-800-FDA-1088

Perm Glue

NDC 11534-180-28

Polyethylene Glycol

3350

Powder for Solution, Osmotic Laxative

- **Relieves Occasional Constipation/Irregularity**
 - **Softens Stool**
 - **Dissolves in Any Beverage**
- Unflavored Powder Grit Free**

Polyethylene Glycol 3350 increases frequency of bowel movements and softens the stool.

NDC 11534-180-28
compare to MiraLAX[®] active ingredient*

Original Prescription Strength

Polyethylene Glycol 3350

Unflavored Powder Grit Free

Powder for Solution, Osmotic Laxative

- Relieves Occasional Constipation (Irregularity)
- Softens Stool
- Dissolves in Any Beverage

Polyethylene Glycol 3350 increases frequency of bowel movements and softens the stool.

SUNRISE PHARMA 14 ONCE-DAILY DOSES
NET WT 8.3 OZ (238 g)

<p>Drug Facts</p> <p>Active ingredient (in each dose) Polyethylene Glycol 3350, 17 g (cap filled to line) Osmotic Laxative</p> <p>Use</p> <ul style="list-style-type: none"> relieves occasional constipation (irregularity) unless advised by your doctor generally produces a bowel movement in 1 to 3 days <p>Warnings</p> <p>Allergy alert: Do not use if you are allergic to polyethylene glycol</p> <p>Do not use if you have kidney disease, except under the advice and supervision of a doctor</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> nausea, vomiting or abdominal pain a sudden change in bowel habits that lasts over 2 weeks irritable bowel syndrome <p>Ask a doctor or pharmacist before use if you are taking a prescription drug</p> <p>When using this product you may have loose, watery, more frequent stools</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition. you get diarrhea you need to use a laxative for longer than 1 week <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Manufactured & Distributed by: Sunrise Pharmaceutical, Inc. Rahway, NJ 07065 *This product is not manufactured or distributed by Bayer Healthcare LLC, distributor of MiraLAX[®]</p> <p>Rev. 05/2017 5152/02</p> <p>MADE IN USA</p> <p>1-800-FDA-1088</p>	<p>Drug Facts (continued)</p> <p>Keep out of reach of children. In case of overdose get medical help or contact a Poison Control Center right away.</p> <p>Directions</p> <ul style="list-style-type: none"> do not take more than directed unless advised by your doctor the bottom cap is measuring cap powder is contained in 17 grams of powder when filled to the indicated line (clear section in cap) adults and children 17 years of age and older: fill to top of clear section in cap which is marked to indicate the correct dose (17g) stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink use once a day use no more than 7 days children 16 years of age or under: ask a doctor <p>Other information</p> <ul style="list-style-type: none"> store at 20° - 25°C (68° - 77°F) tamper-evident: do not use if foil seal under cap, printed with "SEALED FOR YOUR PROTECTION" is missing, open or broken <p>Inactive ingredients none</p> <p>Questions or comments? 1-800-FDA-1088</p>
---	--

3 11534 18028 2

No Varnish

- NDC 11534-180-50
Polyethylene Glycol 3350
Powder for Solution, Osmotic Laxative
- Relieves Occasional Constipation/Irregularity
 - Softens Stool
 - Dissolves in Any Beverage
- Unflavored Powder Grit Free

Polyethylene Glycol 3350 increases frequency of bowel movements and softens the stool.

NDC 11534-180-50
compare to MiraLAX[®] active ingredient*

Original Prescription Strength

Polyethylene Glycol 3350

Unflavored Powder Grit Free

Powder for Solution, Osmotic Laxative

- Relieves Occasional Constipation (Irregularity)
- Softens Stool
- Dissolves in Any Beverage

Polyethylene Glycol 3350 increases frequency of bowel movements and softens the stool.

SUNRISE PHARMA 30 ONCE-DAILY DOSES
NET WT 17.9 OZ (510 g)

<p>Drug Facts</p> <p>Active ingredient (in each dose) Polyethylene Glycol 3350, 17 g (cap filled to line) Osmotic Laxative</p> <p>Use</p> <ul style="list-style-type: none"> relieves occasional constipation (irregularity) unless advised by your doctor generally produces a bowel movement in 1 to 3 days <p>Warnings</p> <p>Allergy alert: Do not use if you are allergic to polyethylene glycol</p> <p>Do not use if you have kidney disease, except under the advice and supervision of a doctor</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> nausea, vomiting or abdominal pain a sudden change in bowel habits that lasts over 2 weeks irritable bowel syndrome <p>Ask a doctor or pharmacist before use if you are taking a prescription drug</p> <p>When using this product you may have loose, watery, more frequent stools</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition. you get diarrhea you need to use a laxative for longer than 1 week <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Manufactured & Distributed by: Sunrise Pharmaceutical, Inc. Rahway, NJ 07065 *This product is not manufactured or distributed by Bayer Healthcare LLC, distributor of MiraLAX[®]</p> <p>Rev. 05/2017 5153/02</p> <p>MADE IN USA</p> <p>1-800-FDA-1088</p>	<p>Drug Facts (continued)</p> <p>Keep out of reach of children. In case of overdose get medical help or contact a Poison Control Center right away.</p> <p>Directions</p> <ul style="list-style-type: none"> do not take more than directed unless advised by your doctor the bottom cap is measuring cap powder is contained in 17 grams of powder when filled to the indicated line (clear section in cap) adults and children 17 years of age and older: fill to top of clear section in cap which is marked to indicate the correct dose (17g) stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink use once a day use no more than 7 days children 16 years of age or under: ask a doctor <p>Other information</p> <ul style="list-style-type: none"> store at 20° - 25°C (68° - 77°F) tamper-evident: do not use if foil seal under cap, printed with "SEALED FOR YOUR PROTECTION" is missing, open or broken <p>Inactive ingredients none</p> <p>Questions or comments? 1-800-FDA-1088</p>
---	--

3 11534 18050 3

No Varnish

POLYETHYLENE GLYCOL 3350 NF

polyethylene glycol 3350 powder, for solution

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:11534-180

Route of Administration	ORAL
--------------------------------	------

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P) (POLYETHYLENE GLYCOL 3350 - UNII:G2M7P15E5P)	POLYETHYLENE GLYCOL 3350	17 g in 17 g

Product Characteristics

Color	WHITE	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11534-180-19	119 g in 1 BOTTLE; Type 0: Not a Combination Product	06/13/2017	
2	NDC:11534-180-28	238 g in 1 BOTTLE; Type 0: Not a Combination Product	06/13/2017	
3	NDC:11534-180-50	510 g in 1 BOTTLE; Type 0: Not a Combination Product	06/13/2017	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA206105	06/13/2017	

Labeler - SUNRISE PHARMACEUTICAL, INC (168522378)

Registrant - SUNRISE PHARMACEUTICAL, INC (168522378)

Establishment

Name	Address	ID/FEI	Business Operations
SUNRISE PHARMACEUTICAL INC.		168522378	MANUFACTURE(11534-180) , ANALYSIS(11534-180) , PACK(11534-180)

Revised: 1/2020

SUNRISE PHARMACEUTICAL, INC