

OXYMORPHONE HYDROCHLORIDE - oxymorphone hydrochloride tablet

Physicians Total Care, Inc.

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use Oxymorphone Hydrochloride Tablets safely and effectively. See full prescribing information for Oxymorphone Hydrochloride Tablets, CII.

Initial U.S. Approval: 1959

----- **INDICATIONS AND USAGE** -----

Oxymorphone Hydrochloride Tablets are an opioid agonist indicated for the relief of moderate to severe acute pain where the use of an opioid is appropriate. (1)

----- **DOSAGE AND ADMINISTRATION** -----

- Titrate the dose base on individual response. (2.1)
- Oxymorphone hydrochloride should be taken on an empty stomach, at least one hour prior to or two hours after eating. (2.2)
- *Opioid-Naïve Patients*: 10 to 20 mg orally every four to six hours.

Initiation of therapy with doses higher than 20 mg is not recommended.

If necessary, may be initiated with 5 mg (e.g., for renal or hepatic impairment, geriatric patients). (2)

- *Conversion to Oxymorphone Hydrochloride Tablets*: Follow recommendations for conversion from other opioids or parenteral oxymorphone. (2.2)
- *Cessation of Therapy*: Taper gradually. (2.4)

----- **DOSAGE FORMS AND STRENGTHS** -----

Tablets: 5 mg and 10 mg. (3)

----- **CONTRAINDICATIONS** -----

- Known hypersensitivity to oxymorphone, any other ingredients in Oxymorphone Hydrochloride Tablets, or morphine analogs (4)
- Respiratory depression in the absence of resuscitative equipment (4)
- Acute or severe bronchial asthma or hypercarbia (4)
- Paralytic ileus (4)
- Moderate or severe hepatic impairment (4)

----- **WARNINGS AND PRECAUTIONS** -----

- Respiratory depression: Increased risk in elderly, debilitated patients, and those suffering from conditions accompanied by hypoxia, hypercapnia, or decreased respiratory reserve. (5.1)
- Misuse, abuse, and diversion: Oxymorphone Hydrochloride is an opioid agonist and a Schedule II controlled substance with an abuse liability similar to morphine. (5.2)
- CNS effects: Additive CNS-depressive effects when used in conjunction with alcohol, other opioids, or illicit drugs. (5.3)
- Head injury: Effects may be markedly exaggerated. Administer with extreme caution. (5.4)
- Hypotensive effect: Increased risk with compromised ability to maintain blood pressure. Administer with caution to patients in circulatory shock. (5.5)
- Mild hepatic impairment: Use with caution and at lower doses due to higher plasma concentrations than in patients with normal hepatic function (5.6)
- Prolonged gastric obstruction: May occur in patients with gastrointestinal obstruction, especially paralytic ileus. (5.8)
- Sphincter of Oddi: Administer with caution in patients with biliary tract disease. (5.9)
- Impaired mental/physical abilities: Caution must be used with potentially hazardous activities. (5.10)

----- **ADVERSE REACTIONS** -----

Adverse reactions ($\geq 2\%$ of patients): Nausea, pyrexia, somnolence, vomiting, pruritus, headache, dizziness, constipation, and confusion. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Roxane Laboratories, Inc. at 1-800-962-8364 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch

----- **DRUG INTERACTIONS** -----

- CNS depressants: Increased risk of respiratory depression, hypotension, profound sedation, coma or death. When combined therapy with CNS depressant is contemplated, the dose of one or both agents should be reduced. (7.1)
- Mixed agonist/antagonist opioids (i.e., pentazocine, nalbuphine, and butorphanol): May reduce analgesic effect and/or precipitate withdrawal symptoms. (7.2)
- Cimetidine: Combination use may precipitate confusion, disorientation, respiratory depression, apnea, seizures. (7.3)

- Anticholinergics: May result in urinary retention and/or severe constipation, which may lead to paralytic ileus. (7.4)
- Monoamine oxidase inhibitors (MAOIs): Potentiate the action of opioids. Oxymorphone Hydrochloride Tablets should not be used in patients taking MAOIs or within 14 days of stopping such treatment. (7.5)

-----**USE IN SPECIFIC POPULATIONS**-----

- Pregnancy: Based on animal data, may cause fetal harm. (8.1)
- Geriatric patients: Oxymorphone Hydrochloride Tablets should be used with caution in elderly patients. (8.5)

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 1/2013

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

- 2.1 Individualization of Dosage
- 2.2 Initiation of Therapy
- 2.3 Maintenance of Therapy
- 2.4 Cessation of Therapy
- 2.5 Patients with Hepatic Impairment
- 2.6 Patients with Renal Impairment
- 2.7 Use with Central Nervous System Depressants
- 2.8 Geriatric Patients

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Respiratory Depression
- 5.2 Misuse, Abuse, and Diversion of Opioids
- 5.3 Additive CNS Depressant Effects
- 5.4 Use in Patients with Head Injury and Increased Intracranial Pressure
- 5.5 Hypotensive Effect
- 5.6 Hepatic Impairment
- 5.7 Special Risk Groups
- 5.8 Gastrointestinal Effects
- 5.9 Use in Pancreatic/Biliary Tract Disease
- 5.10 Driving and Operating Machinery

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience

7 DRUG INTERACTIONS

- 7.1 Use with CNS Depressants
- 7.2 Interactions with Mixed Agonist/Antagonist Opioid Analgesics
- 7.3 Cimetidine
- 7.4 Anticholinergics
- 7.5 MAO Inhibitors

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Labor and Delivery
- 8.3 Nursing Mothers
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Hepatic Impairment
- 8.7 Renal Impairment

9 DRUG ABUSE AND DEPENDENCE

9.1 Controlled Substance

9.2 Abuse

9.3 Dependence

10 OVERDOSAGE

10.1 Symptoms

10.2 Treatment

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

14.1 Orthopedic Surgery

14.2 Abdominal Surgery

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

Oxymorphone Hydrochloride Tablets are indicated for the relief of moderate to severe acute pain where the use of an opioid is appropriate.

2 DOSAGE AND ADMINISTRATION

Selection of patients for treatment with oxymorphone hydrochloride should be governed by the same principles that apply to the use of similar opioid analgesics [see *Indications and Usage (1)*]. Physicians should individualize treatment in every case [see *Dosage and Administration (2.1)*], using non-opioid analgesics, opioids on an as needed basis, combination products, and chronic opioid therapy in a progressive plan of pain management such as outlined by the World Health Organization, the Agency for Healthcare Research and Quality, and the American Pain Society.

Oxymorphone hydrochloride should be administered on an empty stomach, at least one hour prior to or two hours after eating [see *Clinical Pharmacology (12.3)*].

2.1 Individualization of Dosage

As with any opioid drug product, it is necessary to adjust the dosing regimen for each patient individually, taking into account the patient's prior analgesic treatment experience. In the selection of the initial dose of oxymorphone hydrochloride, attention should be given to the following:

- The total daily dose, potency and specific characteristics of the opioid the patient has been taking previously;
- The relative potency estimate used to calculate the equivalent oxymorphone dose needed;
- The patient's degree of opioid tolerance;
- The age, general condition, and medical status of the patient;
- Concurrent non-opioid analgesics and other medications;

- The type and severity of the patient's pain;
- The balance between pain control and adverse experiences;
- Risk factors for abuse or addiction, including a prior history of abuse or addiction.

Once therapy is initiated, frequently assess pain relief and other opioid effects. Titrate dose to adequate pain relief (generally mild or no pain). Patients who experience breakthrough pain may require dosage adjustment.

If signs of excessive opioid-related adverse experiences are observed, the next dose may be reduced. Adjust dosing to obtain an appropriate balance between pain relief and opioid-related adverse experiences. If significant adverse events occur before the therapeutic goal of mild or no pain is achieved, the events should be treated aggressively. Once adverse events are adequately managed, continue upward titration to an acceptable level of pain control.

During periods of changing analgesic requirements, including initial titration, frequent contact is recommended between physician, other members of the healthcare team, the patient, and the caregiver/family. Advise patients and family members of the potential common adverse reactions associated with changing opioid doses.

The dosing recommendations below, therefore, can only be considered as suggested approaches to what is actually a series of clinical decisions over time in the management of the pain of each individual patient.

2.2 Initiation of Therapy

Titrate dose to adequate pain relief (generally mild or no pain).

Opioid-Naïve Patients: Patients who have not been receiving opioid analgesics should be started on oxymorphone hydrochloride in a dosing range of 10 to 20 mg every four to six hours depending on the initial pain intensity. If deemed necessary to initiate therapy at a lower dose (e.g., for renal or hepatic impairment or for geriatric patients), patients may be started with oxymorphone hydrochloride 5 mg. The dose should be titrated based upon the individual patient's response to their initial dose of oxymorphone hydrochloride. This dose can then be adjusted to an acceptable level of analgesia taking into account the pain intensity and adverse reactions experienced by the patient.

Initiation of therapy with doses higher than 20 mg is not recommended because of potential serious adverse reactions [see *Clinical Studies (14.1)*].

Conversion from Parenteral Oxymorphone to Oxymorphone Hydrochloride: Given oxymorphone hydrochloride's absolute oral bioavailability of approximately 10%, patients receiving parenteral oxymorphone may be converted to oxymorphone hydrochloride by administering 10 times the patient's total daily parenteral oxymorphone dose as oxymorphone hydrochloride, in four or six equally divided doses (e.g., [IV dose x 10] divided by 4 or 6). For example, approximately 10 mg of oxymorphone hydrochloride four times daily may be required to provide pain relief equivalent to a total daily IM dose of 4 mg oxymorphone. Due to patient variability with regard to opioid analgesic response, upon conversion patients should be closely monitored to ensure adequate analgesia and to minimize side effects.

Conversion from Other Oral Opioids to Oxymorphone Hydrochloride: For conversion from other opioids to oxymorphone hydrochloride, physicians and other healthcare professionals are advised to refer to published relative potency information, keeping in mind that conversion ratios are only approximate. In general, it is safest to start oxymorphone hydrochloride therapy by administering half of the calculated total daily dose of oxymorphone hydrochloride in 4 to 6 equally divided doses, every 4 to 6 hours. The initial dose of oxymorphone hydrochloride can be gradually adjusted until adequate pain relief and acceptable side effects have been achieved.

2.3 Maintenance of Therapy

During therapy, continual re-evaluation of the patient receiving oxymorphone hydrochloride is important, with special attention to the maintenance of pain control and the relative incidence of side effects associated with therapy. If the level of pain increases, effort should be made to identify the source of increased pain, while adjusting the dose [see *Dosage and Administration (2.1)*].

2.4 Cessation of Therapy

When the patient no longer requires therapy with oxymorphone hydrochloride, doses should be tapered gradually to prevent signs and symptoms of withdrawal in the physically dependent patient [see *Drug Abuse and Dependence (9.3)*].

2.5 Patients with Hepatic Impairment

Oxymorphone hydrochloride is contraindicated in patients with moderate or severe hepatic impairment. Use oxymorphone hydrochloride with caution in patients with mild hepatic impairment, starting with the lowest dose (e.g., 5 mg) and titrating slowly while carefully monitoring side effects [see *Warnings and Precautions (5.6)* and *Clinical Pharmacology (12.3)*].

2.6 Patients with Renal Impairment

There are 57% and 65% increases in oxymorphone bioavailability in patients with moderate and severe renal impairment, respectively; treated with extended-release oxymorphone tablets [see *Clinical Pharmacology (12.3)*]. Accordingly, oxymorphone hydrochloride should be administered cautiously and in reduced dosages to patients with creatinine clearance rates less than 50 mL/min.

2.7 Use with Central Nervous System Depressants

Oxymorphone hydrochloride, like all opioid analgesics, should be started at 1/3 to 1/2 of the usual dose in patients who are concurrently receiving other central nervous system (CNS) depressants including sedatives or hypnotics, general anesthetics, phenothiazines, tranquilizers, and alcohol, because respiratory depression, hypotension and profound sedation, coma or death may result [see *Warnings and Precautions (5.3)* and *Drug Interactions (7.1)*]. When combined therapy with any of the above medications is considered, the dose of one or both agents should be reduced.

Although no specific interaction between oxymorphone and monoamine oxidase inhibitors has been observed, oxymorphone hydrochloride is not recommended for use in patients who have received MAO inhibitors within 14 days [see *Drug Interactions (7.5)*].

2.8 Geriatric Patients

Exercise caution in the selection of the starting dose of oxymorphone hydrochloride for an elderly patient by starting at the low end of the dosing range (e.g., 5 mg) [see *Use in Specific Populations (8.5)*].

3 DOSAGE FORMS AND STRENGTHS

The 5 mg dosage form is a white to off-white, round, standard convex tablet debossed with “54 956” on one side and plain on the other.

The 10 mg dosage form is a white to off-white round, standard convex tablet debossed with “54 814” on one side and plain on the other.

4 CONTRAINDICATIONS

- Oxymorphone Hydrochloride Tablets are contraindicated in patients with a known hypersensitivity to oxymorphone or to any of the other ingredients in Oxymorphone Hydrochloride Tablets, or with known hypersensitivity to morphine analogs such as codeine.

- Oxymorphone Hydrochloride Tablets are contraindicated in patients with respiratory depression, except in monitored settings and in the presence of resuscitative equipment.
- Oxymorphone Hydrochloride Tablets are contraindicated in patients with acute or severe bronchial asthma or hypercarbia.
- Oxymorphone Hydrochloride Tablets are contraindicated in any patient who has or is suspected of having paralytic ileus [see *Warning and Precautions (5.8)*].
- Oxymorphone Hydrochloride Tablets are contraindicated in patients with moderate or severe hepatic impairment [see *Warnings and Precautions (5.6)*].

5 WARNINGS AND PRECAUTIONS

5.1 Respiratory Depression

Respiratory depression is the chief hazard of oxymorphone hydrochloride. Respiratory depression may occur more frequently in elderly or debilitated patients as well as in those suffering from conditions accompanied by hypoxia or hypercapnia, when even moderate therapeutic doses may dangerously decrease pulmonary ventilation.

Administer oxymorphone hydrochloride with extreme caution to patients with conditions accompanied by hypoxia, hypercapnia, or decreased respiratory reserve such as: asthma, chronic obstructive pulmonary disease or cor pulmonale, severe obesity, sleep apnea syndrome, myxedema, kyphoscoliosis, CNS depression, or coma. In these patients, even usual therapeutic doses of oxymorphone may decrease respiratory drive while simultaneously increasing airway resistance to the point of apnea. Consider alternative non-opioid analgesics and use oxymorphone hydrochloride only under careful medical supervision at the lowest effective dose in such patients.

5.2 Misuse, Abuse, and Diversion of Opioids

Oxymorphone hydrochloride contains oxymorphone, a mu opioid agonist and a Schedule II controlled substance with an abuse liability similar to morphine. Opioid agonists are sought by drug abusers and people with addiction disorders and are subject to criminal diversion. Oxymorphone can be abused in a manner similar to other opioid agonists, legal or illicit. This issue should be considered when prescribing or dispensing oxymorphone in situations where the physician or pharmacist is concerned about an increased risk of misuse, abuse, or diversion.

Oxymorphone hydrochloride tablets may be abused by crushing, chewing, snorting, or injecting the product. These practices pose a significant risk to the abuser that could result in overdose and death [see *Drug Abuse and Dependence (9)*].

Oxymorphone hydrochloride may be targeted for theft and diversion. Healthcare professionals should contact their State Medical Board, State Board of Pharmacy, or State Control Board for information on how to detect or prevent diversion of this product, and security requirements for storing and handling of oxymorphone hydrochloride.

Healthcare professionals should advise patients to store oxymorphone hydrochloride in a secure place, preferably locked and out of the reach of children and other noncaregivers.

Concerns about abuse, misuse, diversion and addiction should not prevent the proper management of pain.

5.3 Additive CNS Depressant Effects

The concomitant use of other CNS depressants including other opioids, general anesthetics, phenothiazines, other tranquilizers, sedatives, hypnotics, and alcohol with oxymorphone may produce increased depressant effects including hypoventilation, hypotension, profound sedation, coma and death [see *Drug Interactions (7.1)*].

5.4 Use in Patients with Head Injury and Increased Intracranial Pressure

In the presence of head injury, intracranial lesions or a preexisting increase in intracranial pressure, the respiratory depressant effects of opioid analgesics and their potential to elevate cerebrospinal fluid pressure (resulting from vasodilation following CO₂ retention) may be markedly exaggerated.

Furthermore, opioid analgesics can produce effects on papillary response and consciousness, which may obscure neurologic signs of further increases in intracranial pressure in patients with head injuries.

Administer oxymorphone hydrochloride with extreme caution in patients who may be particularly susceptible to the intracranial effects of CO₂ retention, such as those with evidence of increased intracranial pressure or impaired consciousness. Opioids may obscure the clinical course of a patient with a head injury and should be used only if clinically warranted.

5.5 Hypotensive Effect

Oxymorphone hydrochloride, like all opioid analgesics, may cause severe hypotension in a patient whose ability to maintain blood pressure has been compromised by a depleted blood volume, or after concurrent administration with drugs such as phenothiazines or other agents that compromise vasomotor tone. Administer oxymorphone hydrochloride with caution to patients in circulatory shock, since vasodilation produced by the drug may further reduce cardiac output and blood pressure.

5.6 Hepatic Impairment

A study of extended-release oxymorphone tablets in patients with hepatic disease indicated greater plasma concentrations than in those with normal hepatic function [see *Clinical Pharmacology (12.3)*]. Use oxymorphone hydrochloride with caution in patients with mild impairment, starting with the lowest dose and titrating slowly while carefully monitoring for side effects [see *Dosage and Administration (2.2, 2.5)*]. Oxymorphone hydrochloride is contraindicated in patients with moderate or severe hepatic impairment.

5.7 Special Risk Groups

Use oxymorphone hydrochloride with caution in the following conditions: adrenocortical insufficiency (e.g., Addison's disease), prostatic hypertrophy or urethral stricture, severe impairment of pulmonary or renal function, and toxic psychosis.

Opioids may aggravate convulsions in patients with convulsive disorders, and may induce or aggravate seizures in some clinical settings.

5.8 Gastrointestinal Effects

Opioids diminish propulsive peristaltic waves in the gastrointestinal tract. Monitor for decreased bowel motility in post-operative patients receiving opioids. The administration of oxymorphone hydrochloride may obscure the diagnosis or clinical course in patients with acute abdominal conditions. Oxymorphone hydrochloride is contraindicated in patients with paralytic ileus.

5.9 Use in Pancreatic/Biliary Tract Disease

Oxymorphone hydrochloride, like other opioids, may cause spasm of the sphincter of Oddi and should be used with caution in patients with biliary tract disease, including acute pancreatitis.

5.10 Driving and Operating Machinery

Opioid analgesics impair the mental and physical abilities needed to perform potentially hazardous activities such as driving a car or operating machinery.

6 ADVERSE REACTIONS

The following serious adverse reactions are discussed elsewhere in the labeling:

- Respiratory depression [see Warnings and Precautions (5.1)]
- Misuse and abuse [see Warnings and Precautions (5.2) and Drug Abuse and Dependence (9)]
- CNS depressant effects [see Warnings and Precautions (5.3)]

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

6.1 Clinical Trials Experience

A total of 591 patients were treated with oxymorphone hydrochloride in controlled clinical trials. The clinical trials consisted of patients with acute post-operative pain (n=557) and cancer pain (n=34) trials.

The following table lists adverse reactions that were reported in at least 2% of patients receiving oxymorphone hydrochloride in placebo-controlled trials (acute post-operative pain (N=557)).

Table 1: Adverse Reactions Reported in Placebo-Controlled Trials

	Oxymorphone HCl	Placebo
MedDRA Preferred Term	(N=557)	(N=270)
Nausea	19%	12%
Pyrexia	14%	8%
Somnolence	9%	2%
Vomiting	9%	7%
Pruritus	8%	4%
Headache	7%	4%
Dizziness (Excluding Vertigo)	7%	2%
Constipation	4%	1%
Confusion	3%	<1%

The **common** ($\geq 1\%$ to $>10\%$) adverse drug reactions reported at least once by patients treated with oxymorphone hydrochloride in the clinical trials organized by MedDRA's (Medical Dictionary for Regulatory Activities) System Organ Class were and not represented in Table 1:

Cardiac disorders: tachycardia

Gastrointestinal disorders: dry mouth, abdominal distention, and flatulence

General disorders and administration site conditions: sweating increased

Nervous system disorders: anxiety and sedation

Respiratory, thoracic and mediastinal disorders: hypoxia

Vascular disorders: hypotension

Other **less common** adverse reactions known with opioid treatment that were seen $<1\%$ in the oxymorphone hydrochloride trials includes the following: Abdominal pain, ileus, diarrhea, agitation, disorientation, restlessness, feeling jittery, hypersensitivity, allergic reactions, bradycardia, central nervous system depression, depressed level of consciousness, lethargy, mental impairment, mental status changes, fatigue, depression, clamminess, flushing, hot flashes, dehydration, dermatitis, dyspepsia, dysphoria, edema, euphoric mood, hallucination, hypertension, insomnia, miosis, nervousness, palpitation, postural hypotension, syncope, dyspnea, respiratory depression, respiratory

distress, respiratory rate decreased, oxygen saturation decreased, difficult micturition, urinary retention, urticaria, vision blurred, visual disturbances, weakness, appetite decreased, and weight decreased.

7 DRUG INTERACTIONS

7.1 Use with CNS Depressants

The concomitant use of other CNS depressants including sedatives, hypnotics, tranquilizers, general anesthetics, phenothiazines, other opioids, and alcohol may produce additive CNS depressant effects. Oxymorphone hydrochloride, like all opioid analgesics, should be started at 1/3 to 1/2 of the usual dose in patients who are concurrently receiving other central nervous system depressants because respiratory depression, hypotension, and profound sedation, coma and death may result and titrated slowly as necessary for adequate pain relief.

When combined therapy with any of the above medications is considered, the dose of one or both agents should be reduced [*see Dosage and Administration (2.7) and Warnings and Precautions (5.3)*].

7.2 Interactions with Mixed Agonist/Antagonist Opioid Analgesics

Agonist/antagonist analgesics (i.e., pentazocine, nalbuphine, butorphanol, and buprenorphine) should be administered with caution to a patient who has received or is receiving a course of therapy with a pure opioid agonist analgesic such as oxymorphone. In this situation, mixed agonist/antagonist analgesics may reduce the analgesic effect of oxymorphone and/or may precipitate withdrawal symptoms in these patients.

7.3 Cimetidine

CNS side effects have been reported (e.g., confusion, disorientation, respiratory depression, apnea, seizures) following coadministration of cimetidine with opioid analgesics; a causal relationship has not been established.

7.4 Anticholinergics

Anticholinergics or other medications with anticholinergic activity when used concurrently with opioid analgesics may result in increased risk of urinary retention and/or severe constipation, which may lead to paralytic ileus.

7.5 MAO Inhibitors

Oxymorphone hydrochloride is not recommended for use in patients who have received MAO inhibitors within 14 days, because severe and unpredictable potentiation by MAO inhibitors has been reported with opioid analgesics. No specific interaction between oxymorphone and monoamine oxidase inhibitors has been observed, but caution in the use of any opioid in patients taking this class of drugs is appropriate.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

The safety of using oxymorphone in pregnancy has not been established with regard to possible adverse effects on fetal development. The use of oxymorphone hydrochloride in pregnancy, in nursing mothers, or in women of child-bearing potential requires that the possible benefits of the drug be weighted against the possible hazards to the mother and the child.

Teratogenic Effects:

Pregnancy Category C: There are no adequate and well-controlled studies of oxymorphone in pregnant women. In animal studies, oxymorphone caused decreased fetal and pup weights, an increase in stillbirth, and a decrease in postnatal pup survival at maternal oxymorphone doses equivalent to 0.4 to 4 times the human daily dose of 120 mg (Based on body surface area). Oxymorphone hydrochloride should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

In embryo-fetal developmental toxicity studies, pregnant rats and rabbits received oxymorphone hydrochloride at doses up to about 2 times (rats) and 8 times (rabbits) total human daily dose of 120 mg (based on body surface area). No malformations occurred, but reduced fetal weights occurred at maternal doses of 0.8 (rat) and 4 (rabbit) times the total human daily dose of 120 mg (based on body surface area). There were no adverse developmental effects in rats that received 0.4 times or rabbits that received less than 4 times the total human dose. There were no effects of oxymorphone hydrochloride on intrauterine survival at doses in rats ≤ 2 times, or in rabbits at ≤ 8 times the human dose (see Non-teratogenic Effects, below). In a study conducted prior to the establishment of Good Laboratory Practices (GLP) and not according to current recommended methodology, a single subcutaneous injection of oxymorphone hydrochloride on gestation day 8 produced malformations in offspring of hamsters that received a dose equivalent to 10 times the total human daily dose of 120 mg (based on body surface area). This dose also produced 83% maternal lethality.

Non-teratogenic Effects:

Oxymorphone hydrochloride administration to female rats during gestation in a pre- and postnatal developmental toxicity study reduced mean litter size (18%) at a dose of 25 mg/kg/day, attributed to an increase in the incidence of stillborn pups. An increase in neonatal death occurred at doses ≥ 5 mg/kg/day (0.4 times a total human daily dose of 120 mg, based on body surface area). Low pup birth weight, decreased post-natal weight gain, and reduced post-natal survival of pups occurred following treatment of the dams with 25 mg/kg/day (about 2 times a total human daily dose of 120 mg, based on body surface area).

Prolonged use of opioid analgesics during pregnancy may cause fetal-neonatal physical dependence. Neonatal withdrawal may occur. Symptoms usually appear during the first days of life and may include convulsions, irritability, excessive crying, tremors, hyperactive reflexes, fever, vomiting, diarrhea, sneezing, yawning, and increased respiratory rate.

8.2 Labor and Delivery

Opioids cross the placenta and may produce respiratory depression in neonates. Oxymorphone hydrochloride is not recommended for use in women during and immediately prior to labor, when use of shorter acting analgesics or other analgesic techniques are more appropriate. Occasionally, opioid analgesics may prolong labor through actions which temporarily reduce the strength, duration, and frequency of uterine contractions. However this effect is not consistent and may be offset by an increased rate of cervical dilatation, which tends to shorten labor. Neonates whose mothers received opioid analgesics during labor should be observed closely for signs of respiratory depression. A specific opioid antagonist, such as naloxone or nalmefene, should be available for reversal of opioid-induced respiratory depression in the neonate.

8.3 Nursing Mothers

It is not known whether oxymorphone is excreted in human milk. Because many drugs, including some opioids, are excreted in human milk, caution should be exercised when oxymorphone hydrochloride is administered to a nursing woman. Infants exposed to oxymorphone hydrochloride through breast milk should be monitored for excess sedation and respiratory depression. Withdrawal symptoms can occur in breast-fed infants when maternal administration of an opioid analgesic is stopped, or when breast-feeding is stopped.

8.4 Pediatric Use

Safety and effectiveness of oxymorphone hydrochloride in pediatric patients below the age of 18 years have not been established.

8.5 Geriatric Use

Oxymorphone hydrochloride should be used with caution in elderly patients [*see Clinical Pharmacology (12.3)*]. Of the total number of subjects in clinical studies of oxymorphone hydrochloride, 31% were 65 and over, while 7% were 75 and over. No overall differences in effectiveness were observed between these subjects and younger subjects. There were several adverse events that were more frequently observed in subjects 65 and over compared to younger subjects. These adverse events included dizziness, somnolence, confusion, and nausea. In general, dose selection for elderly patients should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal or cardiac function, and of concomitant disease or other drug therapy

8.6 Hepatic Impairment

In a study of extended-release oxymorphone tablets, patients with mild hepatic impairment were shown to have an increase in bioavailability of 1.6 fold. Oxymorphone hydrochloride should be used with caution in patients with mild impairment. These patients should be started with the lowest dose and titrated slowly while carefully monitoring for side effects. Oxymorphone hydrochloride is contraindicated for patients with moderate and severe hepatic impairment [*see Contraindications (4), Warnings and Precautions (5.6), and Dosage and Administration (2.5)*].

8.7 Renal Impairment

In a study of extended-release oxymorphone tablets, patients with moderate to severe renal impairment were shown to have an increase in bioavailability ranging from 57 to 65% [*see Clinical Pharmacology (12.3)*]. Such patients should be started cautiously with lower doses of oxymorphone hydrochloride and titrated slowly while monitoring for side effects [*see Dosage and Administration (2.6)*].

9 DRUG ABUSE AND DEPENDENCE

9.1 Controlled Substance

Oxymorphone hydrochloride contains oxymorphone, a mu opioid agonist and a Schedule II controlled substance with an abuse liability similar to morphine and other opioids. Oxymorphone can be abused and is subject to criminal diversion [*see Warnings and Precautions (5.2)*].

9.2 Abuse

All patients treated with opioids require careful monitoring for signs of abuse and addiction, since use of opioid analgesic products carries the risk of addiction even under appropriate medical use. Addiction is a primary, chronic, neurobiologic disease, with genetic, psychosocial, and environmental factors influencing its development and manifestations. Addiction is characterized by one or more of the following: impaired control over drug use, compulsive use, use for non-medical purposes, and continued use despite harm. Drug addiction is a treatable disease, utilizing a multidisciplinary approach, but relapse is common.

“Drug-seeking” behavior is very common to addicts and drug abusers. Drug-seeking tactics include emergency calls or visits near the end of office hours, refusal to undergo appropriate examination, testing or referral, repeated claims of loss of prescriptions, tampering with prescriptions, and reluctance to provide prior medical records or contact information for other treating physician(s). “Doctor shopping” (visiting multiple prescribers) to obtain additional prescriptions is common among drug abusers and people suffering from untreated addiction. Preoccupation with achieving adequate pain relief can be appropriate behavior in a patient with poor pain control. Abuse and addiction are

separate and distinct from physical dependence and tolerance. Physicians should be aware that addiction may not be accompanied by concurrent tolerance and symptoms of physical dependence in all addicts. In addition, abuse of opioids can occur in the absence of true addiction and is characterized by misuse for non-medical purposes, often in combination with other psychoactive substances. Oxymorphone hydrochloride, like other opioids, may be diverted for non-medical use. Careful record-keeping of prescribing information, including quantity, frequency, and renewal requests is strongly advised.

Oxymorphone hydrochloride is intended for oral use only. Abuse of oxymorphone hydrochloride poses a risk of overdose and death. This risk is increased with concurrent abuse of oxymorphone hydrochloride with alcohol and other substances. Parenteral drug abuse is commonly associated with transmission of infectious diseases such as hepatitis and HIV.

Proper assessment of the patient, proper prescribing practices, periodic re-evaluation of therapy, and proper dispensing and storage are appropriate measures that help to limit abuse of opioid drugs.

9.3 Dependence

Opioid analgesics may cause physical dependence. Physical dependence results in withdrawal symptoms after abrupt discontinuation of a drug or upon administration of an opioid antagonist or mixed opioid agonist/antagonist agent. Withdrawal also may be precipitated through the administration of drugs with opioid antagonist activity, e.g., naloxone, nalmefene, or mixed agonist/antagonist analgesics (pentazocine, butorphanol, buprenorphine, nalbuphine). Physical dependence may not occur to a clinically significant degree until after several days to weeks of continued opioid usage.

Tolerance is the need for increasing doses of opioids to maintain a defined effect such as analgesia (in the absence of disease progression or other external factors). The development of physical dependence and/or tolerance is not unusual during chronic opioid therapy.

Oxymorphone hydrochloride should not be abruptly discontinued [*see Dosage and Administration (2.4)*]. If oxymorphone hydrochloride is abruptly discontinued in a physically dependent patient, an abstinence syndrome may occur. Some or all of the following can characterize this syndrome: restlessness, lacrimation, rhinorrhea, yawning, perspiration, chills, myalgia, and mydriasis. Other symptoms also may develop, including: irritability, anxiety, backache, joint pain, weakness, abdominal cramps, insomnia, nausea, anorexia, vomiting, diarrhea, or increased blood pressure, respiratory rate, or heart rate.

Infants born to mothers physically dependent on opioids will also be physically dependent and may exhibit respiratory difficulties and withdrawal symptoms [*see Use In Specific Populations (8.1, 8.2)*].

10 OVERDOSAGE

10.1 Symptoms

Acute overdosage with oxymorphone hydrochloride is characterized by respiratory depression (a decrease in respiratory rate and/or tidal volume, Cheyne-Stokes respiration, cyanosis), extreme somnolence progressing to stupor or coma, skeletal muscle flaccidity, cold and clammy skin, constricted pupils, and sometimes bradycardia and hypotension. In some cases, apnea, circulatory collapse, cardiac arrest, and death may occur.

Oxymorphone hydrochloride may cause miosis, even in total darkness. Pinpoint pupils are a sign of opioid overdose but are not pathognomonic (e.g., pontine lesions of hemorrhagic or ischemic origin may produce similar findings). Marked mydriasis rather than miosis may be seen with hypoxia in overdose situations [*see Clinical Pharmacology (12.2)*].

10.2 Treatment

In the treatment of oxymorphone hydrochloride overdosage, primary attention should be given to the re-establishment of a patent airway and institution of assisted or controlled ventilation. Supportive measures (including oxygen and vasopressors) should be employed in the management of circulatory

shock and pulmonary edema accompanying overdose as indicated. Cardiac arrest or arrhythmias may require cardiac massage or defibrillation.

The opioid antagonist naloxone hydrochloride is a specific antidote against respiratory depression that may result from overdosage or unusual sensitivity to opioids including oxymorphone hydrochloride. Nalmefene is an alternative pure opioid antagonist, which may be administered as a specific antidote to respiratory depression resulting from opioid overdose. Since the duration of action of oxymorphone hydrochloride may exceed that of the antagonist, keep the patient under continued surveillance and administer repeated doses of the antagonist according to the antagonist labeling as needed to maintain adequate respiration.


In patients receiving oxymorphone hydrochloride, opioid antagonists should not be administered in the absence of clinically significant respiratory or circulatory depression. Administer opioid antagonists cautiously to persons who are known, or suspected to be, physically dependent on any opioid agonist including oxymorphone hydrochloride. In such cases, an abrupt or complete reversal of opioid effects may precipitate an acute abstinence syndrome. In an individual physically dependent on opioids, administration of the usual dose of the antagonist will precipitate an acute withdrawal syndrome. The severity of the withdrawal syndrome produced will depend on the degree of physical dependence and the dose of the antagonist administered. If respiratory depression is associated with muscular rigidity, administration of a neuromuscular blocking agent may be necessary to facilitate assisted or controlled ventilation. Muscular rigidity may also respond to opioid antagonist therapy.

11 DESCRIPTION

Oxymorphone Hydrochloride Tablets are a semi-synthetic opioid analgesic supplied in 5 mg and 10 mg tablet strengths for oral administration. The tablet strengths describe the amount of oxymorphone hydrochloride per tablet. The tablets contain the following inactive ingredients: magnesium stearate, microcrystalline cellulose, and pregelatinized starch.

Chemically, oxymorphone hydrochloride is 4, 5 α -epoxy-3, 14-dihydroxy-17-methylmorphinan-6-one hydrochloride, a white or slightly off-white, odorless powder, which is sparingly soluble in alcohol and ether, but freely soluble in water. The molecular weight of oxymorphone hydrochloride is 337.80. The pK_{a1} and pK_{a2} of oxymorphone at 37°C are 8.17 and 9.54, respectively. The octanol/aqueous partition coefficient at 37°C and pH 7.4 is 0.98.

The structural formula for oxymorphone hydrochloride is as follows:



12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Oxymorphone, a pure opioid agonist, is relatively selective for the mu receptor, although it can interact with other opioid receptors at higher doses.

The precise mechanism of analgesia, the principal therapeutic action of oxymorphone, is unknown. Specific CNS opiate receptors and endogenous compounds with morphinelike activity have been identified throughout the brain and spinal cord and are likely to play a role in the expression and perception of analgesic effects.

12.2 Pharmacodynamics

Pharmacological effects of opioid agonists include analgesia, anxiolysis, euphoria, feelings of relaxation, respiratory depression, constipation, miosis, and cough suppression. Like all pure opioid agonist analgesics, with increasing doses there is increasing analgesia, unlike with mixed agonist/antagonists or non-opioid analgesics, where there is a limit to the analgesic effect with increasing doses. With pure opioid agonist analgesics, there is no defined maximum dose; the ceiling to analgesic effectiveness is imposed only by side effects, the more serious of which may include somnolence and respiratory depression.

Concentration-Efficacy Relationships: The minimum effective plasma concentration of oxymorphone for analgesia varies widely among patients, especially among patients who have been previously treated with potent agonist opioids. As a result, individually titrate patients to achieve a balance between therapeutic and adverse effects. The minimum effective analgesic concentration of oxymorphone for any individual patient may increase over time due to an increase in pain, progression of disease, development of a new pain syndrome and/or development of analgesic tolerance.

Concentration-Adverse Experience Relationships: There is a general relationship between increasing opioid plasma concentration and increasing frequency of adverse experiences such as nausea, vomiting, CNS effects, and respiratory depression.

As with all opioids, the dose of oxymorphone hydrochloride must be individualized [*see Dosage and Administration (2.2)*]. The effective analgesic dose for some patients will be too high to be tolerated by other patients.

Effects on the Central Nervous System (CNS): The principal therapeutic action of oxymorphone is analgesia. In common with other opioids, oxymorphone causes respiratory depression, in part by a direct effect on the brainstem respiratory centers. The respiratory depression involves a reduction in the responsiveness of the brain stem respiratory centers to both increases in carbon dioxide tension and electrical stimulation. Opioids depress the cough reflex by direct effect on the cough center in the medulla. Antitussive effects may occur with doses lower than those usually required for analgesia. Oxymorphone causes miosis, even in total darkness. Pinpoint pupils are a sign of opioid overdose but are not pathognomonic (e.g., pontine lesions of hemorrhagic or ischemic origin may produce similar findings). Marked mydriasis rather than miosis may be seen with hypoxia in overdose situations [*see Overdosage (10.1)*]. Other therapeutic effects of oxymorphone include anxiolysis, euphoria and feelings of relaxation.

In addition to analgesia, the widely diverse effects of oxymorphone include drowsiness, changes in mood, decreased gastrointestinal motility, nausea, vomiting, and alterations of the endocrine and autonomic nervous system.

Effects on the Gastrointestinal Tract and on Other Smooth Muscle: Gastric, biliary and pancreatic secretions are decreased by oxymorphone. Oxymorphone causes a reduction in motility and is associated with an increase in tone in the antrum of the stomach and duodenum. Digestion of food in the small intestine is delayed and propulsive contractions are decreased. Propulsive peristaltic waves in the colon are decreased, while tone is increased to the point of spasm. The end result may be constipation. Oxymorphone can cause a marked increase in biliary tract pressure as a result of spasm of the sphincter of Oddi. Oxymorphone may also cause spasm of the sphincter of the urinary bladder.

Cardiovascular System Effects: Opioids produce peripheral vasodilation which may result in orthostatic

hypotension. Release of histamine can occur and may contribute to opioid-induced hypotension. Manifestations of histamine release may include orthostatic hypotension, pruritus, flushing, red eyes, and sweating. Animal studies have shown that oxymorphone has a lower propensity to cause histamine release than other opioids.

Endocrine System Effects: Opioid agonists have been shown to have a variety of effects on the secretion of hormones. Opioids inhibit the secretion of ACTH, cortisol, and luteinizing hormone (LH) in humans. They also stimulate prolactin, growth hormone (GH) secretion, and pancreatic secretion of insulin and glucagon in humans and other species, rats and dogs. Thyroid stimulating hormone (TSH) has been shown to be both inhibited and stimulated by opioids.

Immune System Effects: Opioids have been shown to have a variety of effects on components of the immune system in *in vitro* and animal models. The clinical significance of these findings is unknown.

12.3 Pharmacokinetics

Absorption: The absolute oral bioavailability of oxymorphone is approximately 10%. Studies in healthy volunteers reveal predictable relationships between oxymorphone hydrochloride dosage and plasma oxymorphone concentrations.

Steady-state levels were achieved after three days of multiple dose administration. Under both single-dose and steady-state conditions, dose proportionality has been established for 5 mg, 10 mg and 20 mg doses of oxymorphone hydrochloride, for both peak plasma levels (C_{max}) and extent of absorption (AUC) (see Table 2).

Table 2: Mean (\pm SD) Oxymorphone Hydrochloride Pharmacokinetic Parameters

	Dosage	C_{max} (ng/mL)	AUC (ng•hr/mL)	$T_{1/2}$ (hr)
Single Dose	5 mg	1.1 \pm 0.55	4.48 \pm 2.07	7.25 \pm 4.4
	10 mg	1.93 \pm 0.75	9.1 \pm 3.4	7.78 \pm 3.56
	20 mg	4.39 \pm 1.72	20.07 \pm 5.8	9.43 \pm 3.36
Multiple Dose	5 mg	1.73 \pm 0.62	4.63 \pm 1.49	NA
	10 mg	3.51 \pm 0.91	10.19 \pm 3.34	NA
	20 mg	7.33 \pm 2.93	21.1 \pm 7.59	NA

Food Effect: After oral dosing with 40 mg of oxymorphone hydrochloride in healthy volunteers under fasting conditions or with a high-fat meal, the C_{max} and AUC were increased by approximately 38% in fed subjects relative to fasted subjects. As a result, oxymorphone hydrochloride should be dosed at least one hour prior to or two hours after eating [see *Dosage and Administration (2)*].

Ethanol Effect: The effect of co-ingestion of alcohol with oxymorphone hydrochloride has not been evaluated. However, an *in vivo* study was performed to evaluate the effect of alcohol (40%, 20%, 4% and 0%) on the bioavailability of a single dose of 40 mg of extended-release oxymorphone tablets in healthy, fasted volunteers. Following concomitant administration of 240 mL of 40% ethanol the C_{max} increased on average by 70% and up to 270% in individual subjects. Following the concomitant administration of 240 mL of 20% ethanol, the C_{max} increased on average by 31% and up to 260% in individual subjects. In some individuals there was also a decrease in oxymorphone peak plasma concentrations. No effect on the release of oxymorphone from the extended-release tablet was noted in an *in vitro* alcohol interaction study. The mechanism of the *in vivo* interaction is unknown. Therefore, avoid co-administration of oxymorphone and ethanol.

Distribution: Formal studies on the distribution of oxymorphone in various tissues have not been conducted. Oxymorphone is not extensively bound to human plasma proteins; binding is in the range of 10% to 12%.

Metabolism: Oxymorphone is highly metabolized, principally in the liver, and undergoes reduction or conjugation with glucuronic acid to form both active and inactive products. The two major metabolites of oxymorphone are oxymorphone-3-glucuronide and 6-OH-oxymorphone. The mean plasma AUC for oxymorphone-3-glucuronide is approximately 90-fold higher than the parent compound. The pharmacologic activity of the glucuronide metabolite has not been evaluated. 6-OH-oxymorphone has been shown in animal studies to have analgesic bioactivity. The mean plasma 6-OH-oxymorphone AUC is approximately 70% of the oxymorphone AUC following single oral doses but is essentially equivalent to the parent compound at steady-state.

Excretion: Because oxymorphone is extensively metabolized, <1% of the administered dose is excreted unchanged in the urine. On average, 33% to 38% of the administered dose is excreted in the urine as oxymorphone-3-glucuronide and 0.25% to 0.62% is excreted as 6-OH-oxymorphone in subjects with normal hepatic and renal function. In animals given radiolabeled oxymorphone, approximately 90% of the administered radioactivity was recovered within 5 days of dosing. The majority of oxymorphone-derived radioactivity was found in the urine and feces.

Pharmacokinetics in Special Populations: Elderly: The plasma levels of oxymorphone administered as an extended-release tablet were about 40% higher in elderly (≥ 65 years of age) than in younger subjects [see *Use in Specific Populations (8.5)*].

Gender: The effect of gender on the pharmacokinetics of oxymorphone hydrochloride has not been studied. In a study with an extended-release formulation of oxymorphone, there was a consistent tendency for female subjects to have slightly higher AUC_{ss} and C_{max} values than male subjects. However, gender differences were not observed when AUC_{ss} and C_{max} were adjusted by body weight.

Hepatic Impairment: The liver plays an important role in the pre-systemic clearance of orally administered oxymorphone. Accordingly, the bioavailability of orally administered oxymorphone may be markedly increased in patients with moderate to severe liver disease. The effect of hepatic impairment on the pharmacokinetics of oxymorphone hydrochloride has not been studied. However, in a study with an extended-release formulation of oxymorphone, the disposition of oxymorphone was compared in 6 patients with mild, 5 patients with moderate, and one patient with severe hepatic impairment, and 12 subjects with normal hepatic function. The bioavailability of oxymorphone was increased by 1.6-fold in patients with mild hepatic impairment and by 3.7-fold in patients with moderate hepatic impairment. In one patient with severe hepatic impairment, the bioavailability was increased by 12.2-fold. The half-life of oxymorphone was not significantly affected by hepatic impairment.

Renal Impairment: The effect of renal impairment on the pharmacokinetics of oxymorphone hydrochloride have not been studied. However, in a study with an extended-release formulation of oxymorphone, an increase of 26%, 57%, and 65% in oxymorphone bioavailability was observed in mild (creatinine clearance 51 to 80 mL/min; n=8), moderate (creatinine clearance 30 to 50 mL/min; n=8), and severe (creatinine clearance <30 mL/min; n=8) patients, respectively, compared to healthy controls.

Drug-Drug Interactions: *In vitro* studies revealed little to no biotransformation of oxymorphone to 6-OH-oxymorphone by any of the major cytochrome P450 (CYP P450) isoforms at therapeutically relevant oxymorphone plasma concentrations.

No inhibition of any of the major CYP P450 isoforms was observed when oxymorphone was incubated with human liver microsomes at concentrations of ≤ 50 μ M. An inhibition of CYP 3A4 activity occurred at oxymorphone concentrations ≥ 150 μ M. Therefore, it is not expected that oxymorphone, or its metabolites will act as inhibitors of any of the major CYP P450 enzymes *in vivo*.

Increases in the activity of the CYP 2C9 and CYP 3A4 isoforms occurred when oxymorphone was incubated with human hepatocytes. However, clinical drug interaction studies with oxymorphone hydrochloride ER showed no induction of CYP450 3A4 or 2C9 enzyme activity, indicating that no dose adjustment for CYP 3A4- or 2C9-mediated drug-drug interactions is required.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis: Long-term studies have been completed to evaluate the carcinogenic potential of oxymorphone in both Sprague-Dawley rats and CD-1 mice. Oxymorphone was administered to Sprague-Dawley rats (2.5, 5, and 10 mg/kg/day in males and 5, 10, and 25 mg/kg/day in females) for 2 years by oral gavage. The systemic drug exposure (AUC ng•h/mL) at the 10 mg/kg/day dose in male rats was 0.34-fold and at the 25 mg/kg/day dose in female rats was 1.5-fold the human exposure at a dose of 260 mg/day. No evidence of carcinogenic potential was observed in rats. Oxymorphone was administered to CD-1 mice (10, 25, 75 and 150 mg/kg/day) for 2 years by oral gavage. The systemic drug exposure (AUC ng•h/mL) at the 150 mg/kg/day dose in mice was 14.5-fold (in males) and 17.3-fold (in females) times the human exposure at a dose of 260 mg/day. No evidence of carcinogenic potential was observed in mice.

Mutagenesis: Oxymorphone hydrochloride was not mutagenic when tested in the *invitro* bacterial reverse mutation assay (Ames test) at concentrations of ≤ 5270 $\mu\text{g}/\text{plate}$, or in an *in vitro* mammalian cell chromosome aberration assay performed with human peripheral blood lymphocytes at concentrations ≤ 5000 $\mu\text{g}/\text{ml}$ with or without metabolic activation. Oxymorphone hydrochloride tested positive in both the rat and mouse *in vivo* micronucleus assays. An increase in micronucleated polychromatic erythrocytes occurred in mice given doses of ≥ 250 mg/kg and in rats given doses of 20 and 40 mg/kg. A subsequent study demonstrated that oxymorphone hydrochloride was not aneugenic in mice following administration of up to 500 mg/kg. Additional studies indicate that the increased incidence of micronucleated polychromatic erythrocytes in rats may be secondary to increased body temperature following oxymorphone administration. Doses associated with increased micronucleated polychromatic erythrocytes also produce a marked, rapid increase in body temperature. Pretreatment of animals with sodium salicylate minimized the increase in body temperature and prevented the increase in micronucleated polychromatic erythrocytes after administration of 40 mg/kg oxymorphone.

Impairment of fertility: Oxymorphone did not affect reproductive function or sperm parameters in male rats at any dose tested (≤ 50 mg/kg/day). In female rats, an increase in the length of the estrus cycle and decrease in the mean number of viable embryos, implantation sites and corpora lutea were observed at doses of oxymorphone ≥ 10 mg/kg/day. The dose of oxymorphone associated with reproductive findings in female rats is 0.8 times a total human daily dose of 120 mg based on a body surface area. The dose of oxymorphone that produced no adverse effects on reproductive findings in female rats (i.e. NNOAEL) is 0.4-times a total human daily dose of 120 mg based on body surface area.

14 CLINICAL STUDIES

The analgesic efficacy of oxymorphone hydrochloride has been evaluated in acute pain following orthopedic and abdominal surgeries.

14.1 Orthopedic Surgery

Two double-blind, placebo-controlled, dose-ranging studies in patients with acute moderate to severe pain following orthopedic surgery evaluated the doses of oxymorphone hydrochloride 10 mg and 20 mg, and 30 mg was included in one study. Both studies demonstrated that oxymorphone hydrochloride 20 mg provided greater analgesia as measured by total pain relief based on a weighted analysis over 8 hours using a 0-4 categorical, compared to placebo. Oxymorphone hydrochloride 10 mg provided greater analgesia as compared to placebo in one of the two studies. There was no evidence of superiority of the 30 mg dose over the 20 mg dose. However, there was a high rate of naloxone use in patients receiving the oxymorphone hydrochloride 30 mg dose in the post-operative period [see *Dosage and Administration* (2.2)].

14.2 Abdominal Surgery

In a randomized, double-blind, placebo-controlled, multiple-dose study, the efficacy of oxymorphone hydrochloride 10 mg and 20 mg was assessed in patients with moderate to severe acute pain following abdominal surgery. In this study, patients were dosed every 4 to 6 hours over a 48-hour treatment period. Oxymorphone hydrochloride 10 and 20 mg provided greater analgesia, as measured by the mean average pain intensity on a 0-100 mm visual analog scale, over 48 hours, compared to placebo [see *Dosage and Administration (2.2)*].

16 HOW SUPPLIED/STORAGE AND HANDLING

Oxymorphone Hydrochloride Tablets are supplied as follows:

10 mg Tablet:

White to off-white, round, standard convex debossed with “54 814” on one side and plain on the other.

Bottles of	NDC 54868-
60	6379-0

Oxymorphone hydrochloride may be targeted for theft and diversion. Healthcare professionals should contact their State Medical Board, State Board of Pharmacy, or State Control Board for information on how to detect or prevent diversion of this product, and security requirements for storing and handling of oxymorphone hydrochloride.

Healthcare professionals should advise patients to store oxymorphone hydrochloride in a secure place, preferably locked and out of the reach of children and other noncaregivers.

Store at 25°C (77°F); excursions permitted to 15° to 30°C (59° to 86°F). [See USP Controlled Room Temperature].

Dispense in tight container as defined in the USP, with a child-resistant closure (as required).

Advise patients to dispose of any unused tablets from a prescription by flushing them down the toilet as soon as they are no longer needed [see *Patient Counseling Information (17)*].

17 PATIENT COUNSELING INFORMATION

- Advise patients that oxymorphone hydrochloride contains oxymorphone, which is a morphine-like pain reliever, and should be taken only as directed.
- Advise patients that oxymorphone hydrochloride is a potential drug of abuse. They should protect it from theft, and it should never be given to anyone other than the individual for whom it was prescribed.
- Advise patients to keep oxymorphone hydrochloride in a secure place out of the reach of children and pets. Accidental consumption especially in children may result in overdose or death. When oxymorphone hydrochloride is no longer needed, the unused tablets should be destroyed by flushing down the toilet.
- Advise patients to report episodes of breakthrough pain and adverse experiences occurring during therapy to their doctor. Individualization of dosage is essential to make optimal use of this medication.
- Advise patients not to adjust the dose of oxymorphone hydrochloride without consulting the prescriber.
- Advise patients that oxymorphone hydrochloride may cause drowsiness, dizziness, or

lightheadedness and may impair mental and/or physical abilities required for the performance of potentially hazardous tasks, such as driving a car, operating machinery, etc.

- Advise patients that oxymorphone hydrochloride will add to the effect of alcohol and other CNS depressants (such as antihistamines, sedatives, hypnotics, tranquilizers, general anesthetics, phenothiazines, other opioids, and monoamine oxidase [MAO] inhibitors).
- Advise patients not to combine oxymorphone hydrochloride with alcohol or other central nervous system depressants (sleep aids, tranquilizers) except by the orders of the prescribing physician, because dangerous additive effects may occur, resulting in serious injury or death
- Advise patients of the potential for severe constipation. Appropriate laxatives and/or stool softeners and other therapeutic approaches should be considered for use with the initiation of oxymorphone hydrochloride therapy.
- Advise women of childbearing potential who become or are planning to become pregnant to consult their physician regarding the effects of opioid analgesics and other drug use during pregnancy on themselves and their unborn child.
- Advise women of childbearing potential who become or are planning to become pregnant that safe use in pregnancy has not been established. Prolonged use of opioid analgesics during pregnancy may cause fetal-neonatal physical dependence, and neonatal withdrawal may occur
- Advise patients that if they have been receiving treatment with oxymorphone hydrochloride for more than a few weeks and cessation of therapy is indicated, it may be appropriate to taper the oxymorphone hydrochloride dose, rather than abruptly discontinue it, due to the risk of precipitating withdrawal symptoms. Their physician can provide a dose schedule to accomplish a gradual discontinuation of the medication.

DEA Order Form Required.

10005444/02 Revised May 2010

© RLI, 2010

Distributed and Repackaged by:

Physicians Total Care, Inc.
Tulsa, Oklahoma 74146

PRINCIPAL DISPLAY PANEL

Oxymorphone Hydrochloride Tablets

10 mg

PHYSICIANS TOTAL CARE
NDC 54868-6379-8
OXYMORPHONE HYDROCHLORIDE
OXYMORPHONE HCL TABLETS, USP
(WARNING: MAY BE HABIT FORMING)
10 MG
60 TABLETS
CAUTION: Federal law prohibits dispensing without prescription.

Each Tablet Contains:
OXYMORPHONE HCL, USP 10 mg
(WARNING: MAY BE HABIT FORMING)

Usual Dosage & Warnings:
Read Carefully Accompanying Literature

KEEP THIS AND ALL DRUGS OUT OF THE REACH OF CHILDREN.

DISPENSE CONTENTS IN CHILD-RESISTANT CLOSURE (AS REQUIRED) AND IN TIGHT, LIGHT-RESISTANT CONTAINER AS DEFINED IN THE USP/NF.

STORE AT CONTROLLED ROOM TEMPERATURE 20 - 25 °C (68 - 77 °F) (20 - 25 °C).

6379060000
LOT: 000 EXP. 01/10
MFG. BY: **ORANGE LABORATORIES, INC.**
COLUMBUS OH 43216

Repackaged & distributed by: **Physicians Total Care, Inc.**
Tulsa, OK 74146

OXYMORPHONE HYDROCHLORIDE

oxymorphone hydrochloride tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:54868- 6379(NDC:0054-0284)
Route of Administration	ORAL	DEA Schedule	CII

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OXYMORPHONE HYDROCHLORIDE (UNII: 5Y2EI94NBC) (OXYMORPHONE - UNII:9VXA968E0C)	OXYMORPHONE HYDROCHLORIDE	10 mg

Inactive Ingredients

Ingredient Name	Strength
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
STARCH, CORN (UNII: O8232NY3SJ)	

Product Characteristics

Color	WHITE	Score	no score
Shape	ROUND	Size	8mm
Flavor		Imprint Code	54;814
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:54868-6379-0	60 in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA090964	12/10/2012	

Labeler - Physicians Total Care, Inc. (194123980)

Establishment

Name	Address	ID/FEI	Business Operations
------	---------	--------	---------------------

Revised: 1/2013

Physicians Total Care, Inc.