

**MAGNESIUM SULFATE IN 5% DEXTROSE- magnesium sulfate heptahydrate injection, solution**  
**Fresenius Kabi USA, LLC**

-----  
**Magnesium Sulfate in 5% Dextrose Injection, USP**

**For Intravenous Use Only**


Rx only

**DESCRIPTION**

Magnesium Sulfate in 5% Dextrose Injection, USP is a sterile, nonpyrogenic solution of magnesium sulfate heptahydrate and dextrose in water for injection. Each 100 mL contains 1 g magnesium sulfate heptahydrate and dextrose, hydrous 5 g in water for injection. May contain sulfuric acid and/or sodium hydroxide for pH adjustment. The pH is 4.5 (3.5 to 6.5). It is available in a 1% concentration. See **HOW SUPPLIED** section for the content and characteristics of available dosage form and size.

Magnesium sulfate, USP heptahydrate is chemically designated  $MgSO_4 \cdot 7H_2O$ , colorless crystals or white powder freely soluble in water.

Dextrose, USP is chemically designated D-glucose, monohydrate, a hexose sugar freely soluble in water. It has the following structural formula:



$C_6H_{12}O_6 \cdot H_2O$  M.W. 198.17

Water for injection, USP is chemically designated  $H_2O$ .

The flexible container is fabricated from a specially formulated non-plasticized, film containing polypropylene and thermoplastic elastomers ( **freeflex**<sup>®</sup> bag). The amount of water that can permeate from the container into the overwrap is insufficient to affect the solution significantly. Solutions in contact with the flexible container can leach out certain of the container's chemical components in very small amounts within the expiration period. The suitability of the container material has been confirmed by tests in animals according to USP biological tests for plastic containers. Exposure to temperatures above 25°C/77°F during transport and storage will lead to minor losses in moisture content. Higher temperatures lead to greater losses. It is unlikely that these minor losses will lead to clinically significant changes within the expiration period.

**CLINICAL PHARMACOLOGY**

Magnesium ( $Mg^{++}$ ) is an important cofactor for enzymatic reactions and plays an important role in neurochemical transmission and muscular excitability.

Magnesium prevents or controls convulsions by blocking neuromuscular transmission and decreasing the amount of acetylcholine liberated at the end plate by the motor

nerve impulse. Magnesium is said to have a depressant effect on the central nervous system, but it does not adversely affect the mother, fetus or neonate when used as directed in eclampsia or pre-eclampsia. Normal serum magnesium levels range from 1.3 to 2.1 mEq/liter.

As serum magnesium rises above 4 mEq/liter, the deep tendon reflexes are first decreased and then disappear as the serum level approaches 10 mEq/liter. At this level respiratory paralysis may occur. Heart block also may occur at this or lower serum levels of magnesium.

Magnesium acts peripherally to produce vasodilation. With low doses only flushing and sweating occur, but larger doses cause lowering of blood pressure. The central and peripheral effects of magnesium poisoning are antagonized to some extent by intravenous administration of calcium.

With intravenous administration the onset of anticonvulsant action is immediate and lasts about 30 minutes. Following intramuscular administration the onset of action occurs in about one hour and persists for three to four hours. Effective anticonvulsant serum levels range from 2.5 to 7.5 mEq/liter.

## **Pharmacokinetics**

### **Absorption**

Intravenously administered magnesium is immediately absorbed.

### **Distribution**

Approximately 1 to 2% of total body magnesium is located in the extracellular fluid space. Magnesium is 30% bound to albumin.

### **Metabolism**

Magnesium is not metabolized.

### **Excretion**

Magnesium is excreted solely by the kidney at a rate proportional to the serum concentration and glomerular filtration.

## **Special Populations**

### **Renal Insufficiency**

Magnesium is excreted solely by the kidney. In patients with severe renal insufficiency, the dose should be lower and frequent serum magnesium levels must be obtained (see **DOSAGE AND ADMINISTRATION**).

### **Hepatic Insufficiency**

Magnesium is excreted solely by the kidney. No dosing adjustments are necessary in hepatic insufficiency.

## **Drug-Drug Interactions**

Drug induced renal losses of magnesium occur with the following drugs or drug classes:

Aminoglycosides	Amphotericin B
Cyclosporine	Diuretics
Digitalis	Cisplatin
Alcohol	

## **INDICATIONS AND USAGE**

Magnesium sulfate in 5% dextrose injection, USP is indicated for the prevention and control of seizures in pre-eclampsia and eclampsia, respectively. When used judiciously it effectively prevents and controls the convulsions of eclampsia without producing deleterious depression of the central nervous system of the mother or infant. However, other effective drugs are available for this purpose.

## **CONTRAINDICATIONS**

Intravenous magnesium should not be given to mothers with toxemia of pregnancy during the two hours preceding delivery.

## **WARNINGS**

**FETAL HARM:** Continuous administration of magnesium sulfate beyond 5 to 7 days to pregnant women can lead to hypocalcemia and bone abnormalities in the developing fetus. These bone abnormalities include skeletal demineralization and osteopenia. In addition, cases of neonatal fracture have been reported. The shortest duration of treatment that can lead to fetal harm is not known. Magnesium sulfate should be used during pregnancy only if clearly needed. If magnesium sulfate is given for treatment of preterm labor, the woman should be informed that the efficacy and safety of such use have not been established and that use of magnesium sulfate beyond 5 to 7 days may cause fetal abnormalities.

Parenteral use in the presence of renal insufficiency may lead to magnesium intoxication.

## **PRECAUTIONS**

Because magnesium is removed from the body solely by the kidneys, the drug should be used with caution in patients with renal impairment. Urine output should be maintained at a level of 100 mL every four hours. Monitoring serum magnesium levels and the patient's clinical status is essential to avoid the consequences of overdosage in toxemia. Clinical indications of a safe dosage regimen include the presence of the patellar reflex (knee jerk) and absence of respiratory depression (approximately 16 breaths or more/minute). Serum magnesium levels usually sufficient to control convulsions range from 3 to 6 mg/100 mL (2.5 to 5 mEq/liter). The strength of the deep tendon reflexes begins to diminish when serum magnesium levels exceed 4 mEq/liter.

Reflexes may be absent at 10 mEq magnesium/liter, where respiratory paralysis is a potential hazard. An injectable calcium salt should be immediately available to counteract the potential hazards of magnesium intoxication in eclampsia.

Magnesium sulfate in 5% dextrose injection, USP should be administered slowly to avoid producing hypermagnesemia.

### **Carcinogenesis, Mutagenesis, Impairment of Fertility**

Studies with magnesium sulfate in 5% dextrose injection, USP have not been performed to evaluate carcinogenic potential, mutagenic potential or effects on fertility.

### **Pregnancy**

#### **Teratogenic Effects**

**Pregnancy Category D (see WARNINGS and PRECAUTIONS)**

See **WARNINGS** and **PRECAUTIONS**.

Magnesium sulfate in 5% dextrose injection, USP can cause fetal abnormalities when administered beyond 5 to 7 days to pregnant women. There are retrospective epidemiological studies and case reports documenting fetal abnormalities such as hypocalcemia, skeletal demineralizations, osteopenia and other skeletal abnormalities with continuous maternal administration of magnesium sulfate for more than 5 to 7 days.<sup>1-12</sup> Magnesium sulfate in 5% dextrose injection, USP should be used during pregnancy only if clearly needed. If this drug is used during pregnancy the woman should be apprised of the potential harm to the fetus.

### **Nonteratogenic Effects**

When administered by continuous IV infusion (especially for more than 24 hours preceding delivery) to control convulsions in a toxemic woman, the newborn may show signs of magnesium toxicity, including neuromuscular or respiratory depression (see **OVERDOSAGE**).

### **Labor and Delivery**

Continuous administration of magnesium sulfate is an unapproved treatment for preterm labor. The safety and efficacy of such use have not been established. The administration of magnesium sulfate in 5% dextrose injection, USP outside of its approved indication in pregnant women should be by trained obstetrical personnel in a hospital setting with appropriate obstetrical care facilities.

### **Nursing Mothers**

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when magnesium sulfate in 5% dextrose injection, USP is administered to a nursing woman.

## **ADVERSE REACTIONS**

The adverse effects of parenterally administered magnesium usually are the result of magnesium intoxication. These include flushing, sweating, hypotension, depressed reflexes, flaccid paralysis, hypothermia, circulatory collapse, cardiac and central nervous system depression proceeding to respiratory paralysis. Hypocalcemia with signs of tetany secondary to magnesium sulfate therapy for eclampsia has been reported.

## **OVERDOSAGE**

Magnesium intoxication is manifested by a sharp drop in blood pressure and respiratory paralysis. Disappearance of the patellar reflex is a useful clinical sign to detect the onset of magnesium intoxication. In the event of overdose, artificial ventilation must be provided until a calcium salt can be injected IV to antagonize the effects of magnesium.

### **For Treatment of Overdose**

Artificial respiration is often required. Intravenous calcium, 10 to 20 mL of a 5% solution (diluted if desirable) with isotonic sodium chloride for injection) is used to counteract effects of hypermagnesemia. Subcutaneous physostigmine, 0.5 to 1 mg may be helpful.

Hypermagnesemia in the newborn may require resuscitation and assisted ventilation via endotracheal intubation or intermittent positive pressure ventilation as well as IV calcium.

## DOSAGE AND ADMINISTRATION

Magnesium sulfate in 5% dextrose injection, USP is intended for intravenous use only. For the management of pre-eclampsia or eclampsia, intravenous infusions of dilute solutions of magnesium (1% to 8%) are often given in combination with intramuscular injections of 50% magnesium sulfate injection, USP. Therefore, in the clinical conditions cited below, both forms of therapy are noted, as appropriate. Continuous maternal administration of magnesium sulfate in pregnancy beyond 5 to 7 days can cause fetal abnormalities.

### In Pre-eclampsia or Eclampsia

In severe pre-eclampsia or eclampsia, the total initial dose is 10 to 14 g of magnesium sulfate. To initiate therapy, 4 g of magnesium sulfate in 5% dextrose injection, USP may be administered intravenously. The rate of IV infusion should generally not exceed 150 mg/minute, or 7.5 mL of a 2% concentration (or its equivalent) per minute, except in severe eclampsia with seizures. Simultaneously, 4 to 5 g (32.5 to 40.6 mEq) of magnesium sulfate may be administered intramuscularly into each buttock using undiluted 50% magnesium sulfate injection, USP. After the initial IV dose, some clinicians administer 1 to 2 g/hour by constant IV infusion.

Subsequent intramuscular doses of 4 to 5 g of magnesium sulfate may be injected into alternate buttocks every four hours, depending on the continuing presence of the patellar reflex, adequate respiratory function, and absence of signs of magnesium toxicity. Therapy should continue until paroxysms cease.

A serum magnesium level of 6 mg/100 mL is considered optimal for control of seizures. A total daily (24 hr) dose of 30 to 40 g magnesium sulfate should not be exceeded. In the presence of severe renal insufficiency, frequent serum magnesium concentrations must be obtained, and the maximum recommended dosage of magnesium sulfate is 20 g per 48 hours.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit.

Do not administer unless solution is clear. Discard unused portion.

## HOW SUPPLIED

Magnesium Sulfate in 5% Dextrose Injection, USP is supplied in a single-dose flexible plastic container as follows:

Product Code	Unit of Sale	Total Magnesium Sulfate*	Total Magnesium Ion	Magnesium Sulfate* Concentration	Magnesium Ion Concentration	Osmolarity (calc.)	Unit of Use
610801	NDC 63323-108-01 Packages of 24	1 g	8.1 mEq	1% (10 mg/mL)	8.1 mEq/100 mL	333 mOsmol/Liter	NDC 63323-108-00 100 mL fill in a 100 mL <b>freeflex</b> Bag

\*As the heptahydrate.

**WARNING: DO NOT USE FLEXIBLE CONTAINER IN SERIES CONNECTIONS.**

The container closure is not made with natural rubber latex. Non-PVC, Non-DEHP, Sterile.

Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Protect from freezing.

## REFERENCES

1. Yokoyama K, Takahashi N, Yada Y. Prolonged maternal magnesium administration and bone metabolism in neonates. *Early Human Dev.* 2010; 86(3):187-91. Epub 2010 Mar 12.
2. Wedig KE, Kogan J, Schorry EK et al. Skeletal demineralization and fractures caused by fetal magnesium toxicity. *J Perinatol.* 2006; 26(6):371-4.
3. Nassar AH, Sakhel K, Maarouf H, et al. Adverse maternal and neonatal outcome of prolonged course of magnesium sulfate tocolysis. *Acta Obstet Gynecol Scan.* 2006; 85(9):1099-103.
4. Malaeb SN, Rassi A, Haddad MC. Bone mineralization in newborns whose mothers received magnesium sulphate for tocolysis of premature labor. *Pediatr Radiol.* 2004; 34(5):384-6. Epub 2004 Feb 18.
5. Matsuda Y, Maeda Y, Ito M, et al. Effect of magnesium sulfate treatment on neonatal bone abnormalities. *Gynecol Obstet Invest.* 1997; 44(2):82-8.
6. Schanler RJ, Smith LG, Burns PA. Effects of long-term maternal intravenous magnesium sulfate therapy on neonatal calcium metabolism and bone mineral content. *Gynecol Obstet Invest.* 1997; 43(4):236-41.
7. Santi MD, Henry GW, Douglas GL. Magnesium sulfate treatment of preterm labor as a cause of abnormal neonatal bone mineralization. *J Pediatr Orthop.* 1994; 14(2):249-53.
8. Holcomb WL, Shackelford GD, Petrie RH. Magnesium tocolysis and neonatal bone abnormalities: a controlled study. *Obstet Gynecol.* 1991; 78(4):611-4.
9. Cumming WA, Thomas VJ. Hypermagnesemia: a cause of abnormal metaphyses in the neonate. *Am J Roentgenol.* 1989; 152(5):1071-2.
10. Lamm CL, Norton KL, Murphy RJ. Congenital rickets associated with magnesium sulfate infusion for tocolysis. *J Pediatr.* 1988; 113(6):1078-82.
11. McGuinness GA, Weinstein MM, Cruikshank DP, et al. Effects of magnesium sulfate treatment on perinatal calcium metabolism. II. Neonatal responses. *Obstet Gynecol.* 1980; 56(5):595-600.
12. Riaz M, Porat R, Brodsky NL, et al. The effect of maternal magnesium sulfate treatment on newborns: a prospective controlled study. *J Perinatol.* 1998; 18(6 pt 1):449-54.

The brand names mentioned in this document are the trademarks of their respective owners.

Manufactured for:



[www.fresenius-kabi.com/us](http://www.fresenius-kabi.com/us)

Made in Norway

451340B

Revised: November 2018

**PACKAGE LABEL - PRINCIPAL DISPLAY PANEL - Magnesium Sulfate 100 mL Foil Overwrap**

To Open Overwrap – Tear at Notch

NDC 63323-108-00

# Magnesium Sulfate

*in 5% Dextrose Injection, USP*

**1g total in 100 mL**  
(0.081 mEq Mg<sup>++</sup>/mL)  
10 mg per mL



For Intravenous Infusion

Rx Only

**USE IMMEDIATELY ONCE REMOVED FROM OVERWRAP.**

Each 100 mL of sterile solution contains: Magnesium Sulfate Heptahydrate 1 g (equivalent to 8.1 mEq magnesium) and dextrose, hydrous 5 g in water for injection. May contain sulfuric acid and/or sodium hydroxide for pH adjustment. pH 4.5 (3.5 to 6.5) 333 mOsmol/Liter (calc.)

**Single Use Only – Discard Unused Portion.**

**Usual dosage:** see insert. Use only if solution is clear and container is undamaged.

**DO NOT ADD SUPPLEMENTARY MEDICATION. WHENEVER POSSIBLE USE CENTRAL ROUTE.**

After removing the overwrap, check for leaks by squeezing container. If leaks are found, discard, as sterility may be impaired.

Overwrap is a moisture barrier and is heat-sterilized.

Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Protect from freezing.

The container closure is not made with natural rubber latex. Non-PVC, Non-DEHP, Sterile.

Manufactured for:



Fresenius Kabi USA, LLC  
Lake Zurich, IL 60047  
Made in Norway



To Open Overwrap – Tear at Notch

47017C  
0179-13-001



# Magnesium Sulfate

in **5% Dextrose Injection, USP**

**1 g total in 100 mL**

(0.081 mEq Mg<sup>++</sup>/mL)

10 mg per mL

**1g**

**TOTAL**

For Intravenous Infusion

Rx Only

01-79-13-002 403016C



**PACKAGE LABEL - PRINCIPAL DISPLAY -Magnesium Sulfate 100 mL bag**

**NDC 63323-108-00**

**Magnesium Sulfate in 5% Dextrose Injection, USP**

**1 g total in 100 mL (0.081 mEq Mg<sup>++</sup>/mL) 10 mg per mL**

For Intravenous Infusion

Rx only



NDC 63323-108-00

# Magnesium Sulfate

*in 5% Dextrose Injection, USP*

**1 g total in 100 mL**



(0.081 mEq Mg<sup>++</sup>/mL) 10 mg per mL

For Intravenous Infusion Rx Only

**Each 100 mL of sterile solution contains:**

Magnesium Sulfate Heptahydrate 1 g (equivalent to 8.1 mEq magnesium) and dextrose, hydrous 5 g in water for injection. May contain sulfuric acid and/or sodium hydroxide for pH adjustment.

pH 4.5 (3.5 to 6.5) 333 mOsmol/Liter (calc.)

**Single Use Only – Discard Unused Portion.**

**Usual dosage:** see insert. Use only if solution is clear and container is undamaged.

The container closure is not made with natural rubber latex. Non-PVC, Non-DEHP, Sterile.



Manufactured for: **Fresenius Kabi USA, LLC**  
Lake Zurich, IL 60047  
Made in Norway

**EXP**  
402855C  
01-69-13-001

1234567890



**PACKAGE LABEL - PRINCIPAL DISPLAY Magnesium Sulfate 100 mL Bag Case Label**

**Product No. 610801**

**Magnesium Sulfate in 5% Dextrose Injection, USP**

**1 g total in 100 mL (10 mg per mL)**

NDC 63323-108-01 Rx only

Product No. 610801

# Magnesium Sulfate

in 5% Dextrose Injection, USP  
1 g total in 100 mL (10 mg per mL)

100 mL x 24

Store at 20° to 25°C  
(68° to 77°F) [see USP  
Controlled Room  
Temperature].  
Protect from freezing.

Manufactured for:

**FRESENIUS  
KABI**  
Fresenius Kabi USA, LLC  
Lake Zurich, IL 60047  
www.fresenius-kabi.com/us  
Made in Norway



NDC 63323-108-01

Rx only

EXP: MM-DD-YYYY

LOT: 0000000

QTY: 24



( 1 7 ) Y Y M M D D ( 1 0 ) 0 0 0 0 0 0 0 ( 3 0 ) 2 4



( 0 1 ) 3 0 3 6 3 3 2 3 1 0 8 0 1 5 ( 2 1 ) n 2 a n 2 0

63697  
FUM 0220 01-89-13-001

## MAGNESIUM SULFATE IN 5% DEXTROSE

magnesium sulfate heptahydrate injection, solution

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:63323-108
<b>Route of Administration</b>	INTRAVENOUS		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>MAGNESIUM SULFATE HEPTAHYDRATE</b> (UNII: SK47B8698T) (MAGNESIUM CATION - UNII:T6V3LHY838)	MAGNESIUM SULFATE HEPTAHYDRATE	1 g in 100 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>DEXTROSE MONOHYDRATE</b> (UNII: LX22YL083G)	
<b>SULFURIC ACID</b> (UNII: O40UQP6WCF)	
<b>SODIUM HYDROXIDE</b> (UNII: 55X04QC32I)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63323-108-01	24 in 1 CASE	03/07/2016	
1	NDC:63323-108-00	100 mL in 1 BAG; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date

ANDA	ANDA206486	03/07/2016	
------	------------	------------	--

**Labeler - Fresenius Kabi USA, LLC (608775388)**

**Establishment**

Name	Address	ID/FEI	Business Operations
HP Halden Pharma AS		347747373	MANUFACTURE(63323-108) , ANALYSIS(63323-108) , PACK(63323-108)

Revised: 4/2024

Fresenius Kabi USA, LLC