

PEPTO BISMOL CHEWS- bismuth subsalicylate tablet, chewable
The Procter & Gamble Manufacturing Company

Pepto Bismol® Chews

Drug Facts

Active ingredient (in each chewable tablet)

Bismuth subsalicylate 262 mg

Purpose

Upset stomach reliever and antidiarrheal

Uses

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
 - heartburn
 - indigestion
 - nausea
 - gas
 - belching
 - fullness

Warnings

Reye's syndrome:

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert:

Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use if you have

- an ulcer
- a bleeding problem
- bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool

Ask a doctor or pharmacist before use if you are

taking any drug for

- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

When using this product

a temporary, but harmless, darkening of the stool and/or tongue may occur

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- fully chew before swallowing
- adults and children 12 years and over:
 - 2 chewable tablets every ½ hour or 4 chewable tablets every hour as needed for diarrhea
 - 2 chewable tablets every ½ hour to 1 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- do not exceed 8 doses (16 chewable tablets) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information

- **each chewable tablet contains:**

- calcium 50 mg • salicylate 111 mg
- low sodium
- Store at no greater than 25°C. Avoid excessive heat (over 40°C.)

Inactive ingredients

acacia, beeswax, calcium carbonate, caprylic/capric triglyceride, carmine, carnauba wax, citric acid, corn starch, corn syrup, dextrin, FD&C Blue No. 2 Aluminum Lake, flavor, hydrogenated coconut oil, maltodextrin, potassium hydroxide, propylene glycol, shellac, sodium benzoate, sorbic acid, sorbitol, soy lecithin, sucrose, titanium dioxide, water

Questions?

1-800-717-3786

Tamper Evident: Do not use if inner printed seal on bottle is broken or missing.

**DIST. BY PROCTER & GAMBLE,
CINCINNATI OH 45202**

PRINCIPAL DISPLAY PANEL - 24 Chewable Tablets

Bismuth Subsalicylate

Upset Stomach Reliever/ Antidiarrheal

Pepto Bismol®

CHEWS

BERRY MINT

5 SYMPTOM

RELIEF

NAUSEA

HEARTBURN

INDIGESTION

UPSET STOMACH

DIARRHEA

24 CHEWABLE TABLETS





24 CHEWABLE TABLETS

**INDICATION
UPSET STOMACH
DIARRHEA**



3 |||01490||00202|| 5

DIST. BY: PROCTER & GAMBLE,
CINCINNATI, OH 45202

Drug Facts

Active ingredient (in each chewable tablet)	Purpose
Bismuth subsalicylate 262 mg ...	Upset stomach reliever and antidiarrheal

Uses relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
- heartburn
- indigestion
- nausea
- gas
- belching
- fullness

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)

- taking other salicylate products

Do not use if you have

- an ulcer
- a bleeding problem
- bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool

Ask a doctor or pharmacist before use if you are taking any drug for

- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

Drug Facts (continued)

If pregnant or breast feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- fully chew before swallowing
- adults and children 12 years and over:
 - 2 chewable tablets every ½ hour or 4 chewable tablets every hour as needed for diarrhea
 - 2 chewable tablets every ½ hour to 1 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- do not exceed 8 doses (16 chewable tablets) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration

caused by diarrhea

Other information

- each chewable tablet contains:
- calcium 50 mg
- salicylate 111 mg
- low sodium
- Store at no greater than 25°C. Avoid excessive heat (over 40°C.)

Inactive ingredients acacia, beeswax, calcium carbonate, caprylic/capric triglyceride, carmine, carnauba wax, citric acid, corn starch, corn syrup, dextrin, FD&C Blue No. 2 Aluminum Lake, flavor, hydrogenated coconut oil, maltodextrin, potassium hydroxide, propylene glycol, shellac, sodium benzoate, sorbic acid, sorbitol, soy lecithin, sucrose, titanium dioxide, water

Questions? 1-800-717-3786

P&G
www.pg.com

pepto-bismol.com
Patents: www.pg.com/patents

PEPTO BISMOL CHEWS

bismuth subsalicylate tablet, chewable

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69423-963
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	262 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 2 ALUMINUM LAKE (UNII: 4AQJ3LG584)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SORBIC ACID (UNII: X045WJ989B)	
SORBITOL (UNII: 506T60A25R)	
ICODEXTRIN (UNII: 2NX48Z0A9G)	
ACACIA (UNII: 5C5403N26O)	
YELLOW WAX (UNII: 2ZA36H0S2V)	
CALCIUM CARBONATE (UNII: H0G9379FGK)	
HYDROGENATED COCONUT OIL (UNII: JY81OXM10M)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
STARCH, CORN (UNII: O8232NY3SJ)	
CORN SYRUP (UNII: 9G5L16BK6N)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	
SHELLAC (UNII: 46N107B71O)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
SUCROSE (UNII: C151H8M554)	
CARMINIC ACID (UNII: CID8Z8N95N)	
WATER (UNII: 059QF0KO0R)	

Product Characteristics

Color	pink	Score	no score
Shape	ROUND	Size	18mm
Flavor	WMNTERGREEN	Imprint Code	PB
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69423-963-24	24 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	09/01/2020	
2	NDC:69423-963-72	72 in 1 CELLO PACK	09/01/2020	
2		24 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	09/01/2020	

Labeler - The Procter & Gamble Manufacturing Company (004238200)

Revised: 10/2023

The Procter & Gamble Manufacturing Company