

**SENSICAINE™ EXTRA TOPICAL ANESTHETIC- benzocaine gel**  
**Safco Dental Supply Co.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**SensiCaine™ Extra Topical Anesthetic Gel Bubble Gum Flavor**

**Active ingredients (in each gram)**

Benzocaine 200 mg

**Purpose**

Oral Anesthetic

**Use**

For oral mucosal use only, as directed by dentist. For the temporary relief of pain due to minor dental procedures.

**Warnings**

**Methemoglobinemia warning:** Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray, or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

**Allergy alert:** Do not use on patients with a history of allergies to local anesthetics such as benzocaine or other "caine" anesthetics.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly. If sore mouth symptoms do not improve in 7 days, or if irritation, pain, or redness persist or worsen, see your dentist or doctor promptly.

**Do not use**

- for teething
- in children under 2 years of age

**When using this product** Avoid contact with eyes. If contact occurs, flush with water.

**If pregnant or breast feeding,** consult a doctor before use.

**KEEP OUT OF REACH OF CHILDREN.** In case of overdose, get medical help or contact a Poison Control Center immediately.

**Directions**

- adults and children 2 years of age and older: apply only amount needed to the oral mucosa to prevent or relieve pain
- children under 2 years of age: do not use

**Other information**

- Store at 59 - 86°F (15 - 30°C).
- Protect from freezing.

**Inactive ingredients**

flavoring, PEG 3350, PEG 400, sucralose. May contain D&C Red #33, FD&C Red #40, D&C Green #5, FD&C Yellow #5 as a color additive.

**Questions or comments?**

800-621-2178

www.safcodental.com

**PRINCIPAL DISPLAY PANEL - Jar Label**

SAFCO

**SensiCaine™ Extra**

Topical Anesthetic Gel

Contains 20% Benzocaine

**Bubble Gum Flavor**

*For Professional Use Only*

1 oz. (34 g)

**Reorder #102-0456**

NDC 67239-0211-1

Distributed by:  
**Safco Dental Supply Co.**  
Buffalo Grove, IL 60089 ☎ 800-621-2178  
www.safcodental.com Made in USA

**safco**

**SensiCaine™ Extra**  
Topical Anesthetic Gel

Contains 20% Benzocaine

**Bubble Gum Flavor**

1 oz. (29.6mL) For professional use only

INGREDIENTS: Each gram of 20% benzocaine gel contains between 180-220 mg benzocaine U.S.P., in a water soluble base of PEG 3350 U.S.P., PEG 400 U.S.P., flavor, sucralose, and D&C Red #33.

1800014

Lift Here ▲

## Drug Facts

### Active ingredients (in each gram)

Benzocaine 200 mg

### Purpose

Oral Anesthetic

### Use

For oral mucosal use only, as directed by dentist. For the temporary relief of pain due to minor dental procedures.

### Warnings

#### Methemoglobinemia

**warning:** Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray, or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

**Allergy alert:** Do not use on patients with a history of allergies to local anesthetics

fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly. If sore mouth symptoms do not improve in 7 days, or if irritation, pain, or redness persist or worsen, see your dentist or doctor promptly.

#### Do not use

- for teething
- in children under 2 years of age

#### When using this product

Avoid contact with eyes. If contact occurs, flush with water.

**If pregnant or breast feeding,** consult a doctor before use.

#### KEEP OUT OF REACH OF

**CHILDREN.** In case of overdose, get medical help or contact a Poison Control Center immediately.

#### Directions

- adults and children 2 years of age and older: apply only amount needed to the oral mucosa to prevent or relieve pain
- children under 2 years of age: do not use

#### Other information

- Store at 59 - 86°F (15 - 30°C).
- Protect from freezing.

#### Inactive ingredients

flavoring, PEG 3350, PEG 400, sucralose. May contain D&C Red #33, FD&C Red #40, D&C Green #5, FD&C Yellow #5 as a color additive.

#### Questions or

such as benzocaine or other  
"caine" anesthetics.

**Sore throat warning:** If sore  
throat is severe, persists for  
more than 2 days, is  
accompanied or followed by

**comments?**

800-621-2178 •

www.safcodental.com

## SENSICAINE TM EXTRA TOPICAL ANESTHETIC

benzocaine gel

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:67239-0211
<b>Route of Administration</b>	DENTAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>Benzocaine</b> (UNII: U3RSY48JW5) (Benzocaine - UNII:U3RSY48JW5)	Benzocaine	200 mg in 1 g

### Inactive Ingredients

Ingredient Name	Strength
<b>Polyethylene Glycol 3350</b> (UNII: G2M7P15E5P)	
<b>Polyethylene Glycol 400</b> (UNII: B697894SGQ)	
<b>Sucralose</b> (UNII: 96K6UQ3ZD4)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>D&amp;C RED NO. 33</b> (UNII: 9DBA0SBB0L)	
<b>D&amp;C GREEN NO. 5</b> (UNII: 8J6RDU8L9X)	
<b>FD&amp;C YELLOW NO. 5</b> (UNII: I753WB2F1M)	

### Product Characteristics

<b>Color</b>	pink	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	BUBBLE GUM	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:67239-0211-1	29 g in 1 JAR; Type 0: Not a Combination Product	04/30/2017	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH NOT FINAL	part356	04/30/2017	

**Labeler** - Safco Dental Supply Co. (020417380)

**Registrant** - Safco Dental Supply Co. (020417380)

Revised: 11/2018

Safco Dental Supply Co.