

**EQUALINE STOMACH RELIEF - bismuth subsalicylate liquid**  
**Supervalu Inc.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Equaline Original Strength Stomach Relief Liquid**

**ACTIVE INGREDIENT(in each 30 mL)**

Bismuth subsalicylate 525 mg

**PURPOSE**

Upset stomach reliever and anti-diarrheal

**USE(S)**

relieves:

- diarrhea
- heartburn
- indigestion
- nausea
- upset stomach associated with these symptoms

**WARNINGS**

**Reye's Syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's Syndrome, a rare but serious illness.

**Allergy alert: Contains salicylate. Do not take if you are**

- allergic to salicylates (including aspirin)
- taking other salicylate products

**DO NOT USE IF YOU HAVE**

- an ulcer
- bloody or black stool

- a bleeding problem

### **ASK A DOCTOR BEFORE USE IF YOU HAVE**

- fever
- mucus in the stool

### **ASK A DOCTOR OR PHARMACIST BEFORE USE IF YOU ARE**

taking any drug for

- anticoagulation (thinning of the blood)
- diabetes
- gout
- arthritis

### **WHEN USING THIS PRODUCT**

a temporary, but harmless darkening of the stool and/or tongue may occur

### **STOP USE AND ASK DOCTOR IF**

- symptoms get worse
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

### **IF PREGNANT OR BREAST-FEEDING,**

ask a health professional before use.

### **KEEP OUT OF REACH OF CHILDREN**

In case of overdose, get medical help or contact a Poison Control Center immediately.

### **DIRECTIONS**

- **shake well before use**
- mL = milliliter
- TBSP = tablespoon
- adults and children 12 years and over: 1 dose (2 TBSP or 30 mL) every 1/2 to 1 hour as needed
- do not exceed 8 doses (16 TBSP or 240 mL) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of fluids to help prevent dehydration caused by diarrhea

### **OTHER INFORMATION**

- **each 30 mL or 2 TBSP contains:**
- potassium 25 mg
- salicylate 260 mg
- sodium 8 mg
- protect from freezing
- avoid excessive heat (over 104°F or 40°C)
- dosage cup provided

## **INACTIVE INGREDIENTS**

benzoic acid, D&C red # 22, D&C red # 28, flavor, hydroxyethyl cellulose, potassium hydroxide, purified water, saccharin sodium, salicylic acid, simethicone, xanthan gum

## **PRINCIPAL DISPLAY PANEL**

NDC 41163-936-06

compare to **Pepto-Bismol**<sup>®</sup> active ingredient\*

EQUALINE

original strength

**stomach relief liquid**

bismuth subsalicylate 525 mg per 30 mL

(upset stomach reliever/antidiarrheal)

5 symptom digestive relief:

- heartburn
- indigestion
- nausea
- upset stomach
- diarrhea

**16 FL OZ (1 pt) 473 mL**

DO NOT USE IF IMPRINTED SHRINKBAND IS MISSING OR BROKEN.

NDC 41163-936-06

compare to  
Pepto-Bismol®  
active ingredient\*

**Drug Facts**

**Active ingredient (in each 30 mL)** Purpose  
Bismuth subsalicylate 525 mg ..... Upset stomach reliever  
and anti-diarrheal

**Uses** relieves: ■ diarrhea ■ heartburn ■ indigestion ■ nausea  
■ upset stomach associated with these symptoms

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Contains salicylate. Do not take if you are  
■ allergic to salicylates (including aspirin)  
■ taking other salicylate products

**Do not use if you have** ■ an ulcer ■ bloody or black stool  
■ a bleeding problem

**Ask a doctor before use if you have**  
■ fever ■ mucus in the stool

**Ask a doctor or pharmacist before use if you are taking any drug for** ■ anticoagulation (thinning of the blood) ■ diabetes  
■ gout ■ arthritis

**When using this product** a temporary, but harmless darkening of the stool and/or tongue may occur

**Stop use and ask a doctor if** ■ symptoms get worse  
■ ringing in the ears or loss of hearing occurs  
■ diarrhea lasts more than 2 days

**If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center immediately.

◀ Drug Facts continued

**EQUALINE®**

original strength  
**stomach  
relief liquid**

bismuth subsalicylate  
525mg per 30mL  
(upset stomach reliever/  
anti-diarrheal)

5 symptom digestive relief

- heartburn
- indigestion
- nausea
- upset stomach
- diarrhea

16 FL OZ (1 pt) 473mL

**Drug Facts (continued)**

**Directions** ■ shake well before use  
■ mL = milliliter ■ Tbsp = tablespoon ■ adults and children 12 years and over: 1 dose (2 Tbsp or 30 mL) every 1/2 to 1 hour as needed ■ do not exceed 8 doses (16 Tbsp or 240 mL) in 24 hours ■ use until diarrhea stops but not more than 2 days  
■ children under 12 years: ask a doctor ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

**Other information** ■ each 30 mL or 2 Tbsp contains:  
• potassium 25 mg • salicylate 260 mg • sodium 8 mg  
■ protect from freezing. ■ avoid excessive heat (over 104°F or 40°C). ■ dosage cup provided

**Inactive ingredients** benzoic acid, D&C red # 22, D&C red # 28, flavor, hydroxyethyl cellulose, potassium hydroxide, purified water, saccharin sodium, salicylic acid, simethicone, xanthan gum.

DISTRIBUTED BY SUPERVALU INC., EDEN PRAIRIE, MN 55344 USA  
877-932-7948, [supervaluprivatebrands.com](http://supervaluprivatebrands.com)

\*This product is not manufactured or distributed by The Procter & Gamble Company, owner of the registered trademark Pepto-Bismol®.



**EQUALINE STOMACH RELIEF**

bismuth subsalicylate liquid

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:41163-936
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>BISMUTH SUBSALICYLATE</b> (UNII: 62TEY51RR1) (SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	525 mg in 30 mL

**Inactive Ingredients**

Ingredient Name	Strength
<b>BENZOIC ACID</b> (UNII: 8SKN0B0MIM)	
<b>D&amp;C RED NO. 22</b> (UNII: 1678RKX8RT)	
<b>D&amp;C RED NO. 28</b> (UNII: 767IP0Y5NH)	
<b>POTASSIUM HYDROXIDE</b> (UNII: WZH3C48M4T)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SALICYLIC ACID</b> (UNII: O414PZ4LPZ)	
<b>HYDROXYETHYL CELLULOSE (1500 MPAS AT 1%)</b> (UNII: L605B5892V)	
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O)	

XANTHAN GUM (UNII: TTV12P4NEE)

### Product Characteristics

<b>Color</b>	PINK	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41163-936-05	354 mL in 1 BOTTLE; Type 0: Not a Combination Product	09/23/2018	
2	NDC:41163-936-06	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	09/23/2018	
3	NDC:41163-936-04	237 mL in 1 BOTTLE; Type 0: Not a Combination Product	09/23/2018	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part335	09/23/2018	

**Labeler** - Supervalu Inc. (006961411)

### Establishment

Name	Address	ID/FEI	Business Operations
Guardian Drug Company		119210276	MANUFACTURE(41163-936)

Revised: 11/2022

Supervalu Inc.