

SODIUM CHLORIDE- sodium chloride injection, solution

Medefil, Inc.

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use SODIUM CHLORIDE INJECTION, USP, 0.9% safely and effectively. See full prescribing information for SODIUM CHLORIDE INJECTION, USP, 0.9%.

SODIUM CHLORIDE INJECTION, USP, 0.9%

Initial U.S. Approval: 2012

INDICATIONS AND USAGE

Sodium Chloride Injection, USP, 0.9% is indicated for:

- Dilution or Dissolving the drugs for intravenous, intramuscular or subcutaneous injections (1.1).

DOSAGE AND ADMINISTRATION

- Volume of preparation to be used for diluting or dissolving any drug for injection is dependent on the vehicle concentration, dose and route of administration recommended by the drug manufacturer (2.1).

DOSAGE FORMS AND STRENGTHS

- Injection, 0.9% (3)

CONTRAINDICATIONS

- None (4)

WARNINGS AND PRECAUTIONS

- Consult the drug product manufacturer's instructions for choice of vehicle, appropriate dilution or volume for dissolving drugs to be injected including the route and rate of injection (5.1).
- Do not use Sodium Chloride Injection, USP, 0.9% if the solution is discolored, cloudy, hazy, or contains a precipitate, or if the syringe is damaged (5.1).
- For single use only. Discard unused portion (5.2).

ADVERSE REACTIONS

- Reactions that may occur because of this solution, added drugs or the technique of reconstitution or administration include air embolization, febrile response, local tenderness, abscess, tissue necrosis or infection at the site of injection, venous thrombosis or phlebitis extending from the site of injection and extravasations (6).

To report SUSPECTED ADVERSE REACTIONS, contact MEDEFIL, INC. at tel:1-630-682-4600 and <http://www.medefilinc.com/> or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

- Some drugs or injections may be incompatible when combined with 0.9% sodium chloride.
- Consult with a pharmacist, if unsure of compatibility (7).

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 6/2022

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS & USAGE

1.1 Dilution or Dissolution of Drugs

2 DOSAGE & ADMINISTRATION

2.1 Recommended Dosage

2.2 Instruction for Administration

2.3 Preparation and Handling Precautions

3 DOSAGE FORMS & STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

5.1 General

5.2 For Single Use Only

6 ADVERSE REACTIONS

7 DRUG INTERACTIONS

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS & USAGE

1.1 Dilution or Dissolution of Drugs

Sodium Chloride Injection, USP, 0.9% is indicated for diluting or dissolving drugs for intravenous, intramuscular or subcutaneous injection, according to instructions of the manufacturer of the drug to be administered.

2 DOSAGE & ADMINISTRATION

2.1 Recommended Dosage

- The volume of preparation to be used for diluting or dissolving any drug for injection is dependent on the vehicle concentration, dose and route of administration as recommended by the drug manufacturer.

2.2 Instruction for Administration

- Solution and fluid path are sterile and non-pyrogenic if the tip cap is in place, syringe is intact and there is no evidence of leakage. Use proper aseptic technique.
- Inspect plastic wrapping. Do not use if plastic wrapping is damaged or not intact. (Figure 1)
- Remove plastic packaging by tearing along perforation. (Figure 2)
- Do not use if, solution is discolored, cloudy, hazy, or contains a precipitate, or if the syringe is damaged. (Figure 3)
- With the tip cap of the syringe on, press the syringe forward to properly activate the syringe. Improper activation may difficult syringe use and/or may cause plunger rod separation. **Never draw back rod because the product may become contaminated.** (Figure 4)
- Remove tip cap. Hold the syringe unit upright and prime to expel any air bubble if present. (Figure 5)
- Syringe is now ready to use.

- Use aseptic techniques for diluting or dissolving drugs, mix thoroughly and use according to the drug manufacturer's label instructions.
- Discard empty unit after use. Discard any unused portion. Do not reuse disposable syringes.

2.3 Preparation and Handling Precautions

- Before use check for any possible incompatibility with sodium chloride.
- Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration whenever solution and container permit. Do not administer unless solution is clear and seal is intact.
- Do not store reconstituted solution or drugs for injection unless otherwise directed by the manufacturer of the solute.
- Adding additives may be incompatible.
- Do not use if, solution is discolored, cloudy, hazy, or contains a precipitate, or if the syringe is damaged.
- Do not freeze.
- Exposure of pharmaceutical products to heat should be minimized. Avoid excessive heat.

3 DOSAGE FORMS & STRENGTHS

10 mL fill in 12 mL syringe, 0.9% Sodium Chloride, USP, single use.

4 CONTRAINDICATIONS

None

5 WARNINGS AND PRECAUTIONS

5.1 General

When used to dilute drug products, consult the drug product manufacturer's instructions for choice of vehicle, appropriate dilution or volume for dissolving drugs to be injected including the route and rate of injection. Inspect reconstituted (diluted or dissolved) drugs for clarity (if soluble) and freedom from unexpected precipitation or discoloration prior to administration. Do not use Sodium Chloride Injection, USP 0.9% if the solution is discolored, cloudy, hazy, or contains a precipitate, or if the syringe is damaged.

5.2 For Single Use Only

Re-use of single-use product creates a potential risk to the user. Contamination of product and/or limited functionality of the syringe may lead to injury, illness or death. Discard any unused portion.

6 ADVERSE REACTIONS

Adverse reactions which may occur because of this solution, added drugs or the

technique of reconstitution or administration include, but are not limited to, air embolization, febrile response, local tenderness, abscess, tissue necrosis or infection at the site of injection, venous thrombosis or phlebitis extending from the site of injection and extravasation.

If an adverse reaction does occur, discontinue the infusion, evaluate the patient, institute appropriate therapeutic countermeasures, and if possible, retrieve and save the remainder of unused vehicle for examination if deemed necessary.

7 DRUG INTERACTIONS

Some drugs or injections may be incompatible when combined with 0.9% sodium chloride. Before Sodium Chloride Injection, USP, 0.9% is used as a vehicle for the administration of a drug, the drug product manufacturer's instructions or other specific references should be checked for any possible incompatibility with sodium chloride.

Consult with a pharmacist, if unsure of compatibility.

11 DESCRIPTION

Sodium Chloride Injection, USP 0.9% is a sterile, nonpyrogenic, isotonic solution of sodium chloride and water for injection. It contains no bacteriostatic antimicrobial agents or added buffer. The nominal pH is 5.5 (4.5 to 7.0).

Hydrochloric acid and/or sodium hydroxide may have been added for pH adjustment. Sodium Chloride Injection, USP, 0.9% contains 9 g/L Sodium Chloride, USP (NaCl) with a calculated osmolarity of approximately 308 mOsmol/L. It contains 154 mEq/L sodium and 154 mEq/L chloride.

The syringe component of Sodium Chloride Injection, USP, 0.9% is manufactured with polypropylene and luer lock.

The syringe is not made with natural rubber latex. The syringe is not made with DEHP. This product contains no preservative. The syringes require no vapor barrier to maintain the proper drug concentration. The empirical formula for sodium chloride is NaCl and the molecular weight is 58.44 g/mol.

Supplied as single use syringes.

12 CLINICAL PHARMACOLOGY

Sodium chloride in water dissociates to provide sodium (Na⁺) and chloride (Cl⁻) ions. These ions are normal constituents of the body fluids (principally extracellular) and are essential for maintaining electrolyte balance.

The small volume of fluid and amount of sodium chloride provided by Sodium Chloride Injection, USP, 0.9% when used only as an isotonic vehicle for parenteral injection of drugs or for flushing of indwelling access devices, is unlikely to exert a significant effect on fluid and electrolyte balance except possibly in neonates and very small infants.

16 HOW SUPPLIED/STORAGE AND HANDLING

The drug product is packaged in a clear plastic hypodermic syringe, which consists of a hypodermic barrel with luer lock, plunger stopper, plunger rod, and tip cap. The syringe is intended for single use and subsequent disposal. The 10 mL fill volume syringes are packaged individually in plastic pouches and 60 syringes are packaged into each dispensing box.

DOSAGE FORM

| Product Number | NDC | Fill Volume |
|----------------|--------------|-----------------------------|
| MSD-0230 | 64253-202-30 | 10 mL fill in 12 mL syringe |

The above product is available in boxes of 30, 60 or 120 count each.

STORAGE AND HANDLING

Store at 25°C (77°F); excursions permitted to 15°-30°C (59° - 86°F). Do not freeze.

17 PATIENT COUNSELING INFORMATION


17.1 When using to dilute drug products, consult the drug product manufacturer's instruction to confirm compatibility, appropriate dilution or volume for dissolving drugs including the route and rate of injection. Inspect reconstituted (diluted or dissolved) drugs for clarity (if soluble) and freedom from unexpected precipitation or discoloration prior to administration. Do not use Sodium Chloride Injection, USP 0.9% if the solution is discolored, cloudy, hazy, or contains a precipitate, or if the syringe damaged [see *Warning and Precautions, section 5.1*].

17.2 Syringes are for single use only. Discard unused portions and dispose of the unit in an appropriate sharps container. [see *Warning and Precautions, section 5.2*].

Medefil, Inc., 250 Windy Point Drive, Glendale Heights, IL 60139. Prepared 10/2020 Rev. 003

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

NDC 64253-202-30

 = 60 Syringes

SODIUM CHLORIDE INJECTION, USP, 0.9%
10 mL fill in 12 mL Syringe, Rx Only

For Use as a STERILE DILUENT

This product is not made with natural rubber latex.

No Preservatives

Each mL contains 9 mg Sodium Chloride, USP
(0.308 mOsmol/mL) in Water for Injection

For Single Use Only. Sterile and Non-Pyrogenic

Recommended Storage: Store at 25°C (77°F); excursions permitted to 15° - 30°C (59° - 86°F). Keep from freezing.

Usual Dose: See package insert.

Do not use if solution is discolored, cloudy, hazy, or contains a precipitate.



Medefil, Inc., Glendale Heights, IL 60139



(01) 3 03 64253 20230 9

MSD-10/12 Rev. 003

REF MSD-0230


1mL 2mL 3mL 4mL 5mL 6mL 7mL 8mL 9mL 10mL 11mL 12mL
Approx. mL

NDC 64253-202-30


This product is not made with natural rubber latex.
No Preservatives

SODIUM CHLORIDE INJECTION, USP, 0.9%
10 mL fill in 12 mL Syringe, Rx Only


Each mL contains 9 mg Sodium Chloride, USP in Water for Injection
Single Use. Discard Unused Portion. Sterile, Nonpyrogenic.
For Use as a **STERILE DILUENT**. Usual Dose: See package insert.



(01) 1 03 64253 20230 5

 Medefil, Inc., Glendale Heights, IL 60139

Rev. 003

LOT / **EXP.** 

REF MSD-0230

Manufactured by:

Medefil, Inc., 250 Windy Point Drive, Glendale Heights, IL 60139.

NDA 202832

Rev. 003

Package Insert_Page 1 and 2

SODIUM CHLORIDE

sodium chloride injection, solution

Product Information

| | | | |
|--------------------------------|--|---------------------------|---------------|
| Product Type | HUMAN PRESCRIPTION DRUG | Item Code (Source) | NDC:64253-202 |
| Route of Administration | INTRAMUSCULAR, SUBCUTANEOUS, INTRAVENOUS | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|-------------------|--------------|
| SODIUM CHLORIDE (UNII: 451W47IQ8X) (SODIUM CATION - UNII:LYR4M0NH37) | SODIUM CHLORIDE | 9 mg in 1 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|---------------------------------|----------|
| WATER (UNII: 059QF0KO0R) | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:64253-202-30 | 60 in 1 BOX | 01/13/2012 | |
| 1 | | 10 mL in 1 SYRINGE, PLASTIC; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.) | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| NDA | NDA202832 | 01/13/2012 | |

