

# VICKS NYQUIL SEVERE COLD AND FLU- acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, and phenylephrine hydrochloride solution

The Procter & Gamble Manufacturing Company

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

Vicks® NyQuil®

## Severe Cold & Flu

### Drug Facts

<b>Active ingredients (in each 30 mL dose cup)</b>	<b>Purpose</b>
Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Doxylamine succinate 12.5 mg	Antihistamine
Phenylephrine HCl 10 mg	Nasal decongestant

### Uses

temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion & pressure
- cough due to minor throat & bronchial irritation
- cough to help you sleep
- minor aches & pains
- headache
- fever
- sore throat
- runny nose & sneezing
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage

### Warnings

#### Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

#### Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- to make a child sleep

**Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland
- a sodium-restricted diet

**Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

**When using this product**

- **do not use more than directed**
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

**Stop use and ask a doctor if**

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning**

Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- take only as directed - see Overdose warning
- use dose cup or tablespoon (TBSP)
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL (2 TBSP) every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

- **when using other DayQuil<sup>®</sup> or NyQuil products, carefully read each label to ensure correct dosing**

#### Other information

- **each 30 mL dose cup contains:** sodium 92 mg
- store at room temperature

#### Inactive ingredients

citric acid, FD&C Blue No.1, FD&C Red No. 40, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose, xanthan gum

#### Questions?

**1-800-362-1683**

**Dist. by**

**Procter & Gamble,  
Cincinnati OH 45202**

#### PRINCIPAL DISPLAY PANEL - 354 mL Bottle Label

**NEW**

**Fights More**

**Symptoms**

than NyQuil

Cold & Flu

**VICKS<sup>®</sup>**

**NyQuil<sup>®</sup>**

**SEVERE**

**COLD & FLU**

**Nighttime Relief**

*Acetaminophen, Phenylephrine HCl, Doxylamine Succinate, Dextromethorphan HBr*

**MAX**

**STRENGTH**

- **Headache, Fever, Sore Throat,  
Minor Aches & Pains**
- **Nasal/Sinus Congestion & Sinus Pressure**
- **Sneezing, Runny Nose**
- **Cough**

***Berry Flavor***

**12 FL OZ (354 ml)**

**NEW**  
Fights More  
Symptoms  
than NyQuil  
Cold & Flu



# NyQuil<sup>®</sup>

## SEVERE

**COLD & FLU**

**Nighttime Relief**

*Acetaminophen, Phenylephrine HCl, Doxylamine Succinate, Dextromethorphan HBr*

**MAX  
STRENGTH**

- Headache, Fever, Sore Throat,  
Minor Aches & Pains
- Nasal/Sinus Congestion & Sinus Pressure
- Sneezing, Runny Nose
- Cough

*Berry Flavor*

**12 FL OZ (354 ml)**

**TAMPER EVIDENT:** Do not use if printed shrinkband is missing or broken.  
Failure to follow these warnings could result in serious consequences.

**PARENTS:**  
Learn about teen medicine abuse



## Drug Facts

<b>Active Ingredients (in each 30 mL dose cup)</b>	<b>Purpose</b>
Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Doxylamine succinate 12.5 mg	Antihistamine
Phenylephrine HCl 10 mg	Nasal decongestant

**Uses** temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion & pressure
- cough due to minor throat & bronchial irritation
- cough to help you sleep
- minor aches & pains
- headache
- fever
- sore throat
- runny nose & sneezing
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hours, which is the maximum daily amount for this product

- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- to make a child sleep

**Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland
- a sodium-restricted diet

### Drug Facts (continued)

**Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

**When using this product**

- do not use more than directed
- excitability may occur, especially



Dist. by 92400410  
**Procter & Gamble,**  
 Cincinnati OH 45202



in children • marked drowsiness may occur • avoid alcoholic drinks  
 • be careful when driving a motor vehicle or operating machinery  
 • alcohol, sedatives, and tranquilizers may increase drowsiness

**Stop use and ask a doctor if** • you get nervous, dizzy or sleepless  
 • pain, nasal congestion, or cough gets worse or lasts more than 7 days  
 • fever gets worse or lasts more than 3 days • redness or swelling is present  
 • new symptoms occur • cough comes back or occurs with rash or headache  
 that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use.**  
**Keep out of reach of children. Overdose warning:** Taking more than directed  
 can cause serious health problems. In case of overdose, get medical help or  
 contact a Poison Control Center right away. Quick medical attention is critical for  
 adults as well as for children even if you do not notice any signs or symptoms.

**Directions** • take only as directed - see Overdose warning  
 • use dose cup or tablespoon (Tbsp) • do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL (2 Tbsp) every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

• when using other DayQuil® or NyQuil products,  
 carefully read each label to ensure correct dosing

**Other information** • each 30 mL dose cup  
 contains: sodium 92 mg  
 • store at room temperature

**Inactive ingredients** citric acid, FD&C Blue  
 No.1, FD&C Red No. 40, flavor, glycerin, propylene  
 glycol, purified water, saccharin sodium, sodium  
 benzoate, sodium chloride, sodium citrate, sorbitol,  
 sucralose, xanthan gum

**Questions? 1-800-362-1683** [www.vicks.com](http://www.vicks.com)

**QUIET ZONE 8**  
 P&U  
 TRUNCATED  
 80% SIZE  
 12-digit UPC  
 (non-suppressed)  
 For Position Only  
 23900 03815  
**QUIET ZONE 3**

## VICKS NYQUIL SEVERE COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, and phenylephrine hydrochloride solution

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:37000-812
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH)	DEXTROMETHORPHAN	20 mg

(DEXTROMETHORPHAN - UNII:7355X3ROTS)	HYDROBROMIDE	in 30 mL
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 30 mL

### Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

### Product Characteristics

Color	red	Score	
Shape		Size	
Flavor	BERRY	Imprint Code	
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37000-812-08	236 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/22/2013	
2	NDC:37000-812-12	354 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/22/2013	
3	NDC:37000-812-24	2 in 1 PACKAGE	07/22/2013	
3		354 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
4	NDC:37000-812-36	3 in 1 PACKAGE	07/22/2013	07/10/2019
4		354 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	07/22/2013	



---

**Labeler** - The Procter & Gamble Manufacturing Company (004238200)

Revised: 2/2021

The Procter & Gamble Manufacturing Company