

**OXYTOCIN- oxytocin injection**  
**West-Ward Pharmaceuticals Corp**

-----  
**OXYTOCIN INJECTION, USP**


**(Synthetic)**

**FOR INTRAVENOUS INFUSION OR INTRAMUSCULAR USE**

**Rx Only**

**DESCRIPTION**

Each mL of Oxytocin Injection, USP (synthetic), intended for intravenous infusion or intramuscular injection, possesses an oxytocic activity equivalent to 10 USP Oxytocin Units and contains chlorobutanol (chloral derivative) 0.5%. This product may contain up to 12.5% decomposition products/impurities. Oxytocin injection (synthetic) is a sterile, clear, colorless solution of oxytocin in Water for Injection, USP prepared by synthesis. Acetic acid may have been added for pH adjustment (pH 3.0 - 5.0). The structural formula is:



**CLINICAL PHARMACOLOGY**

Oxytocin injection (synthetic) acts on the smooth muscle of the uterus to stimulate contractions; response depends on the uterine threshold of excitability. It exerts a selective action on the smooth musculature of the uterus, particularly toward the end of pregnancy, during labor and immediately following delivery. Oxytocin stimulates rhythmic contractions of the uterus, increases the frequency of existing contractions and raises the tone of the uterine musculature. Synthetic oxytocin does not possess the cardiovascular effects, such as elevation of blood pressure, as exhibited by vasopressin found in posterior pituitary injection.

**INDICATIONS AND USAGE**

**IMPORTANT NOTICE**

**Oxytocin Injection, USP (synthetic) is indicated for the medical rather than the elective induction of labor. Available data and information are inadequate to define the benefits to risks considerations in the use of the drug product for elective induction. Elective induction of labor is defined as the initiation of labor for convenience in an individual with a term pregnancy who is free of medical indications.**

***Antepartum***

Oxytocin Injection, USP (synthetic) is indicated for the initiation or improvement of uterine contractions, where this is desirable and considered suitable, in order to achieve early vaginal delivery for fetal or maternal reasons. It is indicated for (1) induction of labor in patients with a medical indication for the initiation of labor, such as Rh problems, maternal diabetes, pre-eclampsia at or near term, when delivery is in the best interest of mother and fetus or when membranes are prematurely ruptured and delivery is

indicated; (2) stimulation or reinforcement of labor, as in selected cases of uterine inertia; (3) adjunctive therapy in the management of incomplete or inevitable abortion. In the first trimester, curettage is generally considered primary therapy. In second trimester abortion, oxytocin infusion will often be successful in emptying the uterus. Other means of therapy, however, may be required in such cases.

### ***Postpartum***

Oxytocin Injection, USP (synthetic) is indicated to produce uterine contractions during the third stage of labor and to control postpartum bleeding or hemorrhage.

## **CONTRAINDICATIONS**

Oxytocin injection (synthetic) is contraindicated in any of the following conditions:

- Significant cephalopelvic disproportion;
- Unfavorable fetal positions or presentations which are undeliverable without conversion prior to delivery, i.e., transverse lies;
- In obstetrical emergencies where the benefit-to-risk ratio for either the fetus or the mother favors surgical intervention;
- In cases of fetal distress where delivery is not imminent;
- Prolonged use in uterine inertia or severe toxemia;
- Hypertonic uterine patterns;
- Patients with hypersensitivity to the drug;
- Induction or augmentation of labor in those cases where vaginal delivery is contraindicated, such as cord presentation or prolapse, total placenta previa, and vasa previa.

## **WARNINGS**

Oxytocin injection (synthetic) when given for induction or stimulation of labor, must be administered only by the intravenous route, and with adequate medical supervision in a hospital.

## **PRECAUTIONS**

### ***General***

All patients receiving intravenous oxytocin must be under continuous observation by trained personnel with a thorough knowledge of the drug and qualified to identify complications. A physician qualified to manage any complications should be immediately available.

When properly administered, oxytocin should stimulate uterine contractions similar to those seen in normal labor. Overstimulation of the uterus by improper administration can be hazardous to both mother and fetus. Even with proper administration and adequate supervision, hypertonic contractions can occur in patients whose uteri are hypersensitive to oxytocin.

Except in unusual circumstances, oxytocin should not be administered in the following conditions: prematurity, borderline cephalopelvic disproportion, previous major surgery on the cervix or uterus including Caesarean section, overdilatation of the uterus, grand multiparity or invasive cervical carcinoma. Because of the variability of the combinations of factors which may be present in the conditions above, the definition of "unusual circumstances" must be left to the judgement of the physician. The decision can only be made by carefully weighing the potential benefits which oxytocin can provide in a given case against rare but definite potential for the drug to produce hypertonicity or tetanic spasm.

Maternal deaths due to hypertensive episodes, subarachnoid hemorrhage, rupture of the uterus and fetal deaths due to various causes have been reported associated with the use of parenteral oxytocic drugs for induction of labor and for augmentation in the first and second stages of labor.

Oxytocin has been shown to have an intrinsic antidiuretic effect, acting to increase water reabsorption from the glomerular filtrate. Consideration should, therefore, be given to the possibility of water intoxication, particularly when oxytocin is administered continuously by infusion and the patient is receiving fluids by mouth.

### ***Drug Interactions***

Severe hypertension has been reported when oxytocin was given three to four hours following prophylactic administration of a vasoconstrictor in conjunction with caudal block anesthesia. Cyclopropane anesthesia may modify oxytocin's cardiovascular effects, so as to produce unexpected results such as hypotension. Maternal sinus bradycardia with abnormal atrioventricular rhythms has also been noted when oxytocin was used concomitantly with cyclopropane anesthesia.

### ***Carcinogenesis, Mutagenesis, Impairment of Fertility***

There are no animal or human studies on the carcinogenicity and mutagenicity of this drug, nor is there any information on its effect on fertility.

### ***Pregnancy Category C***

There are no known indications for use of oxytocin in the first and second trimester of pregnancy other than in relation to spontaneous or induced abortion. Based on the wide experience with this drug and its chemical structure and pharmacological properties, it would not be expected to present a risk of fetal abnormalities when used as indicated.

### ***Nonteratogenic Effects-***

See **ADVERSE REACTIONS** in the fetus or infant.

### ***Labor and Delivery-***

See **INDICATIONS AND USAGE**.

### ***Nursing Mothers***

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when oxytocin is administered to a nursing woman.

## **ADVERSE REACTIONS**

The following adverse reactions have been reported in the mother:

---

Anaphylactic reaction	Nausea
Postpartum hemorrhage	Vomiting
Cardiac arrhythmia	Premature ventricular contractions
Fatal afibrinogenemia	Pelvic hematoma

---

Excessive dosage or hypersensitivity to the drug may result in uterine hypertonicity, spasm, tetanic contraction or rupture of the uterus.

The possibility of increased blood loss and afibrinogenemia should be kept in mind when administering the drug.

Severe water intoxication with convulsions and coma has occurred, and is associated with a slow oxytocin infusion over a 24-hour period. Maternal death due to oxytocin-induced water intoxication has been reported.

The following adverse reactions have been reported in the fetus or infant:

<b><i>Due to induced uterine mobility</i></b>	<b><i>Due to use of oxytocin in the mother</i></b>
Bradycardia	Neonatal retinal hemorrhage
Premature ventricular contractions and other arrhythmias	Low Apgar scores at five minutes
Permanent CNS or brain damage	Neonatal jaundice
Fetal death	

## **OVERDOSAGE**

Overdosage with oxytocin injection (synthetic) depends essentially on uterine hyperactivity whether or not due to hypersensitivity to this agent. Hyperstimulation with strong (hypertonic) or prolonged (tetanic) contractions, or a resting tone of 15 to 20 mm H<sub>2</sub>O or more between contractions can lead to tumultuous labor, uterine rupture, cervical and vaginal lacerations, postpartum hemorrhage, uteroplacental hypoperfusion and variable deceleration of fetal heart, fetal hypoxia, hypercapnia or death. Water intoxication with convulsions, which is caused by the inherent antidiuretic effect of oxytocin, is a serious complication that may occur if large doses (40 to 50 milliunits/minute) are infused for long periods. Management consists of immediate discontinuation of oxytocin, and symptomatic and supportive therapy.

## **DOSAGE AND ADMINISTRATION**

Dosage of oxytocin is determined by uterine response. The following dosage information is based upon the various regimens and indications in general use.

### ***Induction or Stimulation of Labor***

Intravenous infusion (drip method) is the only acceptable method of administration for the induction or stimulation of labor.

Accurate control of the rate of infusion flow is essential. An infusion pump or other such device and frequent monitoring of strength of contractions and fetal heart rate are necessary for the safe administration of oxytocin for the induction or stimulation of labor. If uterine contractions become too powerful, the infusion can be abruptly stopped, and oxytocic stimulation of the uterine musculature will soon wane.

An intravenous infusion of a non-oxytocin containing solution should be started. Physiologic electrolyte solutions should be used except under unusual circumstances.

To prepare the usual solution for intravenous infusion - one mL (10 units) is combined aseptically with 1,000 mL of a non-hydrating diluent.

The combined solution, rotated in the infusion bottle to insure thorough mixing, contains 10 mU/mL. Add the container with dilute oxytocic solution to the system through the use of a constant infusion pump or other such device to control accurately the rate of infusion.

The initial dose should be no more than 1 to 2 mU/min. The dose may be gradually increased in increments of no more than 1 to 2 mU/min, until a contraction pattern has been established which is similar to normal labor.

The fetal heart rate, resting uterine tone, and the frequency, duration, and force of contractions should be monitored.

The oxytocin infusion should be discontinued immediately in the event of uterine hyperactivity or fetal distress. Oxygen should be administered to the mother. The mother and fetus must be evaluated by the responsible physician.

### ***Control of Postpartum Uterine Bleeding***

*Intravenous Infusion (Drip Method)* – To control postpartum bleeding, 10 to 40 units of oxytocin may be added to 1,000 mL of a nonhydrating diluent and run at a rate necessary to control uterine atony.

*Intramuscular Administration* – 1 mL (10 units) of oxytocin can be given after delivery of the placenta.

### ***Treatment of Incomplete or Inevitable Abortion***

Intravenous infusion with physiologic saline solution, 500 mL, or 5% dextrose in physiologic saline solution to which 10 units of oxytocin have been added should be infused at a rate of 20 to 40 drops/minute.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit.

### **HOW SUPPLIED**

Oxytocin Injection, USP (synthetic) is supplied as follows:

NDC 0143-9743-25 Package of 25. 1 mL fill in a 2 mL vial, 10 USP Units per mL.

NDC 0143-9742-25 Package of 25. 10 mL fill in a 10 mL vial, 10 USP Units per mL.

Discard unused portion.

Use only if solution is clear and seal is intact.

Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].

Do not permit to freeze.

**Manufactured by:** HIKMA FARMACÊUTICA (PORTUGAL), S.A.

Estrada do Rio da Mó, nº 8, 8A e 8B - Fervença, - 2705 – 906 Terrugem SNT - PORTUGAL

**Distributed by:**

WEST-WARD PHARMACEUTICAL CORP.

Eatontown, NJ 07724 USA

Issued: September 2013

PIN312-WES/1

### **PRINCIPAL DISPLAY PANEL**

NDC 0143-9743-01

**OXYTOCIN**

INJECTION, USP

(SYNTHETIC)

10 USP Units/mL

**STERILE**

FOR IV (INFUSION)

OR IM USE

Rx ONLY

1 mL Single Dose Vial

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].

**USUAL DOSAGE:** See package insert.

Note: Oversize Vial

NDC 0143-9743-01

**OXYTOCIN**  
INJECTION, USP  
(SYNTHETIC)

**10 USP Units/mL**

**STERILE**

FOR IV (INFUSION)  
OR IM USE  
Rx ONLY

1 mL Single Dose Vial  
Store at 20° to 25°C (68° to 77°F) [See USP  
Controlled Room Temperature].  
**USUAL DOSAGE:** See package insert.  
Note: Oversize Vial.  
Mfd by: HIKMA FARMACÊUTICA (PORTUGAL), S.A.  
Dist by: WEST-WARD PHARMACEUTICAL CORP.  
Eatontown, NJ 07724 USA

  
NDC 0143-9743-01  
PLB693-WES/1

Lot: \_\_\_\_\_  
Exp: \_\_\_\_\_

Non-varnish area

**PRINCIPAL DISPLAY PANEL**

NDC 0143-9742-01  
**OXYTOCIN**  
INJECTION, USP  
(SYNTHETIC)  
100 USP Units/10 mL  
(10 USP Units/mL)  
FOR IV (INFUSION)  
OR IM USE  
Rx ONLY  
10 mL Multiple Dose Vial  
Each mL contains Oxytocin activity equivalent  
to 10 USP Oxytocin Units; Chlorobutanol 0.5%;  
Water for Injection q.s. Acetic Acid may have  
been added for pH adjustment.  
**USUAL DOSAGE:** See package insert.  
Store at 20° to 25°C (68° to 77°F) [See USP  
Controlled Room Temperature].  
Do not permit to freeze.

NDC 0143-9742-01

**OXYTOCIN**  
INJECTION, USP  
(SYNTHETIC)

**100 USP Units/10 mL**  
(10 USP Units/mL)

FOR IV (INFUSION)  
OR IM USE  
Rx ONLY

10 mL Multiple Dose Vial  
Each mL contains Oxytocin activity equivalent  
to 10 USP Oxytocin Units; Chlorobutanol 0.5%;  
Water for Injection q.s. Acetic Acid may have  
been added for pH adjustment.  
**USUAL DOSAGE:** See package insert.  
Store at 20° to 25°C (68° to 77°F) [See USP  
Controlled Room Temperature].  
Do not permit to freeze.


Mfd by: HIKMA FARMACÊUTICA (PORTUGAL), SA  
Dist by: WEST-WARD PHARMACEUTICAL CORP.  
Eatontown, NJ 07724 USA

  
NDC 0143-9742-01  
PLB694-WES/1

Lot: \_\_\_\_\_  
Exp: \_\_\_\_\_

Non-varnish area

# SERIALIZATION IMAGE



## OXYTOCIN

oxytocin injection

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0143-9743
Route of Administration	INTRAMUSCULAR, INTRAVENOUS		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OXYTOCIN (UNII: 1JQS135EYN) (OXYTOCIN - UNII:1JQS135EYN)	OXYTOCIN	10 [USPU] in 1 mL

### Inactive Ingredients

Ingredient Name	Strength
CHLOROBUTANOL (UNII: HM4YQM8WRC)	500 mg in 1 mL
ACETIC ACID (UNII: Q40Q9N063P)	
WATER (UNII: 059QF0KO0R)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0143-9743-25	25 in 1 CARTON	02/13/2013	
1	NDC:0143-9743-01	1 mL in 1 VIAL; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA200219	02/13/2013	

## OXYTOCIN

oxytocin injection

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0143-9742
Route of Administration	INTRAMUSCULAR, INTRAVENOUS		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
OXYTOCIN (UNII: 1JQS135EYN) (OXYTOCIN - UNII:1JQS135EYN)	OXYTOCIN	10 [USPU] in 1 mL

**Inactive Ingredients**

Ingredient Name	Strength
CHLOROBUTANOL (UNII: HM4YQM8WRC)	500 mg in 1 mL
ACETIC ACID (UNII: Q40Q9N063P)	
WATER (UNII: 059QF0K00R)	

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0143-9742-25	25 in 1 CARTON	02/13/2013	
1	NDC:0143-9742-01	10 mL in 1 VIAL; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA200219	02/13/2013	

**Labeler** - West-Ward Pharmaceuticals Corp (001230762)**Establishment**

Name	Address	ID/FEI	Business Operations
HIKMA FARMACEUTICA (PORTUGAL), S.A		452742943	ANALYSIS(0143-9743, 0143-9742) , LABEL(0143-9743, 0143-9742) , MANUFACTURE(0143-9743, 0143-9742) , PACK(0143-9743, 0143-9742)

Revised: 12/2018

West-Ward Pharmaceuticals Corp