

GUAIFENESIN- guaifenesin tablet
RITE AID

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

272R RITE AID 11822 7836 GUAIFENESIN 400 mg TABLETS

DRUG FACTS

Active ingredient (in each immediate-release tablet)

Guaifenesin 400 mg

Purpose

Expectorant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive to rid the bronchial passageway of bothersome mucus

Warnings

Ask a doctor before use if you have

- persistent or chronic cough such as occurs with smoking , asthma, chronic bronchitis, or emphysema
- cough accompanied by excessive phlegm (mucus)

When using this product

- **do not exceed recommended dosage**
- do not use for more than 7 days

Stop use and ask a doctor if

- cough lasts for more than 7 days, recurs, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a healthcare professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- adults and children 12 years of age and older: take 1 tablet every 4 hours with a full glass of water while symptoms persist. Do not exceed 6 doses in 24 hours.
- children under 12 years of age: do not use

Other information

- store at 25°C (77°F) excursions permitted between 15°-30°C (59°- 86°F)
- keep in a dry place and do not expose to excessive heat

Inactive ingredients colloidal silicon dioxide, magnesium stearate, maltodextrin, microcrystalline cellulose, povidone, silicon dioxide, sodium starch glycolate, stearic acid

Questions or comments? Call 1-877-290-4008

NDC 11822-7836-5

MUCUS RELIEF

GUAIFENESIN 400 mg

EXPECTORANT

Relieves chest congestion
Thins & loosens mucus
Immediate-release



30
IMMEDIATE-RELEASE TABLETS

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Uses ■ helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive to rid the bronchial passageway of bothersome mucus

Warnings
Ask a doctor before use if you have
■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
■ cough accompanied by excessive phlegm (mucus)

When using this product ■ do not exceed recommended dosage ■ do not use for more than 7 days

Stop use and ask a doctor if
■ cough lasts for more than 7 days, recurs, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.
If pregnant or breast-feeding, ask a healthcare professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ adults and children 12 years of age and older: take 1 tablet every 4 hours with a full glass of water while symptoms persist. Do not exceed 6 doses in 24 hours.
■ children under 12 years of age: do not use

Other information ■ TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING ■ Store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ Keep in a dry place and do not expose to excessive heat


Questions or comments? Call 1-877-290-4008

DISTRIBUTION GUARANTEE:
RITE AID
If you're not satisfied, we'll happily refund your money.

30 HUNTER LANE
CAMP HILL, PA 17011
www.riteaid.com

272R.0821

Lot No.:
Exp. Date:



Drug Facts
Active ingredient (in each immediate-release tablet) Purpose
Guaifenesin 400 mg Expectorant

Uses
■ helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive to rid the bronchial passageway of bothersome mucus

Warnings
Ask a doctor before use if you have
■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema ■ cough accompanied by excessive phlegm (mucus)

When using this product ■ do not exceed recommended dosage ■ do not use for more than 7 days

Stop use and ask a doctor if
■ cough lasts for more than 7 days, recurs, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a healthcare professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-424-1222) right away.

Directions
■ adults and children 12 years of age and older: take 1 tablet every 4 hours with a full glass of water while symptoms persist. Do not exceed 6 doses in 24 hours.
■ children under 12 years of age: do not use

Other information
■ Store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
■ Keep in a dry place and do not expose to excessive heat

Inactive ingredients: colloidal silicon dioxide, magnesium stearate, maltodextrin, microcrystalline cellulose, povidone, silicon dioxide, sodium starch glycolate, stearic acid

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Questions or comments? call 1-877-290-4008
272R.0821

MUCUS RELIEF
GUAIFENESIN 400 mg
EXPECTORANT

NDC 11822-7836-5

MUCUS RELIEF
GUAIFENESIN 400 mg
EXPECTORANT

Relieves chest congestion
Thins & loosens mucus
Immediate-release

ACTUAL SIZE

30
IMMEDIATE-RELEASE TABLETS

MUCUS RELIEF
GUAIFENESIN 400 mg
EXPECTORANT
Relieves chest congestion
Thins & loosens mucus
Immediate-release

DISTRIBUTED BY:
RITE AID, 30 HUNTER LANE,
CAMP HILL, PA 17011
www.riteaid.com

SATISFACTION GUARANTEE
If you're not satisfied, we'll happily refund your money.

Lot No.:
Exp. Date:

0 11822 78935 4

GUAIFENESIN

guaifenesin tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-7836
Route of Administration	ORAL		

Active Ingredient/Active Moiety				
Ingredient Name			Basis of Strength	Strength
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)			GUAIFENESIN	400 mg
Inactive Ingredients				
Ingredient Name			Strength	
MALTODEXTRIN (UNII: 7CVR7L4A2D)				
MAGNESIUM STEARATE (UNII: 70097M6I30)				
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)				
STEARIC ACID (UNII: 4ELV7Z65AP)				
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)				
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
POVIDONE (UNII: FZ989GH94E)				
Product Characteristics				
Color	white	Score	2 pieces	
Shape	ROUND	Size	12mm	
Flavor		Imprint Code	TCL272	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-7836-5	1 in 1 CARTON	12/27/2021	
1		30 in 1 BOTTLE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph final	part341	12/27/2021		

Labeler - RITE AID (014578892)

Registrant - TIME CAP LABORATORIES INC (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LTD.		925822975	manufacture(11822-7836)