

LIQUID WORMER 2X- pyrantel pamoate suspension
Durvet Inc

Liquid
Wormer
2X

INDICATIONS FOR USE

To prevent reinfestation of (Toxocara canis) in puppies and adult dogs and in lactating bitches after whelping.

For the removal of large roundworms, (Toxocara canis and Toxascaris leonina) and hookworms, (Ancylostoma caninum and Uncinaria stenocephala) in dogs and puppies.

Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

WARNING

KEEP OUT OF REACH OF CHILDREN

Read Accompanying Professional Information

Recommended Storage: Store Below 30°C (86°F)

Durvet logo
www.durvet.com

Manufactured for Durvet, Inc.
100 S.E. Magellan Drive,
Blue Springs, Missouri 64014
ISS12XB04

IMPORTANT: SHAKE WELL BEFORE USING

DIRECTIONS FOR USE

For maximum control and prevention of reinfestation, it is recommended that puppies be treated at 2, 3, 4, 6, 8 and 10 weeks of age. Lactating bitches should be treated 2-3 weeks after whelping. Adult dogs kept in heavily contaminated quarters may be treated at monthly intervals to prevent T. canis reinfestation. Administer one full teaspoonful (5 mL) for each 10 lbs. of body weight. For the removal of large roundworms (ascarids) and hookworms. Administer one full teaspoonful (5 mL) for each 10 lbs. of body weight. It is not necessary to withhold food prior to treatment. Dogs usually find this wormer very palatable and will lick the dose from the bowl willingly. If there is a reluctance to accept the dose, mix in a small quantity of dog food to encourage consumption. The presence of large roundworms (ascarids) and hookworms should be confirmed by lab fecal examination. It is recommended that dogs maintained under conditions of constant exposure to worm infestation should have a followup fecal exam within 2 to 4 weeks after first treatment.

Take Time
eye symbol
Observe Label Directions

PRINCIPAL DISPLAY PANEL - 60 mL Bottle Label




Durvet logo

NDC 30798-760-22

Liquid
Wormer
(pyrantel pamoate) 2X

Canine Anthelmintic
Suspension

Active Ingredient: 4.54 mg of
pyrantel base as pyrantel pamoate per mL
FOR ANIMAL USE ONLY
ANADA 200-007 APPROVED BY FDA
NET CONTENTS: 2 FL OZ (60 ML)

<p>INDICATIONS FOR USE</p> <p>To prevent reinfestation of (<i>Toxocara canis</i>) in puppies and adult dogs and in lactating bitches after whelping.</p> <p>For the removal of large roundworms, (<i>Toxocara canis</i> and <i>Toxascaris leonina</i>) and hookworms, (<i>Ancylostoma caninum</i> and <i>Uncinaria stenocephala</i>) in dogs and puppies.</p> <p>Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.</p> <p>WARNING: KEEP OUT OF REACH OF CHILDREN Read Accompanying Professional Information</p> <p>Recommended Storage: Store Below 30°C (86°F)</p> <p> Manufactured for Durvet, Inc. 100 S.E. Magellan Drive, Blue Springs, Missouri 64014 ISS12XB04</p>	<p>NDC 30798-760-22</p> <p> Liquid Wormer 2X (pyrantel pamoate)</p> <p>Canine Anthelmintic Suspension</p> <p>Active Ingredient: 4.54 mg of pyrantel base as pyrantel pamoate per mL FOR ANIMAL USE ONLY ANADA 200-007 APPROVED BY FDA NET CONTENTS: 2 FL OZ (60 ML)</p>	<p>IMPORTANT: SHAKE WELL BEFORE USING</p> <p>DIRECTIONS FOR USE: For maximum control and prevention of reinfestation, it is recommended that puppies be treated at 2, 3, 4, 6, 8 and 10 weeks of age. Lactating bitches should be treated 2-3 weeks after whelping. Adult dogs kept in heavily contaminated quarters may be treated at monthly intervals to prevent <i>T. canis</i> reinfestation. Administer one full teaspoonful (5 mL) for each 10 lbs. of body weight.</p> <p>For the removal of large roundworms (ascarids) and hookworms, Administer one full teaspoonful (5 mL) for each 10 lbs. of body weight. It is not necessary to withhold food prior to treatment. Dogs usually find this wormer very palatable and will lick the dose from the bowl willingly. If there is a reluctance to accept the dose, mix in a small quantity of dog food to encourage consumption. The presence of large roundworms (ascarids) and hookworms should be confirmed by lab fecal examination.</p> <p>It is recommended that dogs maintained under conditions of constant exposure to worm infestation should have a follow-up fecal exam within 2 to 4 weeks after first treatment.</p> <p>Take Time  Observe Label Directions</p> <p>IMPRINT AREA</p>
--	--	--

LIQUID WORMER 2X

pyrantel pamoate suspension

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:30798-760
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
pyrantel pamoate (UNII: 81BK194Z5M) (pyrantel - UNII:4QIH0N49E7)	pyrantel	4.54 mg in 1 mL

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:30798-760-22	60 mL in 1 BOTTLE, PLASTIC		
2	NDC:30798-760-28	237 mL in 1 BOTTLE, PLASTIC		
3	NDC:30798-760-31	473 mL in 1 BOTTLE, PLASTIC		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANADA	ANADA200007	08/24/2012	

Labeler - Durvet Inc (056387798)

Registrant - Happy Jack Inc (003189354)

Establishment

Name	Address	ID/FEI	Business Operations
ABS Corporation		048022727	MANUFACTURE

Establishment

Name	Address	ID/FEI	Business Operations
COSMA Spa		428655732	API MANUFACTURE

Revised: 7/2013

Durvet Inc