

# MOUNJARO- tirzepatide injection, solution

## Eli Lilly and Company

-----

### HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use MOUNJARO safely and effectively. See full prescribing information for MOUNJARO.

MOUNJARO® (tirzepatide) Injection, for subcutaneous use  
Initial U.S. Approval: 2022

#### WARNING: RISK OF THYROID C-CELL TUMORS

*See full prescribing information for complete boxed warning.*

- Tirzepatide causes thyroid C-cell tumors in rats. It is unknown whether MOUNJARO causes thyroid C-cell tumors, including medullary thyroid carcinoma (MTC), in humans as the human relevance of tirzepatide-induced rodent thyroid C-cell tumors has not been determined (5.1, 13.1).
- MOUNJARO is contraindicated in patients with a personal or family history of MTC or in patients with Multiple Endocrine Neoplasia syndrome type 2 (MEN 2). Counsel patients regarding the potential risk of MTC and symptoms of thyroid tumors (4, 5.1).

#### RECENT MAJOR CHANGES

-----

##### Warnings and Precautions

Severe Gastrointestinal Adverse Reactions (5.6)

11/2024

##### Warnings and Precautions

Pulmonary Aspiration During General Anesthesia or Deep Sedation (5.9)

11/2024

#### INDICATIONS AND USAGE

-----

MOUNJARO® is a glucose-dependent insulinotropic polypeptide (GIP) receptor and glucagon-like peptide-1 (GLP-1) receptor agonist indicated as an adjunct to diet and exercise to improve glycemic control in adults with type 2 diabetes mellitus. (1)

#### DOSAGE AND ADMINISTRATION

-----

- The recommended starting dosage is 2.5 mg injected subcutaneously once weekly. (2.1)
- After 4 weeks, increase to 5 mg injected subcutaneously once weekly. (2.1)
- If additional glycemic control is needed, increase the dosage in 2.5 mg increments after at least 4 weeks on the current dose. (2.1)
- The maximum dosage is 15 mg subcutaneously once weekly. (2.1)
- Administer once weekly at any time of day, with or without meals. (2.2)
- Inject subcutaneously in the abdomen, thigh, or upper arm. (2.2)
- Rotate injection sites with each dose. (2.2)

#### DOSAGE FORMS AND STRENGTHS

-----

Injection: 2.5 mg, 5 mg, 7.5 mg, 10 mg, 12.5 mg, or 15 mg per 0.5 mL in single-dose pen or single-dose vial (3)

#### CONTRAINDICATIONS

-----

- Personal or family history of medullary thyroid carcinoma or in patients with Multiple Endocrine Neoplasia syndrome type 2 (4)
- Known serious hypersensitivity to tirzepatide or any of the excipients in MOUNJARO (4)

#### WARNINGS AND PRECAUTIONS

-----

- *Acute Pancreatitis*: Has been observed in patients treated with GLP-1 receptor agonists, or MOUNJARO. Discontinue if pancreatitis is suspected. (5.2)
- *Hypoglycemia with Concomitant Use of Insulin Secretagogues or Insulin*: Concomitant use with an insulin secretagogue or insulin may increase the risk of hypoglycemia, including severe hypoglycemia. Reducing dose of insulin secretagogue or insulin may be necessary. (5.3)

- *Hypersensitivity Reactions:* Serious hypersensitivity reactions (e.g., anaphylaxis and angioedema) have been reported. Discontinue MOUNJARO if suspected and promptly seek medical advice. (5.4)
- *Acute Kidney Injury Due to Volume Depletion:* Monitor renal function in patients reporting adverse reactions that could lead to volume depletion. (5.5)
- *Severe Gastrointestinal Adverse Reactions:* Use may be associated with gastrointestinal adverse reactions, sometimes severe. MOUNJARO is not recommended in patients with severe gastroparesis. (5.6)
- *Diabetic Retinopathy Complications in Patients with a History of Diabetic Retinopathy:* Has not been studied in patients with non-proliferative diabetic retinopathy requiring acute therapy, proliferative diabetic retinopathy, or diabetic macular edema. Monitor patients with a history of diabetic retinopathy for progression. (5.7)
- *Acute Gallbladder Disease:* Has occurred in clinical trials. If cholelithiasis is suspected, gallbladder studies and clinical follow-up are indicated. (5.8)
- *Pulmonary Aspiration During General Anesthesia or Deep Sedation:* Has been reported in patients receiving GLP-1 receptor agonists undergoing elective surgeries or procedures. Instruct patients to inform healthcare providers of any planned surgeries or procedures. (5.9)

#### ADVERSE REACTIONS

The most common adverse reactions, reported in  $\geq 5\%$  of patients treated with MOUNJARO are nausea, diarrhea, decreased appetite, vomiting, constipation, dyspepsia, and abdominal pain. (6.1)

**To report SUSPECTED ADVERSE REACTIONS, contact Eli Lilly and Company at 1-800-LillyRx (1-800-545-5979) or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

#### DRUG INTERACTIONS

MOUNJARO delays gastric emptying and has the potential to impact the absorption of concomitantly administered oral medications. (7.2)

#### USE IN SPECIFIC POPULATIONS

- *Pregnancy:* Based on animal studies, may cause fetal harm. (8.1)
- *Females of Reproductive Potential:* Advise females using oral contraceptives to switch to a non-oral contraceptive method or add a barrier method of contraception for 4 weeks after initiation and for 4 weeks after each dose escalation. (8.3)

**See 17 for PATIENT COUNSELING INFORMATION and Medication Guide.**

**Revised: 6/2025**

## FULL PRESCRIBING INFORMATION: CONTENTS\*

### WARNING: RISK OF THYROID C-CELL TUMORS

#### 1 INDICATIONS AND USAGE

#### 2 DOSAGE AND ADMINISTRATION

##### 2.1 Dosage

##### 2.2 Important Administration Instructions

#### 3 DOSAGE FORMS AND STRENGTHS

#### 4 CONTRAINDICATIONS

#### 5 WARNINGS AND PRECAUTIONS

##### 5.1 Risk of Thyroid C-Cell Tumors

##### 5.2 Acute Pancreatitis

##### 5.3 Hypoglycemia with Concomitant Use of Insulin Secretagogues or Insulin

##### 5.4 Hypersensitivity Reactions

##### 5.5 Acute Kidney Injury Due to Volume Depletion

##### 5.6 Severe Gastrointestinal Adverse Reactions

##### 5.7 Diabetic Retinopathy Complications in Patients with a History of Diabetic Retinopathy

##### 5.8 Acute Gallbladder Disease

5.9 Pulmonary Aspiration During General Anesthesia or Deep Sedation

## **6 ADVERSE REACTIONS**

6.1 Clinical Trials Experience

6.2 Postmarketing Experience

## **7 DRUG INTERACTIONS**

7.1 Concomitant Use with an Insulin Secretagogue (e.g., Sulfonylurea) or with Insulin

7.2 Oral Medications

## **8 USE IN SPECIFIC POPULATIONS**

8.1 Pregnancy

8.2 Lactation

8.3 Females and Males of Reproductive Potential

8.4 Pediatric Use

8.5 Geriatric Use

8.6 Renal Impairment

8.7 Hepatic Impairment

## **10 OVERDOSAGE**

## **11 DESCRIPTION**

## **12 CLINICAL PHARMACOLOGY**

12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

12.6 Immunogenicity

## **13 NONCLINICAL TOXICOLOGY**

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

## **14 CLINICAL STUDIES**

14.1 Overview of Clinical Studies

14.2 Monotherapy Use of MOUNJARO in Adult Patients with Type 2 Diabetes Mellitus

14.3 MOUNJARO Use in Combination with Metformin, Sulfonylureas, and/or SGLT2 Inhibitors in Adult Patients with Type 2 Diabetes Mellitus

14.4 MOUNJARO Use in Combination with Basal Insulin with or without Metformin in Adult Patients with Type 2 Diabetes Mellitus

## **16 HOW SUPPLIED/STORAGE AND HANDLING**

16.1 How Supplied

16.2 Storage and Handling

## **17 PATIENT COUNSELING INFORMATION**

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## **FULL PRESCRIBING INFORMATION**

### **WARNING: RISK OF THYROID C-CELL TUMORS**

- In both male and female rats, tirzepatide causes dose-dependent and treatment-duration-dependent thyroid C-cell tumors at clinically relevant exposures. It is unknown whether MOUNJARO causes thyroid C-cell tumors, including medullary thyroid carcinoma (MTC), in humans as human relevance of tirzepatide-induced rodent thyroid C-cell tumors has not been determined [*see Warnings and Precautions (5.1) and Nonclinical Toxicology (13.1)*].
- MOUNJARO is contraindicated in patients with a personal or family history of MTC or in patients with Multiple Endocrine Neoplasia syndrome type 2 (MEN 2) [*see Contraindications (4)*]. Counsel patients regarding the potential risk for MTC with the use of MOUNJARO and inform them of symptoms of thyroid tumors (e.g., a mass in the neck, dysphagia, dyspnea, persistent hoarseness). Routine monitoring of serum calcitonin or using thyroid ultrasound is of uncertain value for early detection of MTC in patients treated with MOUNJARO [*see Contraindications (4) and Warnings and Precautions (5.1)*].

## **1 INDICATIONS AND USAGE**

MOUNJARO® is indicated as an adjunct to diet and exercise to improve glycemic control in adults with type 2 diabetes mellitus.

## **2 DOSAGE AND ADMINISTRATION**

### **2.1 Dosage**

- The recommended starting dosage of MOUNJARO is 2.5 mg injected subcutaneously once weekly. Follow the dosage escalation below to reduce the risk of gastrointestinal adverse reactions [*see Warnings and Precautions (5.6) and Adverse Reactions (6.1)*]. The 2.5 mg dosage is for treatment initiation and is not intended for glycemic control.
- After 4 weeks, increase the dosage to 5 mg injected subcutaneously once weekly.
- If additional glycemic control is needed, increase the dosage in 2.5 mg increments after at least 4 weeks on the current dose.
- The maximum dosage of MOUNJARO is 15 mg injected subcutaneously once weekly.
- If a dose is missed, instruct patients to administer MOUNJARO as soon as possible within 4 days (96 hours) after the missed dose. If more than 4 days have passed, skip the missed dose and administer the next dose on the regularly scheduled day. In each case, patients can then resume their regular once weekly dosing schedule.
- The day of weekly administration can be changed, if necessary, as long as the time between the two doses is at least 3 days (72 hours).

### **2.2 Important Administration Instructions**

- Prior to initiation, train patients and caregivers on proper injection technique [*see Instructions for Use*].
- Instruct patients using the single-dose vial to use a syringe appropriate for dose administration (e.g., a 1 mL syringe capable of measuring a 0.5 mL dose).

- Administer MOUNJARO once weekly, any time of day, with or without meals.
- Inject MOUNJARO subcutaneously in the abdomen, thigh, or upper arm.
- Rotate injection sites with each dose.
- Inspect MOUNJARO visually before use. It should appear clear and colorless to slightly yellow. Do not use MOUNJARO if particulate matter or discoloration is seen.
- When using MOUNJARO with insulin, administer as separate injections and never mix. It is acceptable to inject MOUNJARO and insulin in the same body region, but the injections should not be adjacent to each other.

### 3 DOSAGE FORMS AND STRENGTHS

Injection: Clear, colorless to slightly yellow solution in pre-filled single-dose pens or single-dose vials, each available in the following strengths:

- 2.5 mg/0.5 mL
- 5 mg/0.5 mL
- 7.5 mg/0.5 mL
- 10 mg/0.5 mL
- 12.5 mg/0.5 mL
- 15 mg/0.5 mL

### 4 CONTRAINDICATIONS

MOUNJARO is contraindicated in patients with:

- A personal or family history of medullary thyroid carcinoma (MTC) or in patients with Multiple Endocrine Neoplasia syndrome type 2 (MEN 2) [*see Warnings and Precautions (5.1)*].
- Known serious hypersensitivity to tirzepatide or any of the excipients in MOUNJARO. Serious hypersensitivity reactions, including anaphylaxis and angioedema, have been reported with MOUNJARO [*see Warnings and Precautions (5.4)*].

### 5 WARNINGS AND PRECAUTIONS

#### 5.1 Risk of Thyroid C-Cell Tumors

In both sexes of rats, tirzepatide caused a dose-dependent and treatment-duration-dependent increase in the incidence of thyroid C-cell tumors (adenomas and carcinomas) in a 2-year study at clinically relevant plasma exposures [*see Nonclinical Toxicology (13.1)*]. It is unknown whether MOUNJARO causes thyroid C-cell tumors, including medullary thyroid carcinoma (MTC), in humans as human relevance of tirzepatide-induced rodent thyroid C-cell tumors has not been determined.

MOUNJARO is contraindicated in patients with a personal or family history of MTC or in patients with MEN 2. Counsel patients regarding the potential risk for MTC with the use of MOUNJARO and inform them of symptoms of thyroid tumors (e.g., a mass in the neck, dysphagia, dyspnea, persistent hoarseness).

Routine monitoring of serum calcitonin or using thyroid ultrasound is of uncertain value for early detection of MTC in patients treated with MOUNJARO. Such monitoring may increase the risk of unnecessary procedures, due to the low test specificity for serum

calcitonin and a high background incidence of thyroid disease. Significantly elevated serum calcitonin values may indicate MTC and patients with MTC usually have calcitonin values >50 ng/L. If serum calcitonin is measured and found to be elevated, the patient should be further evaluated. Patients with thyroid nodules noted on physical examination or neck imaging should also be further evaluated.

## **5.2 Acute Pancreatitis**

Acute pancreatitis, including fatal and non-fatal hemorrhagic or necrotizing pancreatitis, has been observed in patients treated with GLP-1 receptor agonists, or MOUNJARO [see *Adverse Reactions (6.1)*].

After initiation of MOUNJARO, observe patients carefully for signs and symptoms of pancreatitis (including persistent severe abdominal pain, sometimes radiating to the back and which may or may not be accompanied by vomiting). If pancreatitis is suspected, discontinue MOUNJARO and initiate appropriate management.

## **5.3 Hypoglycemia with Concomitant Use of Insulin Secretagogues or Insulin**

Patients receiving MOUNJARO in combination with an insulin secretagogue (e.g., sulfonylurea) or insulin may have an increased risk of hypoglycemia, including severe hypoglycemia [see *Adverse Reactions (6.1)*, *Drug Interactions (7.1)*].

The risk of hypoglycemia may be lowered by a reduction in the dose of sulfonylurea (or other concomitantly administered insulin secretagogue) or insulin. Inform patients using these concomitant medications of the risk of hypoglycemia and educate them on the signs and symptoms of hypoglycemia.

## **5.4 Hypersensitivity Reactions**

Serious hypersensitivity reactions (e.g., anaphylaxis, angioedema) have been reported in patients treated with MOUNJARO. If hypersensitivity reactions occur, discontinue use of MOUNJARO; treat promptly per standard of care, and monitor until signs and symptoms resolve. Do not use in patients with a previous serious hypersensitivity reaction to tirzepatide or any of the excipients in MOUNJARO [see *Contraindications (4)*, *Adverse Reactions (6.2)*].

Anaphylaxis and angioedema have been reported with GLP-1 receptor agonists. Use caution in patients with a history of angioedema or anaphylaxis with a GLP-1 receptor agonist because it is unknown whether such patients will be predisposed to these reactions with MOUNJARO.

## **5.5 Acute Kidney Injury Due to Volume Depletion**

There have been postmarketing reports of acute kidney injury, in some cases requiring hemodialysis, in patients treated with GLP-1 receptor agonists, or MOUNJARO [see *Adverse Reactions (6.2)*]. The majority of the reported events occurred in patients who experienced gastrointestinal adverse reactions leading to dehydration such as nausea, vomiting, or diarrhea [see *Adverse Reactions (6.1)*]. Monitor renal function in patients reporting adverse reactions to MOUNJARO that could lead to volume depletion, especially during dosage initiation and escalation of MOUNJARO.

## **5.6 Severe Gastrointestinal Adverse Reactions**

Use of MOUNJARO has been associated with gastrointestinal adverse reactions, sometimes severe [see *Adverse Reactions (6)*]. In the pool of placebo-controlled trials, severe gastrointestinal adverse reactions occurred more frequently among patients receiving MOUNJARO (5 mg 1.3%, 10 mg 0.4%, 15 mg 1.2%) than placebo (0.9%).

MOUNJARO is not recommended in patients with severe gastroparesis.

### **5.7 Diabetic Retinopathy Complications in Patients with a History of Diabetic Retinopathy**

Rapid improvement in glucose control has been associated with a temporary worsening of diabetic retinopathy. MOUNJARO has not been studied in patients with non-proliferative diabetic retinopathy requiring acute therapy, proliferative diabetic retinopathy, or diabetic macular edema. Patients with a history of diabetic retinopathy should be monitored for progression of diabetic retinopathy.

### **5.8 Acute Gallbladder Disease**

Acute events of gallbladder disease such as cholelithiasis or cholecystitis have been reported in GLP-1 receptor agonist trials and postmarketing.

In MOUNJARO placebo-controlled clinical trials, acute gallbladder disease (cholelithiasis, biliary colic, and cholecystectomy) was reported by 0.6% of MOUNJARO-treated patients and 0% of placebo-treated patients. If cholelithiasis is suspected, gallbladder diagnostic studies and appropriate clinical follow-up are indicated.

### **5.9 Pulmonary Aspiration During General Anesthesia or Deep Sedation**

MOUNJARO delays gastric emptying [see *Clinical Pharmacology (12.2)*]. There have been rare postmarketing reports of pulmonary aspiration in patients receiving GLP-1 receptor agonists undergoing elective surgeries or procedures requiring general anesthesia or deep sedation who had residual gastric contents despite reported adherence to preoperative fasting recommendations.

Available data are insufficient to inform recommendations to mitigate the risk of pulmonary aspiration during general anesthesia or deep sedation in patients taking MOUNJARO, including whether modifying preoperative fasting recommendations or temporarily discontinuing MOUNJARO could reduce the incidence of retained gastric contents. Instruct patients to inform healthcare providers prior to any planned surgeries or procedures if they are taking MOUNJARO.

## **6 ADVERSE REACTIONS**

The following serious adverse reactions are described below or elsewhere in the prescribing information:

- Risk of Thyroid C-cell Tumors [see *Warnings and Precautions (5.1)*]
- Acute Pancreatitis [see *Warnings and Precautions (5.2)*]
- Hypoglycemia with Concomitant Use of Insulin Secretagogues or Insulin [see *Warnings and Precautions (5.3)*]
- Hypersensitivity Reactions [see *Warnings and Precautions (5.4)*]
- Acute Kidney Injury Due to Volume Depletion [see *Warnings and Precautions (5.5)*]
- Severe Gastrointestinal Adverse Reactions [see *Warnings and Precautions (5.6)*]

- Diabetic Retinopathy Complications in Patients with a History of Diabetic Retinopathy [see Warnings and Precautions (5.7)]
- Acute Gallbladder Disease [see Warnings and Precautions (5.8)]
- Pulmonary Aspiration During General Anesthesia or Deep Sedation [see Warnings and Precautions (5.9)]

## 6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

### Pool of Two Placebo-Controlled Clinical Trials

The data in Table 1 are derived from 2 placebo-controlled trials [1 monotherapy trial (SURPASS-1) and 1 trial in combination with basal insulin with or without metformin (SURPASS-5)] in adult patients with type 2 diabetes mellitus [see Clinical Studies (14.2, 14.4)]. These data reflect exposure of 718 patients to MOUNJARO and a mean duration of exposure to MOUNJARO of 36.6 weeks. The mean age of patients was 58 years, 4% were 75 years or older and 54% were male. The population was 57% White, 27% Asian, 13% American Indian or Alaska Native, and 3% Black or African American; 25% identified as Hispanic or Latino ethnicity. At baseline, patients had type 2 diabetes mellitus for an average of 9.1 years with a mean HbA1c of 8.1%. As assessed by baseline fundoscopic examination, 13% of the population had retinopathy. At baseline, eGFR was  $\geq 90$  mL/min/1.73 m<sup>2</sup> in 53%, 60 to 90 mL/min/1.73 m<sup>2</sup> in 39%, 45 to 60 mL/min/1.73 m<sup>2</sup> in 7%, and 30 to 45 mL/min/1.73 m<sup>2</sup> in 1% of patients.

### Pool of Seven Controlled Clinical Trials

Adverse reactions were also evaluated in a larger pool of adult patients with type 2 diabetes mellitus participating in seven controlled clinical trials which included two placebo-controlled trials (SURPASS-1 and -5), three trials of MOUNJARO in combination with metformin, sulfonylureas, and/or SGLT2 Inhibitors (SURPASS-2, -3, -4) [see Clinical Studies (14.3)] and two additional trials conducted in Japan. In this pool, a total of 5119 adult patients with type 2 diabetes mellitus were treated with MOUNJARO for a mean duration of 48.1 weeks. The mean age of patients was 58 years, 4% were 75 years or older and 58% were male. The population was 65% White, 24% Asian, 7% American Indian or Alaska Native, and 3% Black or African American; 38% identified as Hispanic or Latino ethnicity. At baseline, patients had type 2 diabetes mellitus for an average of 9.1 years with a mean HbA1c of 8.3%. As assessed by baseline fundoscopic examination, 15% of the population had retinopathy. At baseline, eGFR was  $\geq 90$  mL/min/1.73 m<sup>2</sup> in 52%, 60 to 90 mL/min/1.73 m<sup>2</sup> in 40%, 45 to 60 mL/min/1.73 m<sup>2</sup> in 6%, and 30 to 45 mL/min/1.73 m<sup>2</sup> in 1% of patients.

### Common Adverse Reactions

Table 1 shows common adverse reactions, not including hypoglycemia, associated with the use of MOUNJARO in the pool of placebo-controlled trials. These adverse reactions occurred more commonly on MOUNJARO than on placebo and occurred in at least 5% of patients treated with MOUNJARO.

**Table 1: Adverse Reactions in Pool of Placebo-Controlled Trials Reported in  $\geq 5\%$  of MOUNJARO-treated Adult Patients with Type 2 Diabetes Mellitus**



<b>Adverse Reaction</b>	<b>Placebo (N=235) %</b>	<b>MOUNJARO 5 mg (N=237) %</b>	<b>MOUNJARO 10 mg (N=240) %</b>	<b>MOUNJARO 15 mg (N=241) %</b>
Nausea	4	12	15	18
Diarrhea	9	12	13	17
Decreased Appetite	1	5	10	11
Vomiting	2	5	5	9
Constipation	1	6	6	7
Dyspepsia	3	8	8	5
Abdominal Pain	4	6	5	5

Note: Percentages reflect the number of patients who reported at least 1 occurrence of the adverse reaction.

In the pool of seven clinical trials, the types and frequency of common adverse reactions, not including hypoglycemia, were similar to those listed in Table 1.

#### *Gastrointestinal Adverse Reactions*

In the pool of placebo-controlled trials, gastrointestinal adverse reactions occurred more frequently among patients receiving MOUNJARO than placebo (placebo 20.4%, MOUNJARO 5 mg 37.1%, MOUNJARO 10 mg 39.6%, MOUNJARO 15 mg 43.6%). More patients receiving MOUNJARO 5 mg (3.0%), MOUNJARO 10 mg (5.4%), and MOUNJARO 15 mg (6.6%) discontinued treatment due to gastrointestinal adverse reactions than patients receiving placebo (0.4%). The majority of reports of nausea, vomiting, and/or diarrhea occurred during dose escalation and decreased over time.

The following gastrointestinal adverse reactions were reported more frequently in MOUNJARO-treated patients than placebo-treated patients (frequencies listed, respectively, as: placebo; 5 mg; 10 mg; 15 mg): eructation (0.4%, 3.0%, 2.5%, 3.3%), flatulence (0%, 1.3%, 2.5%, 2.9%), gastroesophageal reflux disease (0.4%, 1.7%, 2.5%, 1.7%), abdominal distension (0.4%, 0.4%, 2.9%, 0.8%).

#### Other Adverse Reactions

##### *Hypoglycemia*

Table 2 summarizes the incidence of hypoglycemic events in the placebo-controlled trials.

**Table 2: Hypoglycemia Adverse Reactions in Placebo-Controlled Trials in Adult Patients with Type 2 Diabetes Mellitus**

	<b>Placebo %</b>	<b>MOUNJARO 5 mg %</b>	<b>MOUNJARO 10 mg %</b>	<b>MOUNJARO 15 mg %</b>
Monotherapy (40 weeks)*	N=115	N=121	N=119	N=120
Blood glucose <54 mg/dL	1	0	0	0
Severe hypoglycemia**	0	0	0	0

Add-on to Basal Insulin with or without Metformin				
(40 weeks)*	N=120	N=116	N=119	N=120
Blood glucose <54 mg/dL	13	16	19	14
Severe hypoglycemia**	0	0	2	1

\* Reflects the study treatment period. Data include events occurring during 4 weeks of treatment-free safety follow up. Events after introduction of a new glucose-lowering treatment are excluded.

\*\* Episodes requiring the assistance of another person to actively administer carbohydrate, glucagon, or other resuscitative actions.

Hypoglycemia was more frequent when MOUNJARO was used in combination with a sulfonylurea [see *Clinical Studies (14)*]. In a clinical trial up to 104 weeks of treatment, when administered with a sulfonylurea, hypoglycemia (glucose level <54 mg/dL) occurred in 13.8%, 9.9%, and 12.8%, and severe hypoglycemia occurred in 0.5%, 0%, and 0.6% of patients treated with MOUNJARO 5 mg, 10 mg, and 15 mg, respectively.

### *Acute Pancreatitis*

In clinical studies, 14 events of acute pancreatitis were confirmed by adjudication in 13 MOUNJARO-treated patients (0.23 patients per 100 years of exposure) versus 3 events in 3 comparator-treated patients (0.11 patients per 100 years of exposure).

### *Heart Rate Increase*

In the pool of placebo-controlled trials, treatment with MOUNJARO resulted in a mean increase in heart rate of 2 to 4 beats per minute compared to a mean increase of 1 beat per minute in placebo-treated patients. Episodes of sinus tachycardia, associated with a concomitant increase from baseline in heart rate of  $\geq 15$  beats per minute, also were reported in 4.3%, 4.6%, 5.9% and 10% of subjects treated with placebo, MOUNJARO 5 mg, 10 mg, and 15 mg, respectively. For patients enrolled in Japan, these episodes were reported in 7% (3/43), 7.1% (3/42), 9.3% (4/43), and 23% (10/43) of patients treated with placebo, MOUNJARO 5 mg, 10 mg, and 15 mg, respectively. The clinical relevance of heart rate increases is uncertain.

### *Hypersensitivity Reactions*

Hypersensitivity reactions have been reported with MOUNJARO in the pool of placebo-controlled trials, sometimes severe (e.g., urticaria and eczema); hypersensitivity reactions were reported in 3.2% of MOUNJARO-treated patients compared to 1.7% of placebo-treated patients.

In the pool of seven clinical trials, hypersensitivity reactions occurred in 106/2,570 (4.1%) of MOUNJARO-treated patients with anti-tirzepatide antibodies and in 73/2,455 (3.0%) of MOUNJARO-treated patients who did not develop anti-tirzepatide antibodies [see *Clinical Pharmacology (12.6)*].

### *Injection Site Reactions*

In the pool of placebo-controlled trials, injection site reactions were reported in 3.2% of MOUNJARO-treated patients compared to 0.4% of placebo-treated patients.

In the pool of seven clinical trials, injection site reactions occurred in 119/2,570 (4.6%) of MOUNJARO-treated patients with anti-tirzepatide antibodies and in 18/2,455 (0.7%) of

MOUNJARO-treated patients who did not develop anti-tirzepatide antibodies [see *Clinical Pharmacology* (12.6)].

#### *Acute Gallbladder Disease*

In the pool of placebo-controlled clinical trials, acute gallbladder disease (cholelithiasis, biliary colic and cholecystectomy) was reported by 0.6% of MOUNJARO-treated patients and 0% of placebo-treated patients.

#### *Dysesthesia*

In the pool of placebo-controlled clinical trials, dysesthesia was reported by 0.4%, 0.4%, and 0.4% of patients treated with MOUNJARO 5 mg, 10 mg, and 15 mg, respectively. No events were reported by patients receiving placebo.

#### *Dysgeusia*

In the pool of placebo-controlled clinical trials, dysgeusia was reported by 0.1% of MOUNJARO-treated patients and 0% of placebo-treated patients.

#### Laboratory Abnormalities

##### *Amylase and Lipase Increase*

In the pool of placebo-controlled clinical trials, treatment with MOUNJARO resulted in mean increases from baseline in serum pancreatic amylase concentrations of 33% to 38% and serum lipase concentrations of 31% to 42%. Placebo-treated patients had a mean increase from baseline in pancreatic amylase of 4% and no changes were observed in lipase. The clinical significance of elevations in lipase or amylase with MOUNJARO is unknown in the absence of other signs and symptoms of pancreatitis.

## **6.2 Postmarketing Experience**

The following adverse reactions have been reported during post-approval use of MOUNJARO. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

*Hypersensitivity:* anaphylaxis, angioedema

*Gastrointestinal:* acute pancreatitis, hemorrhagic and necrotizing pancreatitis sometimes resulting in death, ileus

*Pulmonary:* Pulmonary aspiration has occurred in patients receiving GLP-1 receptor agonists undergoing elective surgeries or procedures requiring general anesthesia or deep sedation

*Renal:* acute renal failure or worsening of chronic renal failure, sometimes requiring hemodialysis

*Skin and Subcutaneous Tissue:* alopecia

## **7 DRUG INTERACTIONS**

### **7.1 Concomitant Use with an Insulin Secretagogue (e.g., Sulfonylurea) or with Insulin**

When initiating MOUNJARO, consider reducing the dose of concomitantly administered insulin secretagogues (e.g., sulfonylureas) or insulin to reduce the risk of hypoglycemia [see *Warnings and Precautions* (5.3)].

## 7.2 Oral Medications

MOUNJARO delays gastric emptying and thereby has the potential to impact the absorption of concomitantly administered oral medications. Caution should be exercised when oral medications are concomitantly administered with MOUNJARO.

Monitor patients on oral medications dependent on threshold concentrations for efficacy and those with a narrow therapeutic index (e.g., warfarin) when concomitantly administered with MOUNJARO.

Advise patients using oral hormonal contraceptives to switch to a non-oral contraceptive method or add a barrier method of contraception for 4 weeks after initiation and for 4 weeks after each dose escalation with MOUNJARO. Hormonal contraceptives that are not administered orally should not be affected [see *Use in Specific Populations* (8.3) and *Clinical Pharmacology* (12.2, 12.3)].

## 8 USE IN SPECIFIC POPULATIONS

### 8.1 Pregnancy

#### Risk Summary

Available data with MOUNJARO use in pregnant women are insufficient to evaluate for a drug-related risk of major birth defects, miscarriage, or other adverse maternal or fetal outcomes. There are risks to the mother and fetus associated with poorly controlled diabetes in pregnancy (see *Clinical Considerations*). Based on animal reproduction studies, there may be risks to the fetus from exposure to tirzepatide during pregnancy. MOUNJARO should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

In pregnant rats administered tirzepatide during organogenesis, fetal growth reductions and fetal abnormalities occurred at clinical exposure in maternal rats based on AUC. In rabbits administered tirzepatide during organogenesis, fetal growth reductions were observed at clinically relevant exposures based on AUC. These adverse embryo/fetal effects in animals coincided with pharmacological effects on maternal weight and food consumption (see *Data*).

The estimated background risk of major birth defects is 6–10% in women with pre-gestational diabetes with an HbA1c >7% and has been reported to be as high as 20–25% in women with an HbA1c >10%. The estimated background risk of miscarriage for the indicated population is unknown. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2–4% and 15–20%, respectively.

#### Clinical Considerations

##### *Disease-Associated Maternal and/or Embryo/Fetal Risk*

Poorly controlled diabetes in pregnancy increases the maternal risk for diabetic ketoacidosis, pre-eclampsia, spontaneous abortions, preterm delivery and delivery

complications. Poorly controlled diabetes increases the fetal risk for major birth defects, stillbirth, and macrosomia-related morbidity.

## Data

### *Animal Data*

In pregnant rats given twice weekly subcutaneous doses of 0.02, 0.1, and 0.5 mg/kg tirzepatide (0.03-, 0.07-, and 0.5-fold the MRHD of 15 mg once weekly based on AUC) during organogenesis, increased incidences of external, visceral, and skeletal malformations, increased incidences of visceral and skeletal developmental variations, and decreased fetal weights coincided with pharmacologically-mediated reductions in maternal body weights and food consumption at 0.5 mg/kg. In pregnant rabbits given once weekly subcutaneous doses of 0.01, 0.03, or 0.1 mg/kg tirzepatide (0.01-, 0.06-, and 0.2-fold the MRHD) during organogenesis, pharmacologically-mediated effects on the gastrointestinal system resulting in maternal mortality or abortion in a few rabbits occurred at all dose levels. Reduced fetal weights associated with decreased maternal food consumption and body weights were observed at 0.1 mg/kg. In a pre- and post-natal study in rats administered subcutaneous doses of 0.02, 0.10, or 0.25 mg/kg tirzepatide twice weekly from implantation through lactation, F<sub>1</sub> pups from F<sub>0</sub> maternal rats given 0.25 mg/kg tirzepatide had statistically significant lower mean body weight when compared to controls from post-natal day 7 through post-natal day 126 for males and post-natal day 56 for females.

## **8.2 Lactation**

### Risk Summary

There are no data on the presence of tirzepatide in animal or human milk, the effects on the breastfed infant, or the effects on milk production. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for MOUNJARO and any potential adverse effects on the breastfed infant from MOUNJARO or from the underlying maternal condition.

## **8.3 Females and Males of Reproductive Potential**

### Contraception

Use of MOUNJARO may reduce the efficacy of oral hormonal contraceptives due to delayed gastric emptying. This delay is largest after the first dose and diminishes over time. Advise patients using oral hormonal contraceptives to switch to a non-oral contraceptive method or add a barrier method of contraception for 4 weeks after initiation and for 4 weeks after each dose escalation with MOUNJARO [see *Drug Interactions* (7.2) and *Clinical Pharmacology* (12.2, 12.3)].

## **8.4 Pediatric Use**

Safety and effectiveness of MOUNJARO have not been established in pediatric patients.

## **8.5 Geriatric Use**

In the pool of seven clinical trials, 1539 (30.1%) MOUNJARO-treated patients were 65 years of age or older, and 212 (4.1%) MOUNJARO-treated patients were 75 years of age or older at baseline.

No overall differences in safety or efficacy were detected between these patients and younger patients, but greater sensitivity of some older individuals cannot be ruled out.

## 8.6 Renal Impairment

No dosage adjustment of MOUNJARO is recommended for patients with renal impairment. In subjects with renal impairment including end-stage renal disease (ESRD), no change in tirzepatide pharmacokinetics (PK) was observed [see *Clinical Pharmacology* (12.3)]. Monitor renal function when initiating or escalating doses of MOUNJARO in patients with renal impairment reporting severe adverse gastrointestinal reactions [see *Warnings and Precautions* (5.5)].

## 8.7 Hepatic Impairment

No dosage adjustment of MOUNJARO is recommended for patients with hepatic impairment. In a clinical pharmacology study in subjects with varying degrees of hepatic impairment, no change in tirzepatide PK was observed [see *Clinical Pharmacology* (12.3)].


## 10 OVERDOSAGE

In the event of an overdose, consider contacting the Poison Help line (1-800-222-1222) or a medical toxicologist for additional overdose management recommendations. Initiate appropriate supportive treatment according to the patient's clinical signs and symptoms. A period of observation and treatment for these symptoms may be necessary, taking into account the half-life of tirzepatide of approximately 5 days.

## 11 DESCRIPTION

MOUNJARO (tirzepatide) injection, for subcutaneous use, contains tirzepatide, a once weekly GIP receptor and GLP-1 receptor agonist. Tirzepatide is based on the GIP sequence and contains aminoisobutyric acid (Aib) in positions 2 and 13, a C-terminal amide, and Lys residue at position 20 that is attached to 1,20-eicosanedioic acid via a linker. The molecular weight is 4813.53 Da and the empirical formula is  $C_{225}H_{348}N_{48}O_{68}$ .

Structural formula:



MOUNJARO is a clear, colorless to slightly yellow, sterile, preservative-free solution for subcutaneous use. Each single-dose pen or single-dose vial contains a 0.5 mL solution of 2.5 mg, 5 mg, 7.5 mg, 10 mg, 12.5 mg, or 15 mg of tirzepatide and the following

excipients: sodium chloride (4.1 mg), sodium phosphate dibasic heptahydrate (0.7 mg), and water for injection. Hydrochloric acid solution and/or sodium hydroxide solution may have been added to adjust the pH. MOUNJARO has a pH of 6.5 – 7.5.

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

Tirzepatide is a GIP receptor and GLP-1 receptor agonist. It is an amino-acid sequence including a C20 fatty diacid that enables albumin binding and prolongs the half-life. Tirzepatide selectively binds to and activates both the GIP and GLP-1 receptors, the targets for native GIP and GLP-1.

Tirzepatide enhances first- and second-phase insulin secretion, and reduces glucagon levels, both in a glucose-dependent manner.


### 12.2 Pharmacodynamics

Tirzepatide lowers fasting and postprandial glucose concentration, decreases food intake, and reduces body weight in patients with type 2 diabetes mellitus.

#### *First and Second-Phase Insulin Secretion*

Tirzepatide enhances the first- and second-phase insulin secretion. (Figure 1)

**Figure 1: Mean insulin concentration at 0-120 minutes during hyperglycemic clamp at baseline and Week 28**



### *Insulin Sensitivity*

Tirzepatide increases insulin sensitivity, as demonstrated in a hyperinsulinemic euglycemic clamp study after 28 weeks of treatment.

### *Glucagon Secretion*

Tirzepatide reduces fasting and postprandial glucagon concentrations. Tirzepatide 15 mg reduced fasting glucagon concentration by 28% and glucagon AUC after a mixed meal by 43%, compared with no change for placebo after 28 weeks of treatment.

### *Gastric Emptying*

Tirzepatide delays gastric emptying. The delay is largest after the first dose and this effect diminishes over time. Tirzepatide slows post-meal glucose absorption, reducing postprandial glucose.

## **12.3 Pharmacokinetics**

The pharmacokinetics of tirzepatide is similar between healthy subjects and patients with type 2 diabetes mellitus. Steady-state plasma tirzepatide concentrations were achieved following 4 weeks of once weekly administration. Tirzepatide exposure increases in a dose-proportional manner.

### Absorption

Following subcutaneous administration, the time to maximum plasma concentration of tirzepatide ranges from 8 to 72 hours. The mean absolute bioavailability of tirzepatide following subcutaneous administration is 80%. Similar exposure was achieved with subcutaneous administration of tirzepatide in the abdomen, thigh, or upper arm.

### Distribution

The mean apparent steady-state volume of distribution of tirzepatide following subcutaneous administration in patients with type 2 diabetes mellitus is approximately 10.3 L. Tirzepatide is highly bound to plasma albumin (99%).

### Elimination

The apparent population mean clearance of tirzepatide is 0.061 L/h with an elimination half-life of approximately 5 days, enabling once-weekly dosing.

### *Metabolism*

Tirzepatide is metabolized by proteolytic cleavage of the peptide backbone, beta-oxidation of the C20 fatty diacid and amide hydrolysis.

### *Excretion*

The primary excretion routes of tirzepatide metabolites are via urine and feces. Intact tirzepatide is not observed in urine or feces.

### Specific Populations

The intrinsic factors of age, gender, race, ethnicity, or body weight do not have a clinically relevant effect on the PK of tirzepatide.

### *Patients with Renal Impairment*



Renal impairment does not impact the pharmacokinetics of tirzepatide. The pharmacokinetics of tirzepatide after a single 5 mg dose was evaluated in patients with different degrees of renal impairment (mild, moderate, severe, ESRD) compared with subjects with normal renal function. This was also shown for patients with both type 2 diabetes mellitus and renal impairment based on data from clinical studies [see *Use in Specific Populations* (8.6)].

#### *Patients with Hepatic Impairment*

Hepatic impairment does not impact the pharmacokinetics of tirzepatide. The pharmacokinetics of tirzepatide after a single 5 mg dose was evaluated in patients with different degrees of hepatic impairment (mild, moderate, severe) compared with subjects with normal hepatic function [see *Use in Specific Populations* (8.7)].

#### Drug Interactions Studies

##### *Potential for Tirzepatide to Influence the Pharmacokinetics of Other Drugs*

In vitro studies have shown low potential for tirzepatide to inhibit or induce CYP enzymes, and to inhibit drug transporters.

MOUNJARO delays gastric emptying and thereby has the potential to impact the absorption of concomitantly administered oral medications [see *Drug Interactions* (7.2)].

The impact of tirzepatide on gastric emptying was greatest after a single dose of 5 mg and diminished after subsequent doses.

Following a first dose of tirzepatide 5 mg, acetaminophen maximum concentration ( $C_{max}$ ) was reduced by 50%, and the median peak plasma concentration ( $t_{max}$ ) occurred 1 hour later. After coadministration at week 4, there was no meaningful impact on acetaminophen  $C_{max}$  and  $t_{max}$ . Overall acetaminophen exposure ( $AUC_{0-24hr}$ ) was not influenced.

Following administration of a combined oral contraceptive (0.035 mg ethinyl estradiol and 0.25 mg norgestimate) in the presence of a single dose of tirzepatide 5 mg, mean  $C_{max}$  of ethinyl estradiol, norgestimate, and norelgestromin was reduced by 59%, 66%, and 55%, while mean AUC was reduced by 20%, 21%, and 23%, respectively. A delay in  $t_{max}$  of 2.5 to 4.5 hours was observed.

## **12.6 Immunogenicity**

The observed incidence of anti-drug antibodies is highly dependent on the sensitivity and specificity of the assay. Differences in assay methods preclude meaningful comparisons of the incidence of anti-drug antibodies in the trials described below with the incidence of anti-drug antibodies in other trials.

During the 40- to 104-week treatment periods with ADA sampling conducted up to 44 to 108 weeks in seven clinical trials in adults with type 2 diabetes mellitus [see *Clinical Studies* (14)], 51% (2,570/5,025) of MOUNJARO-treated patients developed anti-tirzepatide antibodies. In these trials, anti-tirzepatide antibody formation in 34% and 14% of MOUNJARO-treated patients showed cross-reactivity to native GIP or native GLP-1, respectively.

Of the 2,570 MOUNJARO-treated patients who developed anti-tirzepatide antibodies during the treatment periods in these seven trials, 2% and 2% developed neutralizing antibodies against tirzepatide activity on the GIP or GLP-1 receptors, respectively, and

0.9% and 0.4% developed neutralizing antibodies against native GIP or GLP-1, respectively.

There was no identified clinically significant effect of anti-tirzepatide antibodies on pharmacokinetics or effectiveness of MOUNJARO. More MOUNJARO-treated patients who developed anti-tirzepatide antibodies experienced hypersensitivity reactions or injection site reactions than those who did not develop these antibodies [see *Adverse Reactions (6.1)*].

## **13 NONCLINICAL TOXICOLOGY**

### **13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility**

A 2-year carcinogenicity study was conducted with tirzepatide in male and female rats at doses of 0.15, 0.50, and 1.5 mg/kg (0.1-, 0.4-, and 1-fold the MRHD of 15 mg once weekly based on AUC) administered by subcutaneous injection twice weekly. A statistically significant increase in thyroid C-cell adenomas was observed in males ( $\geq 0.5$  mg/kg) and females ( $\geq 0.15$  mg/kg), and a statistically significant increase in thyroid C-cell adenomas and carcinomas combined was observed in males and females at all doses examined. In a 6-month carcinogenicity study in rasH2 transgenic mice, tirzepatide at doses of 1, 3, and 10 mg/kg administered by subcutaneous injection twice weekly was not tumorigenic.

Tirzepatide was not genotoxic in a rat bone marrow micronucleus assay.

In fertility and early embryonic development studies, male and female rats were administered twice weekly subcutaneous doses of 0.5, 1.5, or 3 mg/kg (0.3-, 1-, and 2-fold and 0.3-, 0.9-, and 2-fold, respectively, the MRHD of 15 mg once weekly based on AUC). No effects of tirzepatide were observed on sperm morphology, mating, fertility, and conception. In female rats, an increase in the number of females with prolonged diestrus and a decrease in the mean number of corpora lutea resulting in a decrease in the mean number of implantation sites and viable embryos was observed at all dose levels. These effects were considered secondary to the pharmacological effects of tirzepatide on food consumption and body weight.

## **14 CLINICAL STUDIES**

### **14.1 Overview of Clinical Studies**

The effectiveness of MOUNJARO as an adjunct to diet and exercise to improve glycemic control in adults with type 2 diabetes mellitus was established in five trials. In these trials, MOUNJARO was studied as monotherapy (SURPASS-1); as an add-on to metformin, sulfonylureas, and/or sodium-glucose co-transporter 2 inhibitors (SGLT2 inhibitors) (SURPASS-2, -3, and -4); and in combination with basal insulin with or without metformin (SURPASS-5). In these trials, MOUNJARO (5 mg, 10 mg, and 15 mg given subcutaneously once weekly) was compared with placebo, semaglutide 1 mg, insulin degludec, and/or insulin glargine.

In adult patients with type 2 diabetes mellitus, treatment with MOUNJARO produced a statistically significant reduction from baseline in HbA<sub>1c</sub> compared to placebo. The effectiveness of MOUNJARO was not impacted by age, gender, race, ethnicity, region, or

by baseline BMI, HbA1c, diabetes duration, or renal function.

## 14.2 Monotherapy Use of MOUNJARO in Adult Patients with Type 2 Diabetes Mellitus

SURPASS-1 (NCT03954834) was a 40-week double-blind trial that randomized 478 adult patients with type 2 diabetes mellitus with inadequate glycemic control with diet and exercise to MOUNJARO 5 mg, MOUNJARO 10 mg, MOUNJARO 15 mg, or placebo once weekly.

Patients had a mean age of 54 years, and 52% were men. The mean duration of type 2 diabetes mellitus was 4.7 years, and the mean BMI was 32 kg/m<sup>2</sup>. Overall, 36% were White, 35% were Asian, 25% were American Indians/Alaska Natives, and 5% were Black or African American; 43% identified as Hispanic or Latino ethnicity.

Monotherapy with MOUNJARO 5 mg, 10 mg and 15 mg once weekly for 40 weeks resulted in a statistically significant reduction in HbA1c compared with placebo (see Table 3).

**Table 3: Results at Week 40 in a Trial of MOUNJARO as Monotherapy in Adult Patients with Type 2 Diabetes Mellitus with Inadequate Glycemic Control with Diet and Exercise**

	Placebo	MOUNJARO 5 mg	MOUNJARO 10 mg	MOUNJARO 15 mg
Modified Intent-to-Treat (mITT) Population (N) <sup>a</sup>	113	121	121	120
HbA1c (%)				
Baseline (mean)	8.1	8.0	7.9	7.9
Change at Week 40 <sup>b</sup>	-0.1	-1.8	-1.7	-1.7
Difference from placebo <sup>b</sup> (95% CI)	--	-1.7 <sup>c</sup> (-2.0, -1.4)	-1.6 <sup>c</sup> (-1.9, -1.3)	-1.6 <sup>c</sup> (-1.9, -1.3)
Patients (%) achieving HbA1c <7% <sup>d</sup>	23	82 <sup>c</sup>	85 <sup>c</sup>	78 <sup>c</sup>
Fasting Serum Glucose (mg/dL)				
Baseline (mean)	155	154	153	154
Change at Week 40 <sup>b</sup>	4	-40	-40	-39
Difference from placebo <sup>b</sup> (95% CI)	--	-43 <sup>c</sup> (-55, -32)	-43 <sup>c</sup> (-55, -32)	-42 <sup>c</sup> (-54, -30)
Body Weight (kg)				
Baseline (mean)	84.5	87.0	86.2	85.5
Change at Week 40 <sup>b</sup>	-1.0	-6.3	-7.0	-7.8
Difference from placebo <sup>b</sup> (95% CI)	--	-5.3 <sup>c</sup> (-6.8, -3.9)	-6.0 <sup>c</sup> (-7.4, -4.6)	-6.8 <sup>c</sup> (-8.3, -5.4)

<sup>a</sup> The modified intent-to-treat population consists of all randomly assigned participants who were exposed to at least 1 dose of study drug. Patients who discontinued study treatment because they did not meet study enrollment criteria were excluded. During the trial, rescue medication (additional antihyperglycemic medication) was initiated by 25%, 2%, 3%, and 2% of patients randomized to placebo, MOUNJARO 5 mg, 10 mg, and 15 mg, respectively. At Week 40 the HbA1c data were missing for 12%, 6%, 7%, and 14% of patients randomized to placebo, MOUNJARO 5

mg, 10 mg, and 15 mg, respectively. Missing Week 40 data were imputed using placebo-based multiple imputation.

<sup>b</sup> Least-squares mean from ANCOVA adjusted for baseline value and other stratification factors.

<sup>c</sup>  $p < 0.001$  (two-sided) for superiority vs. placebo, adjusted for multiplicity.

<sup>d</sup> Analyzed using logistic regression adjusted for baseline value and other stratification factors.

### 14.3 MOUNJARO Use in Combination with Metformin, Sulfonylureas, and/or SGLT2 Inhibitors in Adult Patients with Type 2 Diabetes Mellitus

#### Add-on to metformin

SURPASS-2 (NCT03987919) was a 40-week open-label trial (double-blind with respect to MOUNJARO dose assignment) that randomized 1879 adult patients with type 2 diabetes mellitus with inadequate glycemic control on stable doses of metformin alone to the addition of MOUNJARO 5 mg, MOUNJARO 10 mg, or MOUNJARO 15 mg once weekly or subcutaneous semaglutide 1 mg once weekly.

Patients had a mean age of 57 years and 47% were men. The mean duration of type 2 diabetes mellitus was 8.6 years, and the mean BMI was 34 kg/m<sup>2</sup>. Overall, 83% were White, 4% were Black or African American, and 1% were Asian; 70% identified as Hispanic or Latino ethnicity.

Treatment with MOUNJARO 10 mg and 15 mg once weekly for 40 weeks resulted in a statistically significant reduction in HbA1c compared with semaglutide 1 mg once weekly (see Table 4 and Figure 2).

**Table 4: Results at Week 40 in a Trial of MOUNJARO versus Semaglutide 1 mg in Adult Patients with Type 2 Diabetes Mellitus Added to Metformin**

	<b>Semaglutide 1 mg</b>	<b>MOUNJARO 5 mg</b>	<b>MOUNJARO 10 mg</b>	<b>MOUNJARO 15 mg</b>
Modified Intent-to-Treat (mITT) Population (N) <sup>a</sup>	468	470	469	469
HbA1c (%)				
Baseline (mean)	8.3	8.3	8.3	8.3
Change at Week 40 <sup>b</sup>	-1.9	-2.0	-2.2	-2.3
Difference from semaglutide <sup>b</sup> (95% CI)	--	-0.2 <sup>c</sup> (-0.3, -0.0)	-0.4 <sup>d</sup> (-0.5, -0.3)	-0.5 <sup>d</sup> (-0.6, -0.3)
Patients (%) achieving HbA1c <7% <sup>e</sup>	79	82	86 <sup>f</sup>	86 <sup>f</sup>
Fasting Serum Glucose (mg/dL)				
Baseline (mean)	171	174	174	172
Change at Week 40 <sup>b</sup>	-49	-55	-59	-60
Body Weight (kg)				
Baseline (mean)	93.7	92.5	94.8	93.8
Change at Week 40 <sup>b</sup>	-5.7	-7.6	-9.3	-11.2
Difference from semaglutide <sup>b</sup> (95% CI)	--	-1.9 <sup>c</sup> (-2.8, -1.0)	-3.6 <sup>d</sup> (-4.5, -2.7)	-5.5 <sup>d</sup> (-6.4, -4.6)

<sup>a</sup> The modified intent-to-treat population consists of all randomly assigned participants who were exposed to at least 1 dose of study drug. Patients who discontinued study treatment because they did not meet study enrollment criteria were excluded. During the trial, rescue medication

(additional antihyperglycemic medication) was initiated by 3%, 2%, 1%, and 1% of patients randomized to semaglutide 1 mg, MOUNJARO 5 mg, 10 mg, and 15 mg, respectively. At Week 40 the HbA1c endpoint was missing for 5%, 4%, 5%, and 5% of patients randomized to semaglutide 1 mg, MOUNJARO 5 mg, 10 mg, and 15 mg, respectively. Missing Week 40 data were imputed using multiple imputation with retrieved dropout.

<sup>b</sup> Least-squares mean from ANCOVA adjusted for baseline value and other stratification factors.


<sup>c</sup>  $p < 0.05$  (two-sided) for superiority vs. semaglutide, adjusted for multiplicity.

<sup>d</sup>  $p < 0.001$  (two-sided) for superiority vs. semaglutide, adjusted for multiplicity.

<sup>e</sup> Analyzed using logistic regression adjusted for baseline value and other stratification factors.

<sup>f</sup>  $p < 0.01$  (two-sided) for superiority vs. semaglutide, adjusted for multiplicity.

**Figure 2: Mean HbA1c (%) Over Time - Baseline to Week 40**



#### Number of patients

<b>MOUNJARO 5mg</b>	470	451	470
<b>MOUNJARO 10mg</b>	469	445	469
<b>MOUNJARO 15mg</b>	469	447	469
<b>Semaglutide 1mg</b>	468	443	468

Note: Displayed results are from modified Intent-to-Treat Full Analysis Set. (1) Observed mean value from Week 0 to Week 40, and (2) least-squares mean  $\pm$  standard error at Week 40 multiple imputation (MI).

#### Add-on to metformin with or without SGLT2 inhibitor

SURPASS-3 (NCT03882970) was a 52-week open-label trial that randomized 1444 adult patients with type 2 diabetes mellitus with inadequate glycemic control on stable doses of metformin with or without SGLT2 inhibitor to the addition of MOUNJARO 5 mg,

MOUNJARO 10 mg, MOUNJARO 15 mg once weekly, or insulin degludec 100 units/mL once daily. In this trial, 32% of patients were on SGLT2 inhibitor. Insulin degludec was initiated at 10 units once daily and adjusted weekly throughout the trial using a treat-to-target algorithm based on self-measured fasting blood glucose values. At Week 52, 26% of patients randomized to insulin degludec achieved the fasting serum glucose target of <90 mg/dL, and the mean daily insulin degludec dose was 49 U (0.5 U per kilogram).

Patients had a mean age of 57 years, and 56% were men. The mean duration of type 2 diabetes mellitus was 8.4 years, and the mean baseline BMI was 34 kg/m<sup>2</sup>. Overall, 91% were White, 3% were Black or African American, and 5% were Asian; 29% identified as Hispanic or Latino ethnicity.

Treatment with MOUNJARO 10 mg and 15 mg once weekly for 52 weeks resulted in a statistically significant reduction in HbA1c compared with daily insulin degludec (see Table 5).

**Table 5: Results at Week 52 in a Trial of MOUNJARO versus Insulin Degludec in Adult Patients with Type 2 Diabetes Mellitus Added to Metformin with or without SGLT2 Inhibitor**

	<b>Insulin Degludec</b>	<b>MOUNJARO 5 mg</b>	<b>MOUNJARO 10 mg</b>	<b>MOUNJARO 15 mg</b>
Modified Intent-to-Treat (mITT) <sup>a</sup> Population (N)	359	358	360	358
HbA1c (%)				
Baseline (mean)	8.1	8.2	8.2	8.2
Change at Week 52 <sup>b</sup>	-1.3	-1.9	-2.0	-2.1
Difference from insulin degludec <sup>b</sup> (95% CI)	--	-0.6 <sup>c</sup> (-0.7, -0.5)	-0.8 <sup>c</sup> (-0.9, -0.6)	-0.9 <sup>c</sup> (-1.0, -0.7)
Patients (%) achieving HbA1c <7% <sup>d</sup>	58	79 <sup>c</sup>	82 <sup>c</sup>	83 <sup>c</sup>
Fasting Serum Glucose (mg/dL)				
Baseline (mean)	167	172	170	168
Change at Week 52 <sup>b</sup>	-51	-47	-50	-54
Body Weight (kg)				
Baseline (mean)	94.0	94.4	93.8	94.9
Change at Week 52 <sup>b</sup>	1.9	-7.0	-9.6	-11.3
Difference from insulin degludec <sup>b</sup> (95% CI)	--	-8.9 <sup>c</sup> (-10.0, -7.8)	-11.5 <sup>c</sup> (-12.6, -10.4)	-13.2 <sup>c</sup> (-14.3, -12.1)

<sup>a</sup> The modified intent-to-treat population consists of all randomly assigned participants who were exposed to at least 1 dose of study drug. Patients who discontinued study treatment because they did not meet study enrollment criteria were excluded. During the trial, rescue medication (additional antihyperglycemic medication) was initiated by 1%, 1%, 1%, and 2% of patients randomized to insulin degludec, MOUNJARO 5 mg, 10 mg, and 15 mg, respectively. At Week 52 the HbA1c endpoint was missing for 9%, 6%, 10%, and 5% of patients randomized to insulin degludec, MOUNJARO 5 mg, 10 mg, and 15 mg, respectively. Missing Week 52 data were imputed using multiple imputation with retrieved dropout.

<sup>b</sup> Least-squares mean from ANCOVA adjusted for baseline value and other stratification factors.

<sup>c</sup> p<0.001 (two-sided) for superiority vs. insulin degludec, adjusted for multiplicity.

<sup>d</sup> Analyzed using logistic regression adjusted for baseline value and other stratification factors.

Add-on to 1-3 oral anti-hyperglycemic agents (metformin, sulfonylurea, or SGLT-2 inhibitor)

SURPASS-4 (NCT03730662) was a 104-week open-label trial (52-week primary endpoint) that randomized 2002 adult patients with type 2 diabetes mellitus with increased cardiovascular risk to MOUNJARO 5 mg, MOUNJARO 10 mg, MOUNJARO 15 mg once weekly, or insulin glargine 100 units/mL once daily (1:1:1:3 ratio) on a background of metformin (95%) and/or sulfonylureas (54%) and/or SGLT2 inhibitors (25%).

Patients had a mean age of 64 years, and 63% were men. The mean duration of type 2 diabetes mellitus was 11.8 years, and the mean baseline BMI was 33 kg/m<sup>2</sup>. Overall, 82% were White, 4% were Black or African American, and 4% were Asian; 48% identified as Hispanic or Latino ethnicity. Across all treatment groups, 87% had a history of cardiovascular disease. At baseline, eGFR was  $\geq 90$  mL/min/1.73 m<sup>2</sup> in 43%, 60 to 90 mL/min/1.73 m<sup>2</sup> in 40%, 45 to 60 mL/min/1.73 m<sup>2</sup> in 10%, and 30 to 45 mL/min/1.73 m<sup>2</sup> in 6% of patients.

Insulin glargine was initiated at 10 U once daily and adjusted weekly throughout the trial using a treat-to-target algorithm based on self-measured fasting blood glucose values. At Week 52, 30% of patients randomized to insulin glargine achieved the fasting serum glucose target of <100 mg/dL, and the mean daily insulin glargine dose was 44 U (0.5 U per kilogram).

Treatment with MOUNJARO 10 mg and 15 mg once weekly for 52 weeks resulted in a statistically significant reduction in HbA1c compared with insulin glargine once daily (see Table 6).

**Table 6: Results at Week 52 in a Trial of MOUNJARO versus Insulin Glargine in Adult Patients with Type 2 Diabetes Mellitus Added to Metformin and/or Sulfonylurea and/or SGLT2 Inhibitor**

	<b>Insulin Glargine</b>	<b>MOUNJARO 5 mg</b>	<b>MOUNJARO 10 mg</b>	<b>MOUNJARO 15 mg</b>
Modified Intent-to-Treat (mITT) Population (N) <sup>a</sup>	998	328	326	337
HbA1c (%)				
Baseline (mean)	8.5	8.5	8.6	8.5
Change at Week 52 <sup>b</sup>	-1.4	-2.1	-2.3	-2.4
Difference from insulin glargine <sup>b</sup> (95% CI)	--	-0.7 <sup>c</sup> (-0.9, -0.6)	-0.9 <sup>c</sup> (-1.1, -0.8)	-1.0 <sup>c</sup> (-1.2, -0.9)
Patients (%) achieving HbA1c <7% <sup>d</sup>	49	75 <sup>c</sup>	83 <sup>c</sup>	85 <sup>c</sup>
Fasting Serum Glucose (mg/dL)				
Baseline (mean)	168	172	176	174
Change at Week 52 <sup>b</sup>	-49	-44	-50	-55
Body Weight (kg)				
Baseline (mean)	90.2	90.3	90.6	90.0
Change at Week 52 <sup>b</sup>	1.7	-6.4	-8.9	-10.6
Difference from insulin glargine <sup>b</sup> (95% CI)	--	-8.1 <sup>c</sup> (-8.9, -7.3)	-10.6 <sup>c</sup> (-11.4, -9.8)	-12.2 <sup>c</sup> (-13.0, -11.5)

- <sup>a</sup> The modified intent-to-treat population consists of all randomly assigned participants who were exposed to at least 1 dose of study drug. Patients who discontinued study treatment because they did not meet study enrollment criteria were excluded. During the trial, rescue medication (additional antihyperglycemic medication) was initiated by 1%, 0%, 0%, and 1% of patients randomized to insulin glargine, MOUNJARO 5 mg, 10 mg, and 15 mg, respectively. At Week 52 the HbA1c endpoint was missing for 9%, 9%, 6%, and 4% of patients randomized to insulin glargine, MOUNJARO 5 mg, 10 mg, and 15 mg, respectively. Missing Week 52 data were imputed using multiple imputation with retrieved dropout.
- <sup>b</sup> Least-squares mean from ANCOVA adjusted for baseline value and other stratification factors.
- <sup>c</sup>  $p < 0.001$  (two-sided) for superiority vs. insulin glargine, adjusted for multiplicity.
- <sup>d</sup> Analyzed using logistic regression adjusted for baseline value and other stratification factors.

#### 14.4 MOUNJARO Use in Combination with Basal Insulin with or without Metformin in Adult Patients with Type 2 Diabetes Mellitus

SURPASS-5 (NCT04039503) was a 40-week double-blind trial that randomized 475 patients with type 2 diabetes mellitus with inadequate glycemic control on insulin glargine 100 units/mL, with or without metformin, to MOUNJARO 5 mg, MOUNJARO 10 mg, MOUNJARO 15 mg once weekly, or placebo. The dose of background insulin glargine was adjusted using a treat-to-target algorithm based on self-measured fasting blood glucose values, targeting  $<100$  mg/dL.

Patients had a mean age of 61 years, and 56% were men. The mean duration of type 2 diabetes mellitus was 13.3 years, and the mean baseline BMI was  $33 \text{ kg/m}^2$ . Overall, 80% were White, 1% were Black or African American, and 18% were Asian; 5% identified as Hispanic or Latino ethnicity.

The mean dose of insulin glargine at baseline was 34, 32, 35, and 33 units/day for patients receiving MOUNJARO 5 mg, 10 mg, 15 mg, and placebo, respectively. At randomization, the initial insulin glargine dose in patients with  $\text{HbA1c} \leq 8.0\%$  was reduced by 20%. At week 40, mean dose of insulin glargine was 38, 36, 29, and 59 units/day for patients receiving MOUNJARO 5 mg, 10 mg, 15 mg, and placebo, respectively.

Treatment with MOUNJARO 5 mg once weekly, 10 mg once weekly and 15 mg once weekly for 40 weeks resulted in a statistically significant reduction in HbA1c compared with placebo (see Table 7).

**Table 7: Results at Week 40 in a Trial of MOUNJARO Added to Basal Insulin with or without Metformin in Adult Patients with Type 2 Diabetes Mellitus**

	Placebo	MOUNJARO 5 mg	MOUNJARO 10 mg	MOUNJARO 15 mg
Modified Intent-to-Treat (mITT) Population (N) <sup>a</sup>	119	116	118	118
HbA1c (%)				
Baseline (mean)	8.4	8.3	8.4	8.2
Change at Week 40 <sup>b</sup>	-0.9	-2.1	-2.4	-2.3
Difference from placebo <sup>b</sup> (95% CI)	--	-1.2 <sup>c</sup> (-1.5, -1.0)	-1.5 <sup>c</sup> (-1.8, -1.3)	-1.5 <sup>c</sup> (-1.7, -1.2)
Patients (%) achieving HbA1c $<7\%$ <sup>d</sup>	35	87 <sup>c</sup>	90 <sup>c</sup>	85 <sup>c</sup>
Fasting Serum Glucose (mg/dL)				



Baseline (mean)	164	163	163	160
Change at Week 40 <sup>b</sup>	-39	-58	-64	-63
Difference from placebo <sup>b</sup> (95% CI)	--	-19 <sup>c</sup> (-27, -11)	-25 <sup>c</sup> (-32, -17)	-23 <sup>c</sup> (-31, -16)
Body Weight (kg)				
Baseline (mean)	94.2	95.8	94.6	96.0
Change at Week 40 <sup>b</sup>	1.6	-5.4	-7.5	-8.8
Difference from placebo <sup>b</sup> (95% CI)	--	-7.1 <sup>c</sup> (-8.7, -5.4)	-9.1 <sup>c</sup> (-10.7, -7.5)	-10.5 <sup>c</sup> (-12.1, -8.8)

<sup>a</sup> The modified intent-to-treat population consists of all randomly assigned participants who were exposed to at least 1 dose of study drug. Patients who discontinued study treatment because they did not meet study enrollment criteria were excluded. During the trial, rescue medication (additional antihyperglycemic medication) was initiated by 4%, 1%, 0%, and 1% of patients randomized to placebo, MOUNJARO 5 mg, 10 mg, and 15 mg, respectively. At Week 40 the HbA1c endpoint was missing for 2%, 6%, 3%, and 7% of patients randomized to placebo, MOUNJARO 5 mg, 10 mg, and 15 mg, respectively. Missing Week 40 data were imputed using placebo-based multiple imputation.

<sup>b</sup> Least-squares mean from ANCOVA adjusted for baseline value and other stratification factors.

<sup>c</sup> p<0.001 (two-sided) for superiority vs. placebo, adjusted for multiplicity.

<sup>d</sup> Analyzed using logistic regression adjusted for baseline value and other stratification factors.

## 16 HOW SUPPLIED/STORAGE AND HANDLING

### 16.1 How Supplied

MOUNJARO is a clear, colorless to slightly yellow solution available in cartons containing 4 pre-filled single-dose pens or 1 single-dose vial as follows:

<b>Total Strength per Total Volume</b>	<b>Pen NDC</b>	<b>Vial NDC</b>
2.5 mg/0.5 mL	0002-1506-80	0002-1152-01
5 mg/0.5 mL	0002-1495-80	0002-1243-01
7.5 mg/0.5 mL	0002-1484-80	0002-2214-01
10 mg/0.5 mL	0002-1471-80	0002-2340-01
12.5 mg/0.5 mL	0002-1460-80	0002-2423-01
15 mg/0.5 mL	0002-1457-80	0002-3002-01

### 16.2 Storage and Handling

- Store MOUNJARO in a refrigerator at 2°C to 8°C (36°F to 46°F).
- If needed, each single-dose pen or single-dose vial can be stored unrefrigerated at temperatures not to exceed 30°C (86°F) for up to 21 days.
- Do not freeze MOUNJARO. Do not use MOUNJARO if frozen.
- Store MOUNJARO in the original carton to protect from light.

## 17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (*Medication Guide and Instructions for Use*).

### Risk of Thyroid C-Cell Tumors

Inform patients that MOUNJARO causes thyroid C-cell tumors in rats and that the human relevance of this finding has not been determined. Counsel patients to report symptoms of thyroid tumors (e.g., a lump in the neck, persistent hoarseness, dysphagia, or dyspnea) to their healthcare provider *[see Boxed Warning and Warnings and Precautions (5.1)]*.

### Acute Pancreatitis

Inform patients of the potential risk for acute pancreatitis and its symptoms: severe abdominal pain that may radiate to the back, and which may or may not be accompanied by vomiting. Instruct patients to discontinue MOUNJARO promptly and contact their physician if pancreatitis is suspected *[see Warnings and Precautions (5.2)]*.

### Hypoglycemia with Concomitant Use of Insulin Secretagogues or Insulin

Inform patients that the risk of hypoglycemia is increased when MOUNJARO is used with an insulin secretagogue (such as a sulfonylurea) or insulin. Educate patients on the signs and symptoms of hypoglycemia *[see Warnings and Precautions (5.3)]*.

### Hypersensitivity Reactions

Inform patients that serious hypersensitivity reactions have been reported with use of MOUNJARO. Advise patients on the symptoms of hypersensitivity reactions and instruct them to stop taking MOUNJARO and seek medical advice promptly if such symptoms occur *[see Warnings and Precautions (5.4)]*.

### Acute Kidney Injury Due to Volume Depletion

Inform patients of the potential risk of acute kidney injury due to dehydration associated with gastrointestinal adverse reactions. Advise patients to take precautions to avoid fluid depletion. Inform patients of the signs and symptoms of acute kidney injury and instruct them to promptly report any of these signs or symptoms or persistent (or extended) nausea, vomiting, and diarrhea to their healthcare provider *[see Warnings and Precautions (5.5)]*.

### Severe Gastrointestinal Adverse Reactions

Inform patients of the potential risk of severe gastrointestinal adverse reactions. Instruct patients to contact their healthcare provider if they have severe or persistent gastrointestinal symptoms *[see Warnings and Precautions (5.6)]*.

### Diabetic Retinopathy Complications in Patients with a History of Diabetic Retinopathy

Inform patients to contact their healthcare provider if changes in vision are experienced during treatment with MOUNJARO *[see Warnings and Precautions (5.7)]*.

### Acute Gallbladder Disease

Inform patients of the risk of acute gallbladder disease. Instruct patients to contact their healthcare provider for appropriate clinical follow-up if gallbladder disease is suspected *[see Warnings and Precautions (5.8)]*.

### Pulmonary Aspiration During General Anesthesia or Deep Sedation

Inform patients that MOUNJARO may cause their stomach to empty more slowly which may lead to complications with anesthesia or deep sedation during planned surgeries or

procedures. Instruct patients to inform healthcare providers prior to any planned surgeries or procedures if they are taking MOUNJARO [see *Warnings and Precautions* (5.9)].

### Pregnancy

Advise a pregnant woman of the potential risk to a fetus. Advise women to inform their healthcare provider if they are pregnant or intend to become pregnant [see *Use in Specific Populations* (8.1)].

### Contraception

Use of MOUNJARO may reduce the efficacy of oral hormonal contraceptives. Advise patients using oral hormonal contraceptives to switch to a non-oral contraceptive method or add a barrier method of contraception for 4 weeks after initiation and for 4 weeks after each dose escalation with MOUNJARO [see *Drug Interactions* (7.2), *Use in Specific Populations* (8.3), and *Clinical Pharmacology* (12.3)].

### Administration

Instruct patients how to prepare and administer the correct dose of MOUNJARO and assess their ability to inject subcutaneously to ensure the proper administration of MOUNJARO. Instruct patients using the single-dose vial to use a syringe appropriate for dose administration (e.g., a 1 mL syringe capable of measuring a 0.5 mL dose) [see *Dosage and Administration* (2.2)].

### Missed Doses

Inform patients if a dose is missed, it should be administered as soon as possible within 4 days after the missed dose. If more than 4 days have passed, the missed dose should be skipped and the next dose should be administered on the regularly scheduled day. In each case, inform patients to resume their regular once weekly dosing schedule [see *Dosage and Administration* (2.1)].

**Marketed by: Lilly USA, LLC, Indianapolis, IN 46285, USA**

Copyright © 2022, 2025, Eli Lilly and Company. All rights reserved.

Pat.: [www.lilly.com/patents](http://www.lilly.com/patents)

MOU-0006-USPI-20250619

## **Mounjaro Medication Guide**

<p style="text-align: center;"><b>Medication Guide</b> <b>MOUNJARO® [mown-JAHR-OH]</b> <b>(tirzepatide)</b> <b>injection, for subcutaneous use</b></p>
--

**What is the most important information I should know about MOUNJARO?**  
**MOUNJARO may cause serious side effects, including:**

- **Possible thyroid tumors, including cancer.** Tell your healthcare provider if you get a lump or swelling in your neck, hoarseness, trouble swallowing, or shortness of breath. These may be symptoms of thyroid cancer. In studies with rats, MOUNJARO and medicines that work like MOUNJARO caused thyroid tumors, including thyroid cancer. It is not known if MOUNJARO will cause thyroid tumors, or a type of thyroid

cancer called medullary thyroid carcinoma (MTC) in people.

- Do not use MOUNJARO if you or any of your family have ever had a type of thyroid cancer called medullary thyroid carcinoma (MTC), or if you have an endocrine system condition called Multiple Endocrine Neoplasia syndrome type 2 (MEN 2).

### **What is MOUNJARO?**

- MOUNJARO is an injectable prescription medicine that is used along with diet and exercise to improve blood sugar (glucose) in adults with type 2 diabetes mellitus.
- It is not known if MOUNJARO is safe and effective for use in children.

### **Do not use MOUNJARO if:**

- you or any of your family have ever had a type of thyroid cancer called MTC or if you have an endocrine system condition called MEN 2.
- you have had a serious allergic reaction to tirzepatide or any of the ingredients in MOUNJARO. See the end of this Medication Guide for a complete list of ingredients in MOUNJARO. See "**What are the possible side effects of MOUNJARO?**" for symptoms of a serious allergic reaction.

### **Before using MOUNJARO, tell your healthcare provider about all of your medical conditions, including if you:**

- have or have had problems with your pancreas.
- have severe problems with your stomach, such as slowed emptying of your stomach (gastroparesis) or problems with digesting food.
- have a history of diabetic retinopathy.
- are scheduled to have surgery or other procedures that use anesthesia or deep sleepiness (deep sedation).
- are pregnant or plan to become pregnant. It is not known if MOUNJARO will harm your unborn baby. Tell your healthcare provider if you become pregnant while using MOUNJARO.
  - **Birth control pills by mouth may not work as well while using MOUNJARO.** If you take birth control pills by mouth, your healthcare provider may recommend another type of birth control for 4 weeks after you start MOUNJARO and for 4 weeks after each increase in your dose of MOUNJARO. Talk to your healthcare provider about birth control methods that may be right for you while using MOUNJARO.
- are breastfeeding or plan to breastfeed. It is not known if MOUNJARO passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby while using MOUNJARO.

**Tell your healthcare provider about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. MOUNJARO may affect the way some medicines work, and some medicines may affect the way MOUNJARO works.

**Before using MOUNJARO, talk to your healthcare provider about low blood sugar and how to manage it.** Tell your healthcare provider if you are taking other medicines to treat diabetes including insulin or sulfonylureas. Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist when you get a new medicine.

### **How should I use MOUNJARO?**

- Read the **Instructions for Use** that comes with MOUNJARO.
- Use MOUNJARO exactly as your healthcare provider tells you to.
- **Your healthcare provider should show you how to prepare and inject your dose of MOUNJARO before injecting for the first time.**
- Inject MOUNJARO under the skin (subcutaneously) of your stomach (abdomen), thigh, or upper arm. **Do not** inject MOUNJARO into a muscle (intramuscularly) or vein (intravenously).
- **Use MOUNJARO 1 time each week, at any time of the day.**
- You may change the day of the week you use MOUNJARO as long as the time between the 2 doses is at least **3** days (72 hours).
- If you miss a dose of MOUNJARO, take the missed dose as soon as possible within 4 days (96 hours) after the missed dose. If more than 4 days have passed, skip the missed dose and take your next dose on the regularly scheduled day. **Do not** take **2** doses of MOUNJARO within **3** days of each other.
- MOUNJARO may be taken with or without food.
- **Do not** mix insulin and MOUNJARO together in the same injection.
- You may give an injection of MOUNJARO and insulin in the same body area (such as your stomach area), but not right next to each other.
- Change (rotate) your injection site with each weekly injection. **Do not** use the same site for each injection.
- If you take too much MOUNJARO, call your healthcare provider or Poison Help line at 1-800-222-1222 or go to the nearest hospital emergency room right away.

## **What are the possible side effects of MOUNJARO?**

### **MOUNJARO may cause serious side effects, including:**

- See **“What is the most important information I should know about MOUNJARO?”**
- **inflammation of your pancreas (pancreatitis).** Stop using MOUNJARO and call your healthcare provider right away if you have severe pain in your stomach area (abdomen) that will not go away, with or without vomiting. You may feel the pain from your abdomen to your back.
- **low blood sugar (hypoglycemia).** Your risk for getting low blood sugar may be higher if you use MOUNJARO with another medicine that can cause low blood sugar, such as a sulfonylurea or insulin.

### **Signs and symptoms of low blood sugar may include:**

- |                                 |                  |  |
|---------------------------------|------------------|--|
| ◦ dizziness or light-headedness | ◦ blurred vision | ◦ anxiety, irritability, or mood changes |
| ◦ sweating                      | ◦ slurred speech | ◦ hunger                                 |
| ◦ confusion or drowsiness       | ◦ shakiness      | ◦ weakness                               |
| ◦ headache                      | ◦ fast heartbeat | ◦ feeling jittery                        |
- **serious allergic reactions.** Stop using MOUNJARO and get medical help right away if you have any symptoms of a serious allergic reaction including:
 

◦ swelling of your face, lips, tongue, or throat	◦ fainting or feeling dizzy
◦ problems breathing or swallowing	◦ very rapid heartbeat
◦ severe rash or itching	

- **dehydration leading to kidney problems.** Diarrhea, nausea, and vomiting may cause a loss of fluids (dehydration) which may cause kidney problems. It is important for you to drink fluids to help reduce your chance of dehydration.
- **severe stomach problems.** Stomach problems, sometimes severe, have been reported in people who use MOUNJARO. Tell your healthcare provider if you have stomach problems that are severe or will not go away.
- **changes in vision.** Tell your healthcare provider if you have changes in vision during treatment with MOUNJARO.
- **gallbladder problems.** Gallbladder problems have happened in some people who use MOUNJARO. Tell your healthcare provider right away if you get symptoms of gallbladder problems which may include:
  - pain in your upper stomach (abdomen)
  - fever
  - yellowing of skin or eyes (jaundice)
  - clay-colored stools
- **food or liquid getting into the lungs during surgery or other procedures that use anesthesia or deep sleepiness (deep sedation).** MOUNJARO may increase the chance of food getting into your lungs during surgery or other procedures. Tell all your healthcare providers that you are taking MOUNJARO before you are scheduled to have surgery or other procedures.

**The most common side effects of MOUNJARO include:**

- nausea
- diarrhea
- decreased appetite
- vomiting
- constipation
- indigestion
- stomach (abdominal) pain

Talk to your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of MOUNJARO. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**How should I store MOUNJARO?**

- Store MOUNJARO in the refrigerator between 36°F to 46°F (2°C to 8°C). Store MOUNJARO in the original carton until use to protect it from light.
- If needed, each single-dose pen or single-dose vial can be stored at room temperature up to 86°F (30°C) for up to 21 days.
- Do not freeze MOUNJARO. Do not use MOUNJARO if frozen.

**Keep MOUNJARO and all medicines out of the reach of children.**

**General information about the safe and effective use of MOUNJARO.**

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use MOUNJARO for a condition for which it was not prescribed. Do not give MOUNJARO to other people, even if they have the same symptoms you have. It may harm them. You can ask your pharmacist or healthcare provider for information about MOUNJARO that is written for health professionals.

**What are the ingredients in MOUNJARO?**

**Active ingredient:** tirzepatide

**Inactive ingredients:** sodium chloride, sodium phosphate dibasic heptahydrate, and water for injection. Hydrochloric acid solution and/or sodium hydroxide solution may

have been added to adjust the pH.

MOUNJARO® is a registered trademark of Eli Lilly and Company.

Marketed by: Lilly USA, LLC Indianapolis, IN 46285, USA

Copyright © 2022, 2025, Eli Lilly and Company. All rights reserved.

For more information, go to [www.MOUNJARO.com](http://www.MOUNJARO.com) or call 1-800-545-5979.

This Medication Guide has been approved by the U.S. Food and  
Drug Administration

Revised: May 2025

MOU-0004-MG-20250528

## **Mounjaro Single-Dose Pen Instructions for Use**

### **INSTRUCTIONS FOR USE**

**MOUNJARO® (mown-JAHR-OH)**

**(tirzepatide)**

**injection, for subcutaneous use**



**2.5 mg/0.5 mL single-dose pen**

**5 mg/0.5 mL single-dose pen**

**7.5 mg/0.5 mL single-dose pen**

**10 mg/0.5 mL single-dose pen**

**12.5 mg/0.5 mL single-dose pen**

**15 mg/0.5 mL single-dose pen**

**use 1 time each week**


### **Important information you need to know before injecting MOUNJARO**

**Read this Instructions for Use and the Medication Guide before using your MOUNJARO Pen and each time you get a refill.** There may be new information. This information does not take the place of talking to your healthcare provider about your medical condition or treatment.

Talk to your healthcare provider about how to inject MOUNJARO the right way.

- MOUNJARO is a single-dose prefilled pen.
- MOUNJARO is used 1 time each week.
- Inject under the skin (subcutaneously) only.
- You or another person can inject into your stomach (abdomen) or thigh.
- Another person can inject into the back of your upper arm.

## Guide to parts



## Preparing to inject MOUNJARO

**Remove the Pen** from the refrigerator.  
Leave the gray base cap on until you are ready to inject.



## Check the Pen

label to make sure you have the right medicine and dose, and that it has not expired.



## Inspect the Pen

to make sure that it is not damaged.

Make sure the medicine:

- is not frozen
- is colorless to slightly yellow
- is not cloudy
- does not have particles


## Wash your hands.

### Step 1

#### Choose your injection site

Your healthcare provider can help you choose the injection site that is best for you.


You or another person can inject the medicine in your stomach (abdomen) or thigh.



Another person should give you the injection in the back of your upper arm.


Change (rotate) your injection site each week.

You may use the same area of your body but be sure to choose a different injection site in that area.



### Step 2

#### Pull off the gray base cap



Make sure the Pen is **locked**.

**Do not** unlock the Pen until you place the clear base on your skin and are ready to inject.

**Pull** the gray base cap straight off and throw it away in your household trash.


**Do not** put the gray base cap back on – this could damage the needle.

**Do not** touch the needle.


### Step 3

#### Place clear base on skin, then unlock

**Place** the clear base flat against your skin at the injection site.



**Unlock** by turning the lock ring.



### Step 4

#### Press and hold up to 10 seconds


**Press and hold** the purple injection button for up to 10 seconds.

**Listen** for:

- First click = injection started
- Second click = injection completed



You will know your injection is complete when the gray plunger is



**After your injection, place the used Pen in a sharps container.**  
See **Disposing of your used Pen.**

### **Disposing of your used Pen**

- Put your used Pen in an FDA-cleared sharps disposal container right away after use. **Do not** throw away (dispose of) Pens in your household trash.
- If you do not have an FDA-cleared sharps disposal container, you may use a household container that is:
  - made of a heavy-duty plastic,
  - can be closed with a tight-fitting, puncture-resistant lid, without sharps being able to come out,
  - upright and stable during use,
  - leak-resistant, and
  - properly labeled to warn of hazardous waste inside the container.
- When your sharps disposal container is almost full, you will need to follow your community guidelines for the right way to dispose of your sharps disposal container. There may be state or local laws about how you should throw away used needles and syringes. For more information about safe sharps disposal, and for specific information about sharps disposal in the state that you live in, go to the FDA's website at: <http://www.fda.gov/safesharpsdisposal>.
- **Do not** recycle your used sharps disposal container.

### **Storage and handling**

- Store your Pen in the refrigerator between 36°F to 46°F (2°C to 8°C).
- You may store your Pen at room temperature up to 86°F (30°C) for up to 21 days.

- **Do not** freeze your Pen. If the Pen has been frozen, throw the Pen away and use a new Pen.
- Store your Pen in the original carton to protect your Pen from light.
- The Pen has glass parts. Handle it carefully. If you drop the Pen on a hard surface, **do not** use it. Use a new Pen for your injection.
- Keep your MOUNJARO Pen and all medicines out of the reach of children.

## Commonly asked questions

### What if I see air bubbles in my Pen?

Air bubbles are normal.

### What if my Pen is not at room temperature?

It is not necessary to warm the Pen to room temperature.

### What if I unlock the Pen and press the purple injection button before pulling off the gray base cap?

**Do not** remove the gray base cap. Throw away the Pen and get a new Pen.

### What if there is a drop of liquid on the tip of the needle when I remove the gray base cap?

A drop of liquid on the tip of the needle is normal. **Do not** touch the needle.

### Do I need to hold the injection button down until the injection is complete?

This is not necessary, but it may help you keep the Pen steady against your skin.

### I heard more than 2 clicks during my injection—2 loud clicks and 1 soft one. Did I get my complete injection?

Some people may hear a soft click right before the second loud click. That is the normal operation of the Pen. **Do not** remove the Pen from your skin until you hear the second loud click.

### I am not sure if my Pen worked the right way.



Check to see if you have received your dose. Your dose was delivered the right way if the gray plunger is visible. Also, see **Step 4** of the instructions.

If you do not see the gray plunger, contact Lilly at 1-800-Lilly-Rx (1-800-545-5979) for further instructions. Until then, store your Pen safely to avoid an accidental needle stick.

### What if there is a drop of liquid or blood on my skin after my injection?

This is normal. Press a cotton ball or gauze over the injection site. **Do not** rub the injection site.

## Other information

- If you have vision problems, **do not** use your Pen without help from a person trained to use the MOUNJARO Pen.

## Where to learn more

- If you have questions or problems with your MOUNJARO Pen, contact Lilly at 1-800-Lilly-Rx (1-800-545-5979) or call your healthcare provider.
- For more information about the MOUNJARO Pen, visit our website at [www.mounjaro.com](http://www.mounjaro.com).



**Scan this code to launch  
[www.mounjaro.com](http://www.mounjaro.com)**

Marketed by:

Lilly USA, LLC

Indianapolis, IN 46285, USA

MOUNJARO is a registered trademark of Eli Lilly and Company.

Copyright © 2022, 2024, Eli Lilly and Company. All rights reserved.

This Instructions for Use has been approved by the U.S. Food and Drug Administration.  
Revised: May 2024

MOU-0003-IFU-20240516



## Mounjaro Single-Dose Vial Instructions for Use

### INSTRUCTIONS FOR USE

**MOUNJARO® [mown-JAHR-OH]**

**(tirzepatide)**

**injection, for subcutaneous use**

**2.5 mg/0.5 mL single-dose vial**

**5 mg/0.5 mL single-dose vial**

**7.5 mg/0.5 mL single-dose vial**

**10 mg/0.5 mL single-dose vial**

**12.5 mg/0.5 mL single-dose vial**

**15 mg/0.5 mL single-dose vial**

## Important information you need to know before injecting MOUNJARO

**Read this Instructions for Use before you start taking MOUNJARO and each time you get a new vial.** There may be new information. This information does not take the place of talking to your healthcare provider about your medical condition or your treatment.

**Do not share your needles or syringes with other people. You may give other people a serious infection or get a serious infection from them.**


Talk to your healthcare provider about how to inject MOUNJARO the right way.

- MOUNJARO is a single-dose vial.
- MOUNJARO is used 1 time each week.
- Inject under the skin (subcutaneously) only.
- You or another person may inject into your stomach (abdomen) or thigh.
- Another person can inject into the back of your upper arm.

### Gather supplies needed to give your injection

- 1 single-dose MOUNJARO vial
- 1 syringe and 1 needle, supplied separately (for example, use a 1 mL syringe and needle as recommended by your healthcare provider)
- 1 alcohol swab
- gauze
- 1 sharps container for throwing away used needles and syringes. **See** “Disposing of used needles and syringes” at the end of these instructions.

### Guide to parts



**Note:** The needle and syringe are not included. The needle and syringe recommended by your healthcare provider may look different than the needle and syringe in this Instructions for Use.

### Preparing to inject MOUNJARO

**Remove the vial** from the refrigerator.

**Check the vial** label to make sure you have the right medicine and dose, and that it has

not expired.

Make sure the medicine:


- is not frozen
- is not cloudy
- is colorless to slightly yellow
- does not have particles

**Always use a new syringe and needle for each injection to prevent infections and blocked needles. Do not reuse or share your syringes or needles with other people. You may give other people a serious infection or get a serious infection from them.**

**Wash your hands with soap and water.**

**Step 1:**

Pull off the plastic protective cap. **Do not** remove the rubber stopper.




**Step 2:**

Wipe the rubber stopper with an alcohol swab.




**Step 3:**


Remove the outer wrapping from the syringe.

**Step 4:**


Remove the outer wrapping from the needle. The syringe that your healthcare provider recommended may have a pre-attached needle. If the needle is attached, skip to step 6.

**Step 5:**

Place the needle on top of the syringe and turn until it is tight and firmly attached.

**Step 6:**


Remove the needle shield by pulling straight off.






**Step 7:**


Hold the syringe in one hand with the needle pointing up. With the other hand pull down on the plunger until the plunger tip reaches the line on the syringe indicating that 0.5 mL of air has been drawn into the syringe.

**Step 8:**

Push the needle through the rubber stopper of the vial.



**Step 9:**

Push the plunger all the way in. This puts air into the vial and makes it easier to pull the solution from the vial.




**Step 10:**

Turn the vial and syringe upside down. Make sure that the tip of the needle is in the liquid and slowly pull the plunger down until the plunger tip is past the 0.5 mL line. If there are air bubbles, tap the syringe gently a few times to let any air bubbles rise to the top.


**Step 11:**

Slowly push the plunger up until the plunger tip reaches the 0.5 mL line.



**Step 12:**

Pull the syringe out of the rubber stopper of the vial.

**Injecting MOUNJARO**


- Inject exactly as your healthcare provider has shown you. Your healthcare provider should tell you if you should pinch the skin before injecting.
- **Change (rotate) your injection site within the area you choose for each dose** to reduce your risk of getting lipodystrophy (pits in skin or thickened skin) and localized cutaneous amyloidosis (skin with lumps) at the injection sites.
- **Do not** inject where the skin has pits, is thickened, or has lumps.
- **Do not** inject where the skin is tender, bruised, scaly or hard, or into scars or damaged skin.
- **Do not** mix MOUNJARO with any other medicine.
- **Do not** inject MOUNJARO in the same injection site used for other medicines.


**Step 13:**

Choose your injection site.

You can inject MOUNJARO under the skin (subcutaneously) of your stomach area (abdomen) or thighs.

Someone else can inject in your stomach area, thighs, or the back of the upper arms.






**Step 14:**

Insert the needle into your skin.



**Step 15:**


Push down on the plunger to inject your dose. The needle should stay in your skin for at least 5 seconds to make sure you have injected all of your dose.



**Step 16:**

Pull the needle out of your skin.

- If you see blood after you take the needle out of your skin, press the injection site with a piece of gauze or an alcohol swab. **Do not** rub the area.
- **Do not** recap the needle. Recapping the needle can lead to a needle stick injury.

**Disposing of used needles and syringes**

- Put your used needle and syringe in an FDA-cleared sharps disposal container right away after use. **Do not** throw away (dispose of) loose needles and syringes in your household trash.
- If you do not have an FDA-cleared sharps disposal container, you may use a household container that is:
  - made of a heavy-duty plastic,
  - can be closed with a tight-fitting, puncture-resistant lid, without sharps being able to come out,
  - upright and stable during use,
  - leak-resistant, and
  - properly labeled to warn of hazardous waste inside the container.
- When your sharps disposal container is almost full, you will need to follow your community guidelines for the right way to dispose of your sharps disposal container. There may be state or local laws about how you should throw away used needles and syringes. For more information about safe sharps disposal, and for specific information about sharps disposal in the state that you live in, go to the FDA's website at:  
<http://www.fda.gov/safesharpsdisposal>.
- **Do not** dispose of your used sharps disposal container in your household trash unless your community guidelines permit this. **Do not** recycle your used sharps disposal container.

**Storing MOUNJARO**

- Store all unopened vials in the refrigerator at 36°F to 46°F (2°C to 8°C).
- You may store the unopened vial at room temperature up to 86°F (30°C) for up to 21 days.
- **Do not** freeze. **Do not** use if MOUNJARO has been frozen.
- Store the vial in the original carton to protect from light.
- Throw away all opened vials after use, even if there is medicine left in the vial.

**Keep MOUNJARO vials, syringes, needles, and all medicines out of the reach**

**of children.**

If you have any questions or problems with your MOUNJARO, contact Lilly at 1-800-Lilly-Rx (1-800-545-5979) or call your healthcare provider for help.

**Marketed by:**

**Lilly USA, LLC**

**Indianapolis, IN 46285, USA**

MOUNJARO is a registered trademark of Eli Lilly and Company.

Copyright © 2023, 2024, Eli Lilly and Company. All rights reserved.

MON-VL-0002-IFU-20240229

This Instructions for Use has been approved by the U.S. Food and Drug Administration.

Revised: February 2024

**PACKAGE LABEL - Mounjaro<sup>®</sup>, 2.5 mg/0.5 mL, Carton, 4 Single-Dose Pens**

NDC 0002-1506-80

Mounjaro<sup>®</sup>

(tirzepatide) injection

2.5 mg/0.5 mL

Use one pen every week.

4 Single-dose prefilled pens

4 x 2.5 mg/0.5 mL prefilled pens

Rx Only

For Subcutaneous Use

Dispense enclosed Medication Guide to each patient.

Lilly



NDC 0002-1495-80

Mounjaro®

(tirzepatide) injection

5 mg/0.5 mL

Use one pen every week.

4 Single-dose prefilled pens

4 x 5 mg/0.5 mL prefilled pens

Rx Only

For Subcutaneous Use

Dispense enclosed Medication Guide to each patient.

Lilly



**mounjaro**<sup>®</sup>  
(tirzepatide) Injection

5 mg  
0.5 mL



4 Single-dose  
prefilled pens

**mounjaro**<sup>®</sup>  
(tirzepatide) Injection

5 mg  
0.5 mL



4 Single-dose  
prefilled pens



3 0002-1495-80 0

www.mounjaro.com

In the map, U.S. 46385, USA

Made by:

Lilly USA, LLC

No U.S. side effect of potency.

No preservative.

MOUNJARO has a pH of 6.5 to 7.5.

hydrolytic solution may have been added to adjust the pH.

Each 0.5 mL of MOUNJARO contains 5 mg tirzepatide, sodium

chloride (4.1 mg), and turn potassium chloride (0.7 mg).

Instructions for use

Package Insert

4 single-dose prefilled pens

Carton contains:

Recommended Dosage: See prescribing information.

Keep out of reach of children.

Remove from the refrigerator.

up to 21 days in the carton. Discard if not used within 21 days after

MOUNJARO is a sterile, non-pyrogenic, non-aqueous solution.

Protect from light. DO NOT FREEZE.

Store refrigerated at 2°C to 8°C (36°F to 46°F) in original carton to

PHARMACY  
LABEL HERE

**mounjaro**<sup>®</sup>  
(tirzepatide) Injection

5 mg  
0.5 mL



4 Single-dose  
prefilled pens

4 Single-dose  
prefilled pens

5 mg  
0.5 mL

**mounjaro**<sup>®</sup>  
(tirzepatide) Injection

LOT: 0002149580  
LAL/EP/SEA/NO

NDC 0002-1495-80

**mounjaro**<sup>®</sup>  
(tirzepatide) injection

4 x 5 mg/0.5 mL prefilled pens

Rx Only

For Subcutaneous Use

Dispense enclosed Medication  
Guide to each patient.

Use one pen every week.

4 Single-dose  
prefilled pens

5 mg  
0.5 mL



Lilly

**PACKAGE LABEL - Mounjaro<sup>®</sup>, 7.5 mg/0.5 mL, Carton, 4 Single-Dose Pens**

NDC 0002-1484-80

Mounjaro<sup>®</sup>

(tirzepatide) injection

7.5 mg/0.5 mL

Use one pen every week.

4 Single-dose prefilled pens

4 x 7.5 mg/0.5 mL prefilled pens

Rx Only

For Subcutaneous Use

Dispense enclosed Medication Guide to each patient.

Lilly

**mounjaro**<sup>®</sup>  
(tirzepatide) Injection

**7.5**  $\frac{\text{mg}}{0.5 \text{ mL}}$



4 Single-dose  
prefilled pens

**mounjaro**<sup>®</sup>  
(tirzepatide) Injection

**7.5**  $\frac{\text{mg}}{0.5 \text{ mL}}$



4 Single-dose  
prefilled pens



3 0002-1484-80 4

www.mounjaro.com

In the map, see, IN 46285, USA

Made by:

Lilly USA, LLC

No U.S. side of pen.

No preservative.

MOUNJARO has a H of 6.5 to 7.5.

Hydrochloric acid solution may have been added to adjust the pH.

Each 0.5 mL of MOUNJARO contains 7.5 mg tirzepatide, sodium

chloride (4.1 mg) and sodium phosphate buffer (0.7 mg).

Instructions for Use

Medication Guide

Package Insert

4 Single-dose prefilled pens

Caution: Do not use

Reconstituted Dosage: See prescribing information.

Keep out of reach of children.

Remove from the refrigerator.

Use within 21 days in the carton. Discard if not used within 21 days after

MOUNJARO is stored at room temperature up to 30 °C (86 °F) for

up to 21 days. DO NOT FREEZE.

Protect from light. DO NOT EXPOSE TO

PHARMACY  
LABEL HERE



4 Single-dose  
prefilled pens

**7.5**  $\frac{\text{mg}}{0.5 \text{ mL}}$

**mounjaro**<sup>®</sup>  
(tirzepatide) Injection

4 Single-dose  
prefilled pens

**7.5**  $\frac{\text{mg}}{0.5 \text{ mL}}$

**mounjaro**<sup>®</sup>  
(tirzepatide) Injection

GTIN: 0 00021484 80  
Lot / Exp / Serial No

NDC 0002-1484-80

**mounjaro**<sup>®</sup>  
(tirzepatide) injection

4 x 7.5 mg/0.5 mL prefilled pens

Rx Only

For Subcutaneous Use

Dispense enclosed Medication  
Guide to each patient.

Use one pen every week.

**4** Single-dose  
prefilled pens

**7.5**  $\frac{\text{mg}}{0.5 \text{ mL}}$



Lilly

**PACKAGE LABEL - Mounjaro<sup>®</sup>, 10 mg/0.5 mL, Carton, 4 Single-Dose Pens**

NDC 0002-1471-80

Mounjaro<sup>®</sup>

(tirzepatide) injection

10 mg/0.5 mL

Use one pen every week.

4 Single-dose prefilled pens

4 x 10 mg/0.5 mL prefilled pens

Rx Only

For Subcutaneous Use

Dispense enclosed Medication Guide to each patient.

Lilly

**mounjaro®**  
(tirzepatide) Injection

**10 mg**  
**0.5 mL**



4 Single-dose  
prefilled pens

**mounjaro®**  
(tirzepatide) Injection

**10 mg**  
**0.5 mL**



4 Single-dose  
prefilled pens



www.mounjaro.com

Lilly USA, LLC  
Incl. map. 01, IN 46285, USA

No U.S. standard of potency.

No preservative.

MOUNJARO has a pH of 6.5 to 7.5.

Each 0.5 mL of MOUNJARO contains 10 mg tirzepatide, sodium chloride, and water for injection. Hydrochloric acid, sodium hydroxide solution may have been added to adjust the pH.

Instructions for Use

Package Insert

Caution: Do not use prefilled pen

Recommended Dosage: See prescribing information.

Keep out of reach of children.

Remove from the refrigerator.

MOUNJARO should be stored at room temperature up to 30°C (86°F) for up to 21 days in the carton. Discard if not used within 21 days after first use.

Protect from light. DO NOT FREEZE.

Store refrigerated at 2°C to 8°C (36°F to 46°F) in original carton.

PHARMACY  
LABEL HERE

**mounjaro®**  
(tirzepatide) Injection

**10 mg**  
**0.5 mL**



4 Single-dose  
prefilled pens

**mounjaro®**  
(tirzepatide) Injection

**10 mg**  
**0.5 mL**

4 Single-dose  
prefilled pens

61N-0000021-0101  
Lot / Exp / Serial No

NDC 0002-1471-80

**mounjaro®**  
(tirzepatide) injection

4 x 10 mg/0.5 mL prefilled pens

Rx Only

For Subcutaneous Use

Dispense enclosed Medication  
Guide to each patient.

Use one pen every week.

**4** Single-dose  
prefilled pens

**10 mg**  
**0.5 mL**



*Lilly*

**PACKAGE LABEL - Mounjaro<sup>®</sup>, 12.5 mg/0.5 mL, Carton, 4 Single-Dose Pens**

NDC 0002-1460-80

Mounjaro<sup>®</sup>

(tirzepatide) injection

12.5 mg/0.5 mL

Use one pen every week.

4 Single-dose prefilled pens

4 x 12.5 mg/0.5 mL prefilled pens

Rx Only

For Subcutaneous Use

Dispense enclosed Medication Guide to each patient.

Lilly

**mounjaro**<sup>®</sup>  
(tirzepatide) Injection

**12.5**  $\frac{\text{mg}}{0.5 \text{ mL}}$



4 Single-dose  
prefilled pens

**mounjaro**<sup>®</sup>  
(tirzepatide) Injection

**12.5**  $\frac{\text{mg}}{0.5 \text{ mL}}$



4 Single-dose  
prefilled pens



3 0002-1460-80 8

www.mounjaro.com

Lilly USA, LLC  
Indianapolis, IN 46205, USA

No preservative.  
No U.S. standard of potency.

MOUNJARO has a pH of 6.5 to 7.5.  
Hydrochloric acid has been added to adjust the pH.  
Each 0.5 mL of MOUNJARO contains 12.5 mg tirzepatide; sodium  
chloride 9.1 mg, sodium phosphate dibasic heptahydrate 10.7 mg.  
Packaging in sets of 4 single-dose prefilled pens.  
Medication Guide  
Instructions for Use  
Each 0.5 mL of MOUNJARO contains 12.5 mg tirzepatide; sodium  
chloride 9.1 mg, sodium phosphate dibasic heptahydrate 10.7 mg.  
Hydrochloric acid has been added to adjust the pH.  
MOUNJARO has a pH of 6.5 to 7.5.  
No preservative.  
No U.S. standard of potency.

PHARMACY  
LABEL HERE



**12.5**  $\frac{\text{mg}}{0.5 \text{ mL}}$

**mounjaro**<sup>®</sup>  
(tirzepatide) Injection

4 Single-dose  
prefilled pens

**12.5**  $\frac{\text{mg}}{0.5 \text{ mL}}$

**mounjaro**<sup>®</sup>  
(tirzepatide) Injection

NDC 0002-1460-80

**mounjaro**<sup>®</sup>  
(tirzepatide) injection

4 x 12.5 mg/0.5 mL prefilled pens

Rx Only

For Subcutaneous Use

Dispense enclosed Medication  
Guide to each patient.

Use one pen every week.

**4** Single-dose  
prefilled pens

**12.5**  $\frac{\text{mg}}{0.5 \text{ mL}}$



*Lilly*

**PACKAGE LABEL - Mounjaro<sup>®</sup>, 15 mg/0.5 mL, Carton, 4 Single-Dose Pens**

NDC 0002-1457-80

Mounjaro<sup>®</sup>

(tirzepatide) injection

15 mg/0.5 mL

Use one pen every week.

4 Single-dose prefilled pens

4 x 15 mg/0.5 mL prefilled pens

Rx Only

For Subcutaneous Use

Dispense enclosed Medication Guide to each patient.

Lilly



**mounjaro**<sup>®</sup>  
(tirzepatide) Injection

**15**  $\frac{\text{mg}}{0.5 \text{ mL}}$



4 Single-dose  
prefilled pens

**mounjaro**<sup>®</sup>  
(tirzepatide) Injection

**15**  $\frac{\text{mg}}{0.5 \text{ mL}}$



4 Single-dose  
prefilled pens

SHOT 10AAM 01



www.mounjaro.com

Incl. 46 285, U.S.A.

Manufactured by:

No U.S. standard of potency.

No preservative.

MOUNJARO has a pH of 6.5 to 7.5.

Hydrochloric acid solution may have been added to adjust the pH.

an dewatered or injection. Hydrochloric acid solution and/or sodium

chloride (0.1 mg), sodium phosphate dibasic heptahydrate (0.7 mg),

Each 0.5 mL of MOUNJARO contains 15 mg tirzepatide, sodium

injection. See package insert for full prescribing information.

Packaging in cartons

4 Single-dose prefilled pens

Carton Contents:

Recommendation: See prescribing information.

Keep out of reach of children.

Remove from the refrigerator

up to 21 days in the carton. Do not use within 21 days after

MOUNJARO can be stored at room temperature up to 30°C (86°F) for

protection from light. DO NOT FREEZE.

Store refrigerated at 2°C to 8°C (36°F to 46°F) in original carton to

PHARMACY  
LABEL HERE

**mounjaro**<sup>®</sup>  
(tirzepatide) Injection

**15**  $\frac{\text{mg}}{0.5 \text{ mL}}$



4 Single-dose  
prefilled pens

4 Single-dose  
prefilled pens

**15**  $\frac{\text{mg}}{0.5 \text{ mL}}$

**mounjaro**<sup>®</sup>  
(tirzepatide) Injection

Lot / Exp / Serial No  
B 1 N: 0000 021 20 808

NDC 0002-1457-80

**mounjaro**<sup>®</sup>  
(tirzepatide) Injection

4 x 15 mg/0.5 mL prefilled pens

Rx Only

For Subcutaneous Use

Dispense enclosed Medication  
Guide to each patient.

Use one pen every week.

**4** Single-dose  
prefilled pens

**15**  $\frac{\text{mg}}{0.5 \text{ mL}}$



*Lilly*

**PACKAGE LABEL - Mounjaro<sup>®</sup>, 2.5 mg/0.5 mL, Carton, Single-dose Vial**

NDC 0002-1152-01

Mounjaro<sup>®</sup>

(tirzepatide) injection

2.5 mg/0.5 mL

Single-dose Vial - discard unused portion


Rx Only

For Subcutaneous Use

Needles and Syringes are not included

Dispense enclosed Medication Guide to each patient.

Lilly



## PACKAGE LABEL - Mounjaro®, 5 mg/0.5 mL, Single-dose Vial

NDC 0002-1243-01

Mounjaro®

(tirzepatide) injection

5 mg/0.5 mL

Single-dose Vial - discard unused portion

Rx Only

For Subcutaneous Use

Needles and Syringes are not included

Dispense enclosed Medication Guide to each patient.

Lilly



## PACKAGE LABEL - Mounjaro®, 7.5 mg/0.5 mL, Single-dose Vial

NDC 0002-2214-01

Mounjaro®

(tirzepatide) injection

7.5 mg/0.5 mL


Single-dose Vial - discard unused portion

Rx Only

For Subcutaneous Use

Needles and Syringes are not included

Dispense enclosed Medication Guide to each patient.



**PACKAGE LABEL - Mounjaro®, 10 mg/0.5 mL, Carton, Single-dose Vial**

NDC 0002-2340-01

Mounjaro®

(tirzepatide) injection

10 mg/0.5 mL

Single-dose Vial - discard unused portion


Rx Only

For Subcutaneous Use

Needles and Syringes are not included

Dispense enclosed Medication Guide to each patient.

Lilly



**PACKAGE LABEL - Mounjaro®, 12.5 mg/0.5 mL, Single-dose Vial**

NDC 0002-2423-01

Mounjaro®

(tirzepatide) injection

12.5 mg/0.5 mL

Single-dose Vial - discard unused portion

Rx Only


For Subcutaneous Use



Needles and Syringes are not included

Dispense enclosed Medication Guide to each patient.

Lilly



**PACKAGE LABEL - Mounjaro®, 15 mg/0.5 mL, Single-dose Vial**

NDC 0002-3002-01

Mounjaro®

(tirzepatide) injection

15 mg/0.5 mL

Single-dose Vial - discard unused portion


Rx Only

For Subcutaneous Use

Needles and Syringes are not included

Dispense enclosed Medication Guide to each patient.

Lilly



MOUNJARO			
tirzepatide injection, solution			
Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0002-1506
Route of Administration	SUBCUTANEOUS		



Active Ingredient/Active Moiety				
Ingredient Name			Basis of Strength	Strength
<b>tirzepatide</b> (UNII: OYN3CCI6QE) (tirzepatide - UNII:OYN3CCI6QE)			tirzepatide	2.5 mg in 0.5 mL
Inactive Ingredients				
Ingredient Name				Strength
<b>Sodium Phosphate, Dibasic, Heptahydrate</b> (UNII: 70WT22SF4B)				0.7 mg in 0.5 mL
<b>Sodium Chloride</b> (UNII: 451W47IQ8X)				4.1 mg in 0.5 mL
<b>Hydrochloric Acid</b> (UNII: QTT17582CB)				
<b>Sodium Hydroxide</b> (UNII: 55X04QC32I)				
<b>Water</b> (UNII: 059QF0KO0R)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0002-1506-80	4 in 1 CARTON	05/13/2022	
1	NDC:0002-1506-01	0.5 mL in 1 SYRINGE; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		
2	NDC:0002-1506-61	4 in 1 CARTON	06/08/2022	
2		0.5 mL in 1 SYRINGE; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA		NDA215866	05/13/2022	

MOUNJARO			
tirzepatide injection, solution			
Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0002-1495
Route of Administration	SUBCUTANEOUS		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
tirzepatide (UNII: OYN3CCI6QE) (tirzepatide - UNII:OYN3CCI6QE)		tirzepatide	5 mg in 0.5 mL
Inactive Ingredients			

Ingredient Name			Strength	
Sodium Phosphate, Dibasic, Heptahydrate (UNII: 70WT22SF4B)			0.7 mg	in 0.5 mL
Sodium Chloride (UNII: 451W47IQ8X)			4.1 mg	in 0.5 mL
Hydrochloric Acid (UNII: QTT17582CB)				
Sodium Hydroxide (UNII: 55X04QC32I)				
Water (UNII: 059QF0KO0R)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0002-1495-80	4 in 1 CARTON	05/13/2022	
1	NDC:0002-1495-01	0.5 mL in 1 SYRINGE; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA		NDA215866	05/13/2022	

MOUNJARO				
tirzepatide injection, solution				
Product Information				
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0002-1484	
Route of Administration	SUBCUTANEOUS			
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
tirzepatide (UNII: OYN3CCI6QE) (tirzepatide - UNII:OYN3CCI6QE)		tirzepatide	7.5 mg in 0.5 mL	
Inactive Ingredients				
Ingredient Name			Strength	
Sodium Phosphate, Dibasic, Heptahydrate (UNII: 70WT22SF4B)			0.7 mg in 0.5 mL	
Sodium Chloride (UNII: 451W47IQ8X)			4.1 mg in 0.5 mL	
Hydrochloric Acid (UNII: QTT17582CB)				
Sodium Hydroxide (UNII: 55X04QC32I)				
Water (UNII: 059QF0KO0R)				
Packaging				
#	Item	Package Description	Marketing	Marketing

#	Code	Package Description	Start Date	End Date
1	NDC:0002-1484-80	4 in 1 CARTON	05/13/2022	
1	NDC:0002-1484-01	0.5 mL in 1 SYRINGE; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		
<b>Marketing Information</b>				
<b>Marketing Category</b>		<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
NDA		NDA215866	05/13/2022	

<b>MOUNJARO</b>				
tirzepatide injection, solution				
<b>Product Information</b>				
<b>Product Type</b>		HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b> NDC:0002-1471	
<b>Route of Administration</b>		SUBCUTANEOUS		
<b>Active Ingredient/Active Moiety</b>				
<b>Ingredient Name</b>			<b>Basis of Strength</b>	<b>Strength</b>
tirzepatide (UNII: OYN3CCI6QE) (tirzepatide - UNII:OYN3CCI6QE)			tirzepatide	10 mg in 0.5 mL
<b>Inactive Ingredients</b>				
<b>Ingredient Name</b>				<b>Strength</b>
Sodium Phosphate, Dibasic, Heptahydrate (UNII: 70WT22SF4B)				0.7 mg in 0.5 mL
Sodium Chloride (UNII: 451W47IQ8X)				4.1 mg in 0.5 mL
Hydrochloric Acid (UNII: QTT17582CB)				
Sodium Hydroxide (UNII: 55X04QC32I)				
Water (UNII: 059QF0KO0R)				
<b>Packaging</b>				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0002-1471-80	4 in 1 CARTON	05/13/2022	
1	NDC:0002-1471-01	0.5 mL in 1 SYRINGE; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		
<b>Marketing Information</b>				
<b>Marketing Category</b>		<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
NDA		NDA215866	05/13/2022	

## MOUNJARO

tirzepatide injection, solution

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0002-1460
Route of Administration	SUBCUTANEOUS		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
tirzepatide (UNII: OYN3CCI6QE) (tirzepatide - UNII:OYN3CCI6QE)	tirzepatide	12.5 mg in 0.5 mL

### Inactive Ingredients

Ingredient Name	Strength
Sodium Phosphate, Dibasic, Heptahydrate (UNII: 70WT22SF4B)	0.7 mg in 0.5 mL
Sodium Chloride (UNII: 451W47IQ8X)	4.1 mg in 0.5 mL
Hydrochloric Acid (UNII: QTT17582CB)	
Sodium Hydroxide (UNII: 55X04QC32I)	
Water (UNII: 059QF0KO0R)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0002-1460-80	4 in 1 CARTON	05/13/2022	
1	NDC:0002-1460-01	0.5 mL in 1 SYRINGE; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA215866	05/13/2022	

## MOUNJARO

tirzepatide injection, solution

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0002-1457
Route of Administration	SUBCUTANEOUS		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
<b>tirzepatide</b> (UNII: OYN3CCI6QE) (tirzepatide - UNII:OYN3CCI6QE)	tirzepatide	15 mg in 0.5 mL

Inactive Ingredients	
Ingredient Name	Strength
<b>Sodium Phosphate, Dibasic, Heptahydrate</b> (UNII: 70WT22SF4B)	0.7 mg in 0.5 mL
<b>Sodium Chloride</b> (UNII: 451W47IQ8X)	4.1 mg in 0.5 mL
<b>Hydrochloric Acid</b> (UNII: QTT17582CB)	
<b>Sodium Hydroxide</b> (UNII: 55X04QC32I)	
<b>Water</b> (UNII: 059QF0KO0R)	

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0002-1457-80	4 in 1 CARTON	05/13/2022	
1	NDC:0002-1457-01	0.5 mL in 1 SYRINGE; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA215866	05/13/2022	

## MOUNJARO

tirzepatide injection, solution

Product Information			
<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:0002-1152
<b>Route of Administration</b>	SUBCUTANEOUS		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
<b>tirzepatide</b> (UNII: OYN3CCI6QE) (tirzepatide - UNII:OYN3CCI6QE)	tirzepatide	2.5 mg in 0.5 mL

Inactive Ingredients	
Ingredient Name	Strength
<b>Sodium Chloride</b> (UNII: 451W47IQ8X)	4.1 mg in 0.5 mL
<b>Sodium Phosphate, Dibasic, Heptahydrate</b> (UNII: 70WT22SF4B)	0.7 mg in 0.5 mL

Hydrochloric Acid (UNII: QTT17582CB)				
Sodium Hydroxide (UNII: 55X04QC32I)				
Water (UNII: 059QF0KO0R)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0002-1152-01	1 in 1 CARTON	07/28/2023	
1		0.5 mL in 1 VIAL, SINGLE-DOSE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA		NDA215866	07/28/2023	

## MOUNJARO

tirzepatide injection, solution

Product Information				
Product Type		HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0002-1243
Route of Administration		SUBCUTANEOUS		
Active Ingredient/Active Moiety				
Ingredient Name			Basis of Strength	Strength
tirzepatide (UNII: OYN3CCI6QE) (tirzepatide - UNII:OYN3CCI6QE)			tirzepatide	5 mg in 0.5 mL
Inactive Ingredients				
Ingredient Name			Strength	
Sodium Chloride (UNII: 451W47IQ8X)			4.1 mg in 0.5 mL	
Sodium Phosphate, Dibasic, Heptahydrate (UNII: 70WT22SF4B)			0.7 mg in 0.5 mL	
Hydrochloric Acid (UNII: QTT17582CB)				
Sodium Hydroxide (UNII: 55X04QC32I)				
Water (UNII: 059QF0KO0R)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0002-1243-01	1 in 1 CARTON	07/28/2023	
		0.5 mL in 1 VIAL, SINGLE DOSE; Type 0: Not a		

1	0.5 mL in 1 VIAL, SINGLE-DOSE; Type 0: Not a Combination Product		
Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA215866	07/28/2023	

## MOUNJARO

tirzepatide injection, solution

Product Information				
Product Type		HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0002-2214
Route of Administration		SUBCUTANEOUS		
Active Ingredient/Active Moiety				
Ingredient Name			Basis of Strength	Strength
tirzepatide (UNII: OYN3CCI6QE) (tirzepatide - UNII:OYN3CCI6QE)			tirzepatide	7.5 mg in 0.5 mL
Inactive Ingredients				
Ingredient Name				Strength
Sodium Chloride (UNII: 451W47IQ8X)				4.1 mg in 0.5 mL
Sodium Phosphate, Dibasic, Heptahydrate (UNII: 70WT22SF4B)				0.7 mg in 0.5 mL
Hydrochloric Acid (UNII: QTT17582CB)				
Sodium Hydroxide (UNII: 55X04QC32I)				
Water (UNII: 059QF0KO0R)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0002-2214-01	1 in 1 CARTON	07/28/2023	
1		0.5 mL in 1 VIAL, SINGLE-DOSE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA		NDA215866	07/28/2023	

# MOUNJARO

tirzepatide injection, solution

## Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0002-2340
Route of Administration	SUBCUTANEOUS		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
tirzepatide (UNII: OYN3CCI6QE) (tirzepatide - UNII:OYN3CCI6QE)	tirzepatide	10 mg in 0.5 mL

## Inactive Ingredients

Ingredient Name	Strength
Sodium Chloride (UNII: 451W47IQ8X)	4.1 mg in 0.5 mL
Sodium Phosphate, Dibasic, Heptahydrate (UNII: 70WT22SF4B)	0.7 mg in 0.5 mL
Hydrochloric Acid (UNII: QTT17582CB)	
Sodium Hydroxide (UNII: 55X04QC32I)	
Water (UNII: 059QF0KO0R)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0002-2340-01	1 in 1 CARTON	07/28/2023	
1		0.5 mL in 1 VIAL, SINGLE-DOSE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA215866	07/28/2023	

# MOUNJARO

tirzepatide injection, solution

## Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0002-2423
Route of Administration	SUBCUTANEOUS		

## Active Ingredient/Active Moiety



Ingredient Name		Basis of Strength	Strength	
tirzepatide (UNII: OYN3CCI6QE) (tirzepatide - UNII:OYN3CCI6QE)		tirzepatide	12.5 mg in 0.5 mL	
Inactive Ingredients				
Ingredient Name			Strength	
Sodium Chloride (UNII: 451W47IQ8X)			4.1 mg in 0.5 mL	
Sodium Phosphate, Dibasic, Heptahydrate (UNII: 70WT22SF4B)			0.7 mg in 0.5 mL	
Hydrochloric Acid (UNII: QTT17582CB)				
Sodium Hydroxide (UNII: 55X04QC32I)				
Water (UNII: 059QF0KO0R)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0002-2423-01	1 in 1 CARTON	07/28/2023	
1		0.5 mL in 1 VIAL, SINGLE-DOSE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA		NDA215866	07/28/2023	

MOUNJARO

tirzepatide injection, solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0002-3002
Route of Administration	SUBCUTANEOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
tirzepatide (UNII: OYN3CCI6QE) (tirzepatide - UNII:OYN3CCI6QE)	tirzepatide	15 mg in 0.5 mL

Inactive Ingredients

Ingredient Name	Strength
Sodium Chloride (UNII: 451W47IQ8X)	4.1 mg in 0.5 mL
Sodium Phosphate, Dibasic, Heptahydrate (UNII: 70WT22SF4B)	0.7 mg in 0.5 mL
Hydrochloric Acid (UNII: QTT17582CB)	
Sodium Hydroxide (UNII: 55X04QC32I)	

**Water** (UNII: 059QF0KO0R)

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0002-3002-01	1 in 1 CARTON	07/28/2023	
1		0.5 mL in 1 VIAL, SINGLE-DOSE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA215866	07/28/2023	

**Labeler** - Eli Lilly and Company (006421325)

Revised: 6/2025

Eli Lilly and Company