

**DRKIDS CHILDREN MULTI SYMPTOM COLD AND FLU- acetaminophen 320mg, chlorpheniramine maleate 2mg, dextromethorphan hbr 10mg, phenylephrine hcl 5mg solution**  
**Calmco**

-----

**DrKids Children's Multi-Symptoms Cold & Flu**

***Active Ingredient (in each 10mL)***

Acetaminophen, 320mg

Chlorpheniramine Maleate, 2mg

Dextromethorphan HBr, 10mg t

Phenylephrine HCl, 5mg

***Purpose***

Pain reliever / fever reducer

Antihistamine

Cough suppressant

Nasal decongestant

***Uses***

- temporarily relieves the following cold/flu symptoms: ■ minor aches and pains ■ headache ■ sore throat ■ cough ■ stuffy nose
- sneezing and runny nose ■ temporarily reduces fever

**Warnings**

**Liver Warning:** this product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen.

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

■ to make a child sleepy

■ in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

■ if your child has is allergic to any of the ingredients in this product

### **Ask a doctor before use if your child has**

■ liver disease ■ high blood pressure ■ heart disease ■ diabetes ■ thyroid disease ■ glaucoma ■ a breathing problem such as chronic bronchitis ■ persistent or chronic cough such as occurs with asthma ■ cough that occurs with too much phlegm (mucus)

### **Ask a doctor or pharmacist before use if your child is**

■ taking the blood thinning drug warfarin ■ taking sedatives or tranquilizers

### **When using this product**

■ **do not exceed recommended dosage (see overdose warning)**

■ excitability may occur, especially in children

■ marked drowsiness may occur

■ sedatives and tranquilizers may increase the drowsiness effect

### **Stop use and ask a doctor if**

■ nervousness, dizziness, or sleeplessness occur ■ pain, nasal congestion or cough gets worse or lasts more than 5 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with fever, rash, or persistent headache. These could be signs of a serious condition.

### **Keep out of reach of children**

### **Overdose section**

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical even if you do not notice any signs or symptoms

### ***Directions***

■ **this product does not contain directions or complete warning for adult use**

■ **do not give more than directed (see overdose warning)**

■ find right dosage on chart below

■ empty each Single-Use Vial of all liquid when dosing, following the instructions under

"Easy to use" on the side panel

- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- children under the age of 6, do not use

### **Dosing Chart**

---

<b>Age (Years)</b>	<b>Dose (mL)</b>
6 - 11 years	1 Single use vial (10mL) every 4 hours. Do not give more than 5 times in 24 hours

---

### ***Inactive ingredients***

citric acid, edetate disodium, glycerin, maltitol, mixed berry flavor, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium metabisulfite, sorbitol, sucralose, trisodium citrate, xanthan gum.

### ***Other information***

- each Single -Use vial contains 11mg of sodium
- store at 20-25°C (68-77°F)
- tamper-evident: do not use if the top flap of the box is open. Do not use if any vial is open or broken.

### **Questions or comments**

Call 941-202-3207 or visit [www.drkids.com](http://www.drkids.com)

### ***Package***

**Drug Facts** (continued)

Redness or swelling is present. New symptoms occur. Cough comes back, or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.

**Keep out of reach of children.**  
**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

**Directions**  
 This product does not contain directions or complete warnings for adult use. Do not give more than directed (see Overdose warning). First right dosage on chart below. Empty each Single-Use Vial required of all liquid when dosing, following the instructions under "Easy to Use" on the side panel. Repeat dose every 4 hours while symptoms last. Do not give more than 5 doses in 24 hours. Children under the age of 6 do not use.

**Dosing Chart**

Age (Years)	Dose (mL)
6-11 years	1 Single use vial (10mL) every 4 hours. Do not give more than 5 times in 24 hours

**Other Information**  
 Each 10 mL vial contains 6 mg sodium. Store between 20-25°C (68-77°F). Tamper evident: do not use if top flap of carton is open. Do not use if any vials are open or broken.

**Inactive Ingredients**  
 citric acid, edetate disodium, glycerin, mixed berry flavor, propylene glycol, purified water, sodium benzoate, sodium metabisulfite, sorbitol, sucralose, titanium dioxide.

**Questions or Comments?**  
 Call 941-202-3207 or visit www.drkids.com

**Made in the USA:**  
 Distributed by CalmCo LLC  
 Sarasota, FL 34240

**DrKids®** Ages 6-11 years

**Children's Multi-Symptom Cold & Flu**

Acetaminophen 320 mg / Pain Reliever, Fever Reducer  
 Chlorpheniramine maleate 2 mg / Antihistamine  
 Dextromethorphan HBr 10 mg / Cough Suppressant  
 Phenylephrine HCl 5 mg / Nasal Decongestant

**Relief of:**

- Coughs
- Sneezing & itchy, watery eyes
- Stuffy & runny nose
- Fever, headaches & body aches

**Easy to Use**

- Tap the vial to eliminate air bubbles
- Twist the top of the vial off
- Squeeze the liquid into child's mouth

**Why DrKids:**  
 Meet DrKids, the safe, clean medicine in a single-use dose. We are parents, too, and our revolutionary vials remove the guesswork of over or under dosing. DrKids is hygienic (no more sticky cups or syringes) and portable, making it safe and easy to administer wherever you are. DrKids is ready so you can be ready at home or on-the-go. Go forth and conquer!

**Free from:**

- Alcohol
- Dyes
- Ibuprofen & Aspirin
- Sugar

**Drug Facts** Keep Outer Package for Complete Product Information

**Active Ingredient (in each 10 mL)**

Active Ingredient	Purpose
Acetaminophen 320 mg	Pain Reliever/Fever Reducer
Chlorpheniramine maleate 2 mg	Antihistamine
Dextromethorphan HBr 10 mg	Cough Suppressant
Phenylephrine HCl 5 mg	Nasal Decongestant

**Uses**  
 Temporarily relieves the following cold/flu symptoms: minor aches and pains, headache, sore throat, cough, stuffy nose, sneezing and runny nose, temporarily reduces fever.

**Warnings**  
**Liver Warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes:  
 more than 5 doses in 24 hours, which is the maximum daily amount  
 with other drugs containing acetaminophen  
**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include: skin redness, blisters, rash  
 If a skin reaction occurs, stop use and seek medical help right away.  
**Severe throat warning:** If a sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**  
 with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
 to make a child sleepy.  
 in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.  
 if your child is allergic to any of the ingredients in this product.

**Ask a doctor before use if your child has**  
 liver disease, high blood pressure, heart disease, diabetes, thyroid disease, epilepsy, breathing problem such as chronic bronchitis, persistent or chronic cough such as occurs with asthma, cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if your child is** taking the blood thinning drug warfarin, taking sedatives or tranquilizers

**When using this product** Do not exceed recommended dose (see overdose warning). Excitability may occur, especially in children. Marked drowsiness may occur. Sedatives and tranquilizers may increase drowsiness.

**Stop use and ask a doctor if** nervousness, dizziness or sleeplessness occur, pain, nasal congestion or cough gets worse or lasts more than 5 days, fever gets worse or lasts more than 5 days.

*Drug Facts Continued to Side Panel*

# DRKIDS CHILDREN MULTI SYMPTOM COLD AND FLU

acetaminophen 320mg, chlorpheniramine maleate 2mg, dextromethorphan hbr 10mg, phenylephrine hcl 5mg solution

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:73282-0113
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	320 mg in 10 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1W5297W6MV)	PHENYLEPHRINE	5 mg in 10 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 10 mL
<b>CHLORPHENIRAMINE MALEATE</b> (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg in 10 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>MALTITOL</b> (UNII: D65DG142WK)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>SORBITOL SOLUTION</b> (UNII: 8KW3E207O2)	
<b>SODIUM METABISULFITE</b> (UNII: 4VON5FNS3C)	

<b>WATER</b> (UNII: 059QF0KO0R)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>TRISODIUM CITRATE DIHYDRATE</b> (UNII: B22547B95K)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	

### Product Characteristics

<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	BERRY (MIXED BERRY FLAVOR)	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73282-0113-6	16 in 1 CARTON	02/15/2024	
1		10 mL in 1 VIAL; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	02/15/2024	

**Labeler** - Calmco (079308308)

Revised: 12/2023

Calmco