

RANITIDINE- ranitidine tablet
Shopko Stores Operating Co., LLC

Drug Facts

ACTIVE INGREDIENT (IN EACH TABLET)

Ranitidine 150 mg (as ranitidine hydrochloride USP, 168 mg)

PURPOSE

Acid reducer

USES

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

WARNINGS

Allergy alert: Do not use if you are allergic to ranitidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers
- if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding

Ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

DIRECTIONS

- adults and children 12 years and over:
 - to **relieve** symptoms, swallow 1 tablet with a glass of water
 - to **prevent** symptoms, swallow 1 tablet with a glass of water **30 to 60 minutes before** eating food or drinking beverages that cause heartburn
 - can be used up to twice daily (do not take more than 2 tablets in 24 hours)
- children under 12 years: ask a doctor

OTHER INFORMATION

- **TAMPER EVIDENT: DO NOT USE IF THE CARTON OR PRINTED FOIL UNDER CAP IS OPEN OR TORN.**
- store at 20° - 25° C (68° - 77° F)
- avoid excessive heat or humidity
- this product is sodium and sugar free

INACTIVE INGREDIENTS

Colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, talc, titanium dioxide

QUESTIONS?

Call **1-800-406-7984**

PRINCIPAL DISPLAY PANEL

SHOPKO®

Compare to the Active Ingredient of Zantac 150®

Maximum Strength

Acid Reducer

Ranitidine Tablets, USP 150 mg

Acid Reducer

Prevents & Relieves: Heartburn Associated with Acid Indigestion & Sour Stomach

50 Tablets

Packaged by: Ohm Laboratories Inc.

5093015/0212

SHOPKO®

Maximum Strength
Acid Reducer

Ranitidine Tablets, USP 150 mg

Acid Reducer

SHOPKO®

[†]Compare to the Active
Ingredient of Zantac 150®

Maximum Strength
Acid Reducer

Ranitidine Tablets, USP 150 mg

Acid Reducer

*Prevents & Relieves: Heartburn Associated
with Acid Indigestion & Sour Stomach*



50 Tablets

Read the directions, consumer information leaflet and warnings before use.
Keep the carton. It contains important information.

Expiration Date:

NON VARNISH

Batch No.:

[†]This product is not manufactured or distributed by Boehringer Ingelheim
Pharmaceuticals Inc., owner of the registered trademark Zantac 150®.



By Return or Exchange

Packaged by:
Olm Laboratories Inc.
1385 Livingston Avenue
North Brunswick, NJ 08902
MADE IN INDIA

©2012 Shopko Stores Operating Co., LLC
www.shopko.com

0212



Drug Facts (continued)

- to prevent symptoms, swallow 1 tablet with a glass of water **30 to 60 minutes before eating**
- food or drinking beverages that cause heartburn
- can be used up to twice daily (do not take more than 2 tablets in 24 hours)
- children under 12 years: ask a doctor

Other information

- TAMPER EVIDENT: DO NOT USE IF THE CARTON OR PRINTED FOIL UNDER CAP IS OPEN OR TORN.**
- avoid excessive heat or humidity
- store at 20° - 25° C (68° - 77° F)
- this product is sodium and sugar free

Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, talc, titanium dioxide

Questions? call 1-800-406-7984

Warnings

- Allergy alert:** Do not use if you are allergic to ranitidine or other acid reducers
- Do not use**
 - If you have trouble or pain or pain or vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor
 - with other acid reducers
 - If you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- frequent chest pain
- unexplained weight loss
- nausea or vomiting
- stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222).


Directions


- adults and children 12 years and over:
 - to relieve symptoms, swallow 1 tablet with a glass of water

5093015



5093015





RANITIDINE

ranitidine tablet

Product Information

| | | | |
|-------------------------|----------------|--------------------|----------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:370 12-351 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|-------------------|----------|
| RANITIDINE HYDRO CHLORIDE (UNII: BK76465IHM) (RANITIDINE - UNII:884KT10 YB7) | RANITIDINE | 150 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|--|----------|
| COLLOIDAL SILICON DIOXIDE (UNII: ETJ7Z6XBU4) | |
| CROSCARMELOSE SODIUM (UNII: M28OL1HH48) | |
| HYPROMELLOSES (UNII: 3NXW29V3WO) | |
| FERRIC OXIDE RED (UNII: 1K09F3G675) | |
| MAGNESIUM STEARATE (UNII: 70097M6I30) | |
| CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U) | |
| POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A) | |
| TALC (UNII: 7SEV7J4R1U) | |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP) | |

Product Characteristics

| | | | |
|----------|----------------------------------|--------------|----------|
| Color | white (Off-White to Pale Yellow) | Score | no score |
| Shape | OVAL | Size | 7mm |
| Flavor | | Imprint Code | 9R |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|-------------------|---------------------|----------------------|--------------------|
| 1 | NDC:370 12-351-50 | 50 in 1 BOTTLE | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| | | | |

| | | | |
|------|------------|------------|--|
| ANDA | ANDA200536 | 03/30/2012 | |
|------|------------|------------|--|

Labeler - Shopko Stores Operating Co., LLC (023252638)

Registrant - Ranbaxy Pharmaceuticals Inc. (937890044)

Establishment

| Name | Address | ID/FEI | Business Operations |
|--------------------------------|---------|-----------|------------------------|
| Shasun Pharmaceuticals Limited | | 915786829 | manufacture(37012-351) |

Revised: 2/2013

Shopko Stores Operating Co., LLC