

**LOW DOSE ASPIRIN- aspirin tablet, delayed release**  
**Family Dollar Services Inc**

-----  
**Family Wellness 44-600A-Low Dose**

***Active ingredient (in each tablet)***

Aspirin 81 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever

***Uses***

for the temporary relief of minor aches and pains or as recommended by your doctor.  
**Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.**

***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take more or for a longer time than directed
- take a blood thinning (anticoagulant) or steroid drug
- are age 60 or older
- have had stomach ulcers or bleeding problems
- have 3 or more alcoholic drinks every day while using this product
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]

**Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if**

- you have asthma
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug for

- gout
- diabetes
- arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - vomit blood
  - have bloody or black stools
  - feel faint
  - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

***Directions***

- **do not take more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

***Other information***

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

- see end flap for expiration date and lot number

### ***Inactive ingredients***

corn starch, D&C yellow #10, FD&C yellow #6, hypromellose, methacrylic acid, microcrystalline cellulose, polydextrose, polyethylene glycol, shellac wax, silica, simethicone, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, triethyl citrate

### ***Questions or comments?***

**1-800-426-9391**

### ***Principal Display Panel***

**FAMILY**  
**Wellness™**

**COMPARE TO THE ACTIVE INGREDIENT IN**  
**BAYER® LOW DOSE ASPIRIN†**

LOW DOSE  
**ASPIRIN 81 mg**  
**Pain Reliever (NSAID)**

Safety Coated  
Aspirin Regimen  
Enteric Coated

**TALK TO YOUR DOCTOR OR OTHER HEALTHCARE PROVIDER**  
**BEFORE USING THIS PRODUCT FOR YOUR HEART.**

**120 TABLETS**

**ACTUAL SIZE**

**NDC 55319-601-32**

**TAMPER EVIDENT: DO NOT USE IF**  
**IMPRINTED SAFETY SEAL UNDER**  
**CAP IS BROKEN OR MISSING**

†This product is not manufactured or distributed by Bayer AG,  
owner of the registered trademark Bayer® Low Dose Aspirin.  
50844      ORG012260032

**DISTRIBUTED BY: MIDWOOD BRANDS LLC,**  
500 VOLVO PKWY, CHESAPEAKE, VA 23320 USA


**NOT 100% SATISFIED?**

Return within 30 days to the store of purchase  
for a refund (with receipt) or exchange.

No print/No varnish  
Lot & Exp date

SKU 913724

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**



0 32251 92399 9

**Drug Facts (continued)**  
Questions or comments?  
1-800-426-9391

**FAMILY Wellness™**

COMPARE TO THE ACTIVE INGREDIENT IN BAYER® LOW DOSE ASPIRIN



LOW DOSE  
**ASPIRIN 81 mg**  
Pain Reliever (NSAID)  
Safety Coated Aspirin Regimen

Enteric Coated

TALK TO YOUR DOCTOR OR OTHER HEALTHCARE PROVIDER BEFORE USING THIS PRODUCT FOR YOUR HEART.

**120 TABLETS**

NDC 55319-601-32

ACTUAL SIZE

B-0500-600A-32-N  
OR6012260032

**Drug Facts (continued)**

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- do not take more than directed
- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

**Other information**

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

**Inactive ingredients** corn starch, D&C yellow #10, FD&C yellow #6, hypromellose, methacrylic acid, microcrystalline cellulose, polydextrose, polyethylene glycol, shellac wax, silica, simethicone, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, triethyl citrate

**Drug Facts (continued)**  
Questions or comments?  
1-800-426-9391

**Family Wellness 44-600A**

**Drug Facts**

**Active ingredient (in each tablet)**  
Aspirin 81 mg (NSAID)\*  
\*nonsteroidal anti-inflammatory drug

**Purpose**  
Pain reliever

**Uses**  
for the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.

**Warnings**  
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:  

- hives
- facial swelling
- shock
- asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you  

- take more or for a longer time than directed
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

**KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION**

**Drug Facts**

**Active ingredient (in each tablet)**  
Aspirin 81 mg (NSAID)\*  
\*nonsteroidal anti-inflammatory drug

**Purpose**  
Pain reliever

**Uses**  
for the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.

**Warnings**  
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:  

- hives
- facial swelling
- shock
- asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you  

- take more or for a longer time than directed
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

**Drug Facts (continued)**  
Questions or comments?  
1-800-426-9391

B-0500-600A-32-N  
OR6012260032

**Drug Facts (continued)**

■ have had stomach ulcers or bleeding problems

■ take a blood thinning (anticoagulant) or steroid drug

■ are age 60 or older

■ have 3 or more alcoholic drinks every day while using this product

**Do not use** ■ if you are allergic to aspirin or any other pain reliever/fever reducer

■ if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if** ■ you have asthma

■ stomach bleeding warning applies to you

■ you have a history of stomach problems, such as heartburn

■ you are taking a diuretic

■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

**Ask a doctor or pharmacist before use if you are** taking a prescription drug for

■ gout ■ diabetes ■ arthritis

**Stop use and ask a doctor if** ■ an allergic reaction occurs. Seek medical help right away.

■ you experience any of the following signs of stomach bleeding: ■ vomit blood

■ have bloody or black stools

■ feel faint

■ have stomach pain that does not get better

■ ringing in the ears or a loss of hearing occurs

■ pain gets worse or lasts more than 10 days

■ fever gets worse or lasts more than 3 days

■ redness or swelling is present

■ new symptoms occur

These could be signs of a serious condition.

**Drug Facts (continued)**  
Questions or comments?  
1-800-426-9391

B-0500-600A-32-N  
OR6012260032

**Drug Facts (continued)**

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- do not take more than directed
- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

**Other information**

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

**Inactive ingredients** corn starch, D&C yellow #10, FD&C yellow #6, hypromellose, methacrylic acid, microcrystalline cellulose, polydextrose, polyethylene glycol, shellac wax, silica, simethicone, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, triethyl citrate

**Drug Facts (continued)**  
Questions or comments?  
1-800-426-9391

<b>LOW DOSE ASPIRIN</b>			
aspirin tablet, delayed release			
<b>Product Information</b>			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:55319-601
<b>Route of Administration</b>	ORAL		
<b>Active Ingredient/Active Moiety</b>			

Ingredient Name	Basis of Strength	Strength
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>METHACRYLIC ACID</b> (UNII: 1CS02G8656)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYDEXTROSE</b> (UNII: VH2XOU12IE)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>SHELLAC</b> (UNII: 46N107B71O)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM BICARBONATE</b> (UNII: 8MDF5V39QO)	
<b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>TRIACETIN</b> (UNII: XHX3C3X673)	
<b>TRIETHYL CITRATE</b> (UNII: 8Z96QXD6UM)	

### Product Characteristics

<b>Color</b>	yellow	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	6mm
<b>Flavor</b>		<b>Imprint Code</b>	L
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55319-601-32	1 in 1 CARTON	11/19/2019	
1		120 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:55319-601-17	1 in 1 CARTON	11/19/2019	02/10/2022
2		300 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	11/19/2019	

**Labeler** - Family Dollar Services Inc (024472631)

**Establishment**

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(55319-601)

**Establishment**

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(55319-601)

**Establishment**

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(55319-601) , pack(55319-601)

**Establishment**

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(55319-601)

Revised: 1/2025

Family Dollar Services Inc