

**ANTI-DIARRHEAL, CIRCLE K- loperamide hcl tablet tablet  
Lil' Drug Store Products, Inc.**

-----  
**Anti-Diarrheal, Circle K**

***Drug Facts***

***Active ingredient (in each caplet)***

Loperamide HCl, USP 2 mg

***Purpose***

Anti-diarrheal

***Use***

Controls symptoms of diarrhea, including Travelers' Diarrhea

***Warnings***

**Allergy alert**

**Allergy alert:** Do not use if you have ever had a rash or other allergic reaction to Loperamide HCl.

**Heart alert**

**Heart alert:** Taking more than directed can cause serious heart problems or death.

**Do not use** if you have bloody or black stool

**Ask a doctor before use if you have**

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

**Ask a doctor or pharmacist before use if you are** taking a prescription drug. Loperamide may interact with certain prescription drugs.

**When using this product** tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

**Stop use and ask a doctor if**

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

**Directions**

■ **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**

■ find right dose on chart. If possible, use weight to dose; otherwise use age.

---

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hour
children 9 - 11 years (60 - 95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6 - 8 years (48 - 59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2 - 5 years (34 - 47 lbs)	ask a doctor
children under 2 (up to 33 lbs)	do not use

---

**Other information**

**Other information**

■ store between 68-77°F (20-25°C)

**Inactive ingredients**

**Inactive ingredients**

anhydrous lactose, croscarmellose sodium, crospovidone, D&C Yellow no. 10 aluminum lake, FD&C Blue no. 1 aluminum lake, hydrogenated vegetable oil, magnesium stearate, powdered cellulose, pregelatinized starch

**Questions or comments?**

**Questions or comments?**

call toll-free **1-877-507-6516 (M-F 8AM - 4:30PM CST)**

**PDP [Circle K Anti-Diarrheal, 2mg Loperamide]**

**CONTROLS THE SYMPTOMS OF DIARRHEA**

Compare to the active ingredient in

**Imodium® A-D\***

**CIRCLE K™**

**antidiarrheal  
Loperamide HCl  
Tablets, USP  
2 mg**

[caplet image]

actual size

**4 caplets\*\* (2 per pouch)**

**Drug Facts (continued)**

**Inactive ingredients**  
anhydrous lactose, croscarmellose sodium, crospovidone, D&C Yellow no. 10 aluminum lake, FD&C Blue no. 1 aluminum lake, hydrogenated vegetable oil, magnesium stearate, powdered cellulose, pregelatinized starch

**Questions or comments?**  
call toll-free 1-877-507-6516 (M-F 8AM - 4:30PM CST)

CONTROLS THE SYMPTOMS OF DIARRHEA  
Compare to the active ingredient in  
**Imodium® A-D\***



**4 caplets\*\* (2 per pouch)**

DO NOT USE IF INNER POUCH OR CARTON IS OPEN OR DAMAGED!

**Drug Facts**

Active ingredient (in each caplet)	Purpose
Loperamide HCl, USP 2 mg	Anti-diarrheal

**Use** Controls symptoms of diarrhea, including Travelers' Diarrhea

**Warnings**

**Allergy alert:** Do not use if you have ever had a rash or other allergic reaction to Loperamide HCl.

**Heart alert:** Taking more than directed can cause serious heart problems or death.

Do not use if you have bloody or black stool

**Ask a doctor before use if you have**

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

**Ask a doctor or pharmacist before use if you are taking a prescription drug.** Loperamide may interact with certain prescription drugs.

**When using this product** tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

**Stop use and ask a doctor if**

- symptoms get worse
- you get abdominal swelling or bulging. These may be signs of a serious condition.
- diarrhea lasts for more than 2 days

LIFT PANEL FOR CONTINUED DRUG FACTS

Product manufactured for:  
Ul' Drug Store Products, Inc., 8900 Earhart Lane SW,  
Cedar Rapids, IA 52404 www.jldrugstore.com 5292C-US-05-19-3

Proudly distributed by Circle K Stores Inc.

\*This product is not manufactured or distributed by JOHNSON & JOHNSON CONSUMER INC, McNeil Consumer Healthcare Division, distributor of Imodium® A-D.

Each caplet (\*\*capsule-shaped tablet) contains Loperamide HCl, USP 2 mg

**Drug Facts (continued)**

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- drink plenty of clear fluids to help prevent dehydration caused by diarrhea
- find right dose on chart. If possible, use weight to dose; otherwise use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9 - 11 years (60 - 95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6 - 8 years (48 - 59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2 - 5 years (34 - 47 lbs)	ask a doctor
children under 2 (up to 33 lbs)	do not use

**Other information**

- store between 68-77°F (20-25°C)

**ANTI-DIARRHEAL, CIRCLE K**  
loperamide hcl tablet tablet

<b>Product Information</b>			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:66715-5529
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>LOPERAMIDE HYDROCHLORIDE</b> (UNII: 77T135393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg

**Inactive Ingredients**

Ingredient Name	Strength
<b>CROSCARMELLOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>CROSPVIDONE (15 MPA.S AT 5%)</b> (UNII: 68401960MK)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>HYDROGENATED COTTONSEED OIL</b> (UNII: Z82Y2C65EA)	
<b>ANHYDROUS LACTOSE</b> (UNII: 3SY5LH9PMK)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>POWDERED CELLULOSE</b> (UNII: SMD1X3XO9M)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	

**Product Characteristics**

<b>Color</b>	green	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	9mm
<b>Flavor</b>		<b>Imprint Code</b>	123
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66715-5529-2	2 in 1 CARTON	03/26/2021	
1		2 in 1 POUCH; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074091	03/26/2021	

**Labeler** - Lil' Drug Store Products, Inc. (093103646)

Revised: 10/2022

Lil' Drug Store Products, Inc.