

**CEBASHIN CAP.- forsythia suspensa fruit, lonicera japonica whole, mentha arvensis top, platycodon grandiflorus root, licorice, arctium lappa fruit, glycine max whole, lophatherum gracile top, schizonepeta tenuifolia spike, saiga tatarica horn capsule
Chunwoo Pharmaceutical Co., Ltd.**

Disclaimer: This homeopathic product has not been evaluated by the Food and Drug Administration for safety or efficacy. FDA is not aware of scientific evidence to support homeopathy as effective.

Cebashin Cap.

Active ingredients

FORSYTHIA SUSPENS A FRUIT }
LONICERA JAPONICA WHOLE }
MENTHA ARVENSIS TOP }
PLATYCODON GRANDIFLORUS ROOT }
LICORICE }
ARCTIUM LAPPA FRUIT } ----- cold / cough aid
GLYCINE MAX WHOLE }
LOPHATHERUM GRACILE TOP }
SCHIZONEPETA TENUFOLIA SPIKE }
SAIGA TATARICA HORN }

Purpose

Cold/cough aid

Directions

For adults, take 1 packet per intake, 3 times a day, before or between meals

Inactive ingredients

Corn Starch, Animal originated ingredients: Lactose Hydrate/Cow Milk

Keep out of reach of children

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Warnings

Do not use if you are

- Children under 5 years old

Do not use if you have

- Galactose intolerance
- Lapp lactase deficiency
- Glucose-galactose malabsorption (this medicine contains lactose)

Ask a doctor before use if you

- have hypertension
- have a heart disease or a kidney disease
- a patient with edema
- pregnant or have a chance to pregnant
- a patient who has rashes, bloodshot, redness, itching, and etc. caused by medication
- a patient with significantly weakened stomachs
- an elderly person (in general, elderly people should take with care, such as reducing their dosage, as their physiological function has deteriorated)
- in a post-disease stage (an abnormal reaction can be easily seen and the symptoms can worsen)
- a patient who is being treated by a doctor or are taking other medications

Stop use and ask a doctor, pharmacist or health professionals if

- if there is no improvement in symptoms after taking it for a few days
- you have following symptoms after use:

- Pseudohyperaldosteronism: Reduced urine volume, swollen face, swollen hands and feet, heavy eyelids, firm hands, increased blood pressure, headache, and etc.

(Medicines with a maximum daily dose of 1g or more may develop hypoglycemia, blood pressure rise, low sodium body fluid, edema, weight gain, and etc. caused by pseudohyperaldosteronism if taken continuously for a long time. Therefore, make sufficient observations (measurement of serum potassium levels) and stop taking if there is any abnormality.)

-Myopathy: Myopathy (muscle disease) may occur as a result of hypokalemia, so take sufficient observation and stop taking if you find any abnormalities such as feeling helplessness, limb cramps, paralysis, and etc.

-Skin trouble: a rash, itch, bloodshot, and etc.

-Digestive system disorder: anorexia, stomach discomfort, nausea, vomiting, and etc.

Other Warnings

- do not take more than directed
- if children are to take this medicine, take it under the supervision of their guardian.
- when taken with potassium-containing agents, licorice-containing agents, glycyrrhizic acid or other related agents, loop diuretics (furosemide, ethacrynic acid), or thiazide diuretics (trichlormethiazide) and myopathy, it is likely to appear due to gastric aldosterone (pseudohyperaldosteronism) or hypokalemia
- If taken with other herbal medicines, be careful of overlapping of the crude medicines.

- In principle, you should not to take it continuously for a long time, but if it is necessary, consult with a doctor, herb doctor, dentist, pharmacist, or other health care professionals.

Uses

a sore throat, thirst, cough, headache from a cold

Cebashin Cap.

Cebashin

Cap.

- **sore throat**
- **bronchitis**
- **tonsillitis**

Cebashin Cap.
10 capsule / 615mg each

Homeopathic medicine

Drug Facts (continued)	Drug Facts (continued)
Inactive ingredients Animal originated ingredients: Lactose Hydrate/Cow Milk, Corn Starch	Questions or Comments? Contact us at 82-2-6263-1001 and chunwopharm@naver.com

This product has not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.

Drug Facts	Drug Facts (continued)
<p>Active ingredients</p> <p>Lonicera Japonica Whole (710 mg) Forsythia Suspensa Fruit (710 mg) Mentha Arvensis Top (426 mg) Licorice (426 mg) Platycodon Grandiflorus Root (426 mg) Glycine Max Whole (356 mg) Arctium Lappa Fruit (356 mg) Schizonepeta Tenuifolia Flowering Top (284 mg) Lophatherum Gracile Top (284 mg) Saiga Tatarica Horn (22 mg)</p> <p>Purpose</p> <p>---- cold/cough aid</p> <p>Uses Temporarily relieves symptoms of common cold: ■ sore throat ■ thirst ■ cough ■ headache</p> <p>Warnings Do not use if you have ■ galactose intolerance ■ lapp lactase deficiency ■ glucose-galactose malabsorption (contains lactose)</p> <p>Ask a doctor before use if you have ■ hypertension ■ a heart disease ■ kidney disease ■ edema ■ rashes ■ bloodshot ■ redness ■ itching ■ weakened stomachs</p> <p>Ask a doctor before use if you are • an elderly person (in general, elderly people should take with care, such as reducing their dosage, as their physiological function has deteriorated) • in a post-disease stage (an abnormal reaction can be easily seen and the symptoms can worsen) • a patient who is being treated by a doctor or are taking other medications</p> <p>Stop use and ask a doctor, pharmacist or health professionals if • there is no improvement of symptoms after taking it for a few days</p>	<p>• you have following symptoms after use :</p> <p>- Pseudohyperaldosteronism : ■ Reduced urine volume ■ swollen face, hands and feet ■ heavy eyelids ■ firm hands ■ increased blood pressure ■ headache and etc.</p> <p>- Medicines with a maximum daily dose of 1g or more may develop : ■ hypoglycemia ■ blood pressure rise ■ low sodium body fluid ■ edema ■ weight gain and etc. caused by pseydohyperaldosteroni sm if taken continuously for a long time.</p> <p>- Make sufficient observations (measurement of serum potassium levels) and stop taking if there is any abnormality</p> <p>- Myopathy : Myopathy (muscle disease) may occur as a result of hypokalemia, so take sufficient observation and stop taking if you find any abnormalities such as: ■ feeling helplessness ■ limb cramps ■ paralysis and etc.</p> <p>- Skin trouble : ■ rash ■ itch ■ bloodshot and etc.</p> <p>- Digestive system disorder: ■ anorexia ■ stomach discomfort ■ nausea ■ vomiting and etc.</p> <p>Other warnings</p> <p>• do not take more than directed • if children are to take this medicine, take it under the supervision of their guardian • when taken with potassium-containing agents, licorice-containing agents, glycinic acid or other related agents, loop diuretics (furosemide, ethacrynic acid), or thiazide diuretics (trichlormethiazide) and myopathy, it is likely to appear due to gastric aldosterone (pseudohyperaldosteronism) or hypokalemia</p> <p>• If taken with other herbal medicines, be careful of overlapping of the crude medicines • In principle, you should not to take it continuously for a long time, but if it is necessary, consult with a doctor, herb doctor, dentist, pharmacist, or other health care professionals</p> <p>If pregnant or breast-feeding, ask a health professional before use</p> <p>Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.</p>

Drug Facts (continued)	Drug Facts (continued)
Directions ■ Take 1 packet per intake ■ 3 times a day ■ before or between meals	Other information • Store in a dry and cool place at 68-77°F (20-25°C) away from the direct sunlight • To avoid misuse, do not change container

Serial No. :
EXP :

Chunwoo Pharmaceutical Co.,Ltd.
17, Gaepo-ro 20-gil, Gangnam-gu, Seoul, Republic of Korea



- **sore throat**
- **bronchitis**
- **tonsillitis**

Cebashin
Cap.

CEBASHIN CAP.

forsythia suspensa fruit, lonicera japonica whole, mentha arvensis top, platycodon grandiflorus root, licorice, arctium lappa fruit, glycine max whole, lophatherum gracile top, schizonepeta tenuifolia

spike, saiga tatarica horn capsule

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:72850-0007
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FORSYTHIA SUSPENSIA FRUIT (UNII: P4793M1E55) (FORSYTHIA SUSPENSIA FRUIT - UNII:P4793M1E55)	FORSYTHIA SUSPENSIA FRUIT	710 mg in 10 g
LICORICE (UNII: 61ZBX54883) (LICORICE - UNII:61ZBX54883)	LICORICE	426 mg in 10 g
PLATYCODON GRANDIFLORUS ROOT (UNII: 2DF0NS002Z) (PLATYCODON GRANDIFLORUS ROOT - UNII:2DF0NS002Z)	PLATYCODON GRANDIFLORUS ROOT	426 mg in 10 g
ARCTIUM LAPPA FRUIT (UNII: EA541308MV) (ARCTIUM LAPPA FRUIT - UNII:EA541308MV)	ARCTIUM LAPPA FRUIT	356 mg in 10 g
LONICERA JAPONICA WHOLE (UNII: UV4X7MQQ43) (LONICERA JAPONICA WHOLE - UNII:UV4X7MQQ43)	LONICERA JAPONICA WHOLE	710 mg in 10 g
MENTHA ARVENSIS TOP (UNII: 8J5Y2F7A5X) (MENTHA ARVENSIS TOP - UNII:8J5Y2F7A5X)	MENTHA ARVENSIS TOP	426 mg in 10 g
GLYCINE MAX WHOLE (UNII: NID9NUV93F) (GLYCINE MAX WHOLE - UNII:NID9NUV93F)	GLYCINE MAX WHOLE	356 mg in 10 g
SCHIZONEPETA TENUFOLIA SPIKE (UNII: 2FN3BA1MZE) (SCHIZONEPETA TENUFOLIA SPIKE - UNII:2FN3BA1MZE)	SCHIZONEPETA TENUFOLIA SPIKE	284 mg in 10 g
LOPHATHERUM GRACILE TOP (UNII: OY50I2L657) (LOPHATHERUM GRACILE TOP - UNII:OY50I2L657)	LOPHATHERUM GRACILE TOP	284 mg in 10 g
SAIGA TATARICA HORN (UNII: PU683NDA7H) (SAIGA TATARICA HORN - UNII:PU683NDA7H)	SAIGA TATARICA HORN	22 mg in 10 g

Inactive Ingredients

Ingredient Name	Strength
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	

Product Characteristics

Color	red, white	Score	no score
Shape	CAPSULE	Size	21mm
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72850-0007-2	1 in 1 BOX	01/12/2022	
1	NDC:72850-0007-1	10 g in 1 CAPSULE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved homeopathic		01/12/2022	

Labeler - Chunwoo Pharmaceutical Co., Ltd. (688729065)

Registrant - Chunwoo Pharmaceutical Co., Ltd. (688729065)

Establishment

Name	Address	ID/FEI	Business Operations
Chunwoo Pharmaceutical Co., Ltd.		688729065	manufacture(72850-0007)

Revised: 1/2022

Chunwoo Pharmaceutical Co., Ltd.