

**BACTIMICINA CHILDRENS COUGH AND COLD- chlorpheniramine maleate, dextromethorphan hydrobromide liquid
DLC Laboratories, Inc.**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Bactimicina Liquid Children's Cough and Cold

Drug Facts

Active ingredients (in each 5 mL = 1 teaspoon)	Purposes
Chlorpheniramine maleate, USP 1 mg	Antihistamine
Dextromethorphan HBr, USP 7.5 mg	Cough suppressant

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 - runny nose
 - itchy, watery eyes
 - sneezing
 - itching of the nose or throat

Warnings

Do not use

- to sedate a child or to make a child sleepy
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- trouble urinating due to an enlarged prostate gland
- glaucoma
- a cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.

When using this product

- **do not use more than directed**
- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than 4 doses in any 24-hour period
- mL=milliliter; tsp=teaspoonful

Age (years)	Dose
Under 6	do not use
6-11	2 teaspoons (10 mL) every 6 hours
12 and over	4 teaspoons (20 mL) every 6 hours

Other information

- **each teaspoon (5 mL) contains:** sodium 1 mg
- store at 15-30°C (59-86°F)
- measure only with dosage cup provided
- **do not use if printed bottle wrap is missing or broken.**

Inactive ingredients

citric acid, FD&C red no. 40, flavor, glycerin, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose

Questions

1-800-858-3889

Manufactured by:
DLC Laboratories, Inc.
Paramount, CA 90723 USA

PRINCIPAL DISPLAY PANEL - 118 mL Bottle Label

Long Acting TRUSTED SINCE 1978

Children's

Bactimicina

Multi-Symptoms • Liquid Ages 6+ Years

Cough & Cold

Dextromethorphan HBr (Cough Suppressant)

Chlorpheniramine Maleate (Antihistamine)

Relieves Cough up to 8hrs • Sneezing • Runny Nose

Alcohol Free

Natural Strawberry Flavor

4 FL OZ (118 mL)

Do not use if printed bottle wrap is missing or broken.

Drug Facts

Active ingredients Purposes
(in each 5 mL = 1 teaspoon)
Chlorpheniramine maleate, USP 1 mg... Antihistamine
Dextromethorphan HBr, USP 7.5 mg... Cough suppressant

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

Warnings

Do not use

- to sedate a child or to make a child sleepy
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping

Long Acting

TRUSTED SINCE 1970

Children's
Bactimicina

Multi-Symptom • Liquid

Ages **6+** Years

Cough & Cold

Dextromethorphan HBr (Cough Suppressant)
Chlorpheniramine Maleate (Antihistamine)

Relieves Cough up to **8 hrs** • Sneezing • Runny Nose

Alcohol Free



Natural
Strawberry Flavor

4 FL OZ (118 mL)

Drug Facts (continued)

the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- trouble urinating due to an enlarged prostate gland
- glaucoma
- a cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.

Manufactured by / Fabricado por:
DLC Laboratories, Inc.
Paramount, CA 90723 USA
Questions / Preguntas:
1-800-858-3889
dlclabs.com

© 2022 DLC Laboratories, Inc.

@delacruzproducts

@DLClaboratories

Read Here
Levante Aquí

Drug Facts
Datos Medicinales

P0173-GYY

Drug Facts (continued)

When using this product

- do not use more than directed
- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than 4 doses in any 24-hour period
- mL = milliliter, tsp = teaspoonful

Age (years)

Age (years)	Dose
Under 6	do not use
6-11	2 teaspoons (10 mL) every 6 hours
12 and over	4 teaspoons (20 mL) every 6 hours

Other information

- each teaspoon (5 mL) contains: sodium 1 mg
- store at 15-30°C (59-86°F)
- measure only with dosage cup provided

Inactive ingredients citric acid, FD&C red no. 40, flavor, glycerin, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose

Datos Medicinales

Ingredientes activos Propósitos

(en cada 5 mL = 1 cucharadita)
Maleato de clorfeniramina, USP 1 mg... Antihistamínico
Dextrometorfano HBr, USP 7.5 mg... Antitusivo

Usos alivia temporalmente la tos causada por una irritación leve de garganta y bronquios, como puede ocurrir por un resfriado

alivia temporalmente estos síntomas de rinitis alérgica o algunas alergias de las vías respiratorias superiores

goteo nasal

picañón de ojos, ojos llorosos

estornudos

comezón nasal o de garganta

Advertencias

No usar

- para sedar a un niño o hacerlo sentir soñoliento

P0173-GYY

Datos Medicinales (continuación)

- si está tomando un medicamento recetado inhibidor de la monoaminoxidasa (MAOI) (ciertos medicamentos para la depresión, afecciones psiquiátricas o emocionales, o enfermedad de Parkinson), o en un período de 2 semanas después de haber suspendido el uso del medicamento MAOI. Si no sabe si su receta contiene MAOI, pregúntele a un médico o farmacéutico antes de usar este producto.

Pregúntele a un médico antes de usar si usted tiene

- problemas al orinar debido al agrandamiento de la glándula prostática
- glaucoma
- tos que ocurre con demasiada flema (mucosidad)
- problemas respiratorios o los crónicos que perdura o se produce por fumar, asma, bronquitis crónica o enfisema

Consulte con un médico o farmacéutico antes de usarlo si usted está tomando sedantes o tranquilizantes.

Cuando use este producto

- no exceda la dosis recomendada
- puede provocar mucha somnolencia
- evite tomar alcohol
- el alcohol, los sedantes o los tranquilizantes pueden incrementar la somnolencia
- tenga cuidado si conduce u opera maquinaria
- puede provocar excitabilidad, especialmente en niños.

Deje de usar y consulte con un médico si

- la tos dura más de 7 días, regresa o está acompañada por fiebre, sarpullido o dolor de cabeza persistente. Estos podrían ser síntomas de una afección grave.

Si está embarazada o lactando, consulte con un profesional de la salud antes de usar. **Manténgase fuera del alcance de los niños.** En caso de sobredosis, pida ayuda médica o contacte con un Centro de Control de Envenenamientos inmediatamente.

Instrucciones

- no tomar más de 4 dosis en un período de 24 horas
- mL = mililitro

Edad Dosis

Edad	Dosis
Menor de 6	no lo use
6-11	2 cucharaditas (10 mL) cada 6 horas
12 y mayores	4 cucharaditas (20 mL) cada 6 horas

Otra información

- cada cucharadita (5 mL) contiene: 1 mg de sodio
- conservese a 15-30°C (59-86°F)
- use únicamente vaso dosificador incluido

Ingredientes inactivos ácido cítrico, FD&C rojo no. 40, saborizante, glicerina, propilenglicol, agua purificada, benzoato de sodio, sorbitol, sucralosa

P0173-GYY

BACTIMICINA CHILDRENS COUGH AND COLD

chlorpheniramine maleate, dextromethorphan hydrobromide liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:24286-1551
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	1 mg in 5 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	7.5 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
GLYCERIN (UNII: PDC6A3C0OX)	

Product Characteristics

Color	red (Red)	Score	
Shape		Size	
Flavor	STRAWBERRY (Natural Strawberry flavor)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:24286-1551-4	1 in 1 BOX	10/23/2009	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
--------------------	--	----------------------	--------------------

OTC monograph final	part341	10/23/2009	
---------------------	---------	------------	--

Labeler - DLC Laboratories, Inc. (093351930)

Establishment

Name	Address	ID/FEI	Business Operations
DLC Laboratories, Inc.		093351930	manufacture(24286-1551) , label(24286-1551)

Revised: 11/2022

DLC Laboratories, Inc.