

PRO-DEN RX- sodium fluoride rinse
Den-mat Holdings, Llc

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, [click here](#).

For topical application to aid in the protection against dental caries. Neutral pH is especially safe for crowns and restorations.

Dosage and Administration

Dispense ½ oz. (approx. 1 pump) of Pro-DenRx 2.0% Neutral Sodium Fluoride Solution into the provided mixing cup. Instruct patient to rinse vigorously for 30 seconds with ½ oz. of the solution around and between teeth, then expectorate. For maximum benefit repeat the rinse procedure with an additional ½ oz. of solution. Pro-DenRx 2.0% Neutral Sodium Fluoride Solution may also be applied full strength, with cotton pledgets, to teeth isolated with cotton rolls.

Recommended Frequency

Do not exceed four (4) treatments per year.

Contraindications

Hypersensitivity to fluoride.

Warnings and Precautions

For Professional Office Use Only. This product is not intended for home or unsupervised consumer use. Do not swallow. Keep out of reach of children. Not recommended for children under the age of 6. Limited to topical use in the mouth only.

Adverse Reactions

The following adverse reactions are possible in individuals hypersensitive to fluoride: eczema, atopic dermatitis, urticaria, gastric distress, headache, and weakness.

Caution

Federal law prohibits dispensing without a prescription.

Overdosage

Accidental ingestion of large amounts of fluoride can cause: nausea, vomiting, abdominal pain, diarrhea, stupor and/or weakness (usually within 30 minutes). These symptoms may persist for 24 hours. If less than 5 mg fluoride/kg body weight (less than 23 mg fluoride/lb body weight) has been ingested, give calcium (milk) orally to relieve symptoms and observe for a few hours. If more than 5 mg fluoride/kg body weight (more than 23 mg fluoride/lb body weight) has been ingested, induce vomiting, give calcium (i.e., milk, 5% calcium gluconate or calcium lactate solution) and immediately seek medical assistance. For accidental ingestion of more than 15 mg fluoride/kg body weight (i.e., more than 6.9 mg fluoride/lb body weight) induce vomiting, transport and admit immediately to a hospital facility.

Ingredients

Water, Sodium Fluoride, PEG-40 Hydrogenated Castor Oil, Sodium Benzoate, Disodium Phosphate, Sodium Saccharin, Flavor, Methylparaben, Sodium Phosphate, Red #33.

How Supplied/Storage and Handling

2.0% Neutral Sodium Fluoride (0.9% Fluoride Ion) oral solution supplied in a plastic bottle with child-resistant closure containing 64 fl. oz. (1.89 L). Store at room temperature. Protect from freezing. Do not store in direct sunlight.

Rx Only

Revised: 01/2015

1-800-433-6628

Reorder Number: 2037RBDT

Manufactured for

Den-Mat Holdings, LLC

1017 W. Central Ave.

Lompoc, CA 93436

©2015 Den-Mat Holdings, LLC. All rights reserved. 001394900 06/15SN

Principal Display Panel - 64 fl. oz. Bottle Label

NDC 59883-912-64

treatment rinse

2.0% neutral sodium fluoride

berry fresh flavor

IMPORTANT:

Read directions for proper use.

Net Wt. 64 fl. oz. (1.89 L)

Indications and Usage
For topical application to aid in the protection against dental caries. Pro-Den Rx is especially safe for children and adolescents.

Dosage and Administration
Dispense 1/2 oz. (approx. 1 pump) of Pro-Den Rx 2.0% Neutral Sodium Fluoride Solution into the provided rinsing cup. Instruct patient to rinse vigorously for 30 seconds with 1/2 oz. of the solution several times before each meal. Then expel into the sink. For maximum benefit repeat this rinse procedure with an additional 1/2 oz. of solution. Pro-Den Rx 2.0% Neutral Sodium Fluoride Solution may also be applied full strength, with cotton swabs, to teeth associated with orthodontics.

Recommended Frequency
Do not exceed four (4) treatments per year.

Contraindications
Hypersensitivity to fluoride.

Warnings and Precautions
For Professional Office Use Only. This product is not intended for home or unsupervised consumer use. Do not swallow. Keep out of reach of children. Not recommended for children under the age of 6. Limited to topical use in the mouth only.

Adverse Reactions
The following adverse reactions are possible in individuals hypersensitive to fluoride: eczema, allergic dermatitis, urticaria, gothic ulcers, hives, and asthma.

Cautions
Federal law prohibits dispensing without a prescription.

Overdosage
Accidental ingestion of large amounts of fluoride can cause nausea, vomiting, abdominal pain, diarrhea, slight muscle weakness (usually within 30 minutes). These symptoms may persist for 24 hours. If less than 0.5 mg fluoride/kg body weight (less than 35 mg fluoride/kg body weight) has been ingested, give sodium (Na+) orally to relieve symptoms and observe for 48 hours. If more than 0.5 mg fluoride/kg body weight (more than 35 mg fluoride/kg body weight) has been ingested, induce vomiting, give sodium (i.e., milk, 5% calcium gluconate or calcium lactate solution) and immediately seek medical assistance. For accidental ingestion of more than 10 mg fluoride/kg body weight (i.e., more than 6 mg fluoride/kg body weight), induce vomiting, transport and admit immediately to a hospital facility.

Ingredients
Water, Sodium Fluoride, PEG-40 Hydrogenated Castor Oil, Sodium Benzoate, Disodium Phosphate, Sodium Saccharin, Flavor, Methylparaben, Sodium Phosphate, Red #33.

How Supplied/Storage and Handling
2.0% Neutral Sodium Fluoride (0.9% Fluoride Ion) oral solution applied in a plastic bottle with child-resistant closure containing 64 fl. oz. (1.89 L). Store at room temperature. Protect from freezing. Do not store in direct sunlight.

Rx Only
Revised: 01/2015

1-800-433-6628
Reorder Number: 2037RBDT

dm
DenMat
Manufactured for
Den-Mat Holdings, LLC
1017 W. Central Ave.
Lompoc, CA 93436
©2015 Den-Mat Holdings, LLC. All rights reserved. 001394900 06/15SN

Net Wt. 64 fl. oz. (1.89 L)

PRO-DEN RX

sodium fluoride rinse

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:59883-912
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SODIUM FLUORIDE (UNII: 8ZYQ1474W7) (FLUORIDE ION - UNII:Q80VPU408O)	FLUORIDE ION	0.9 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0K00R)	
POLYOXYL 40 HYDROGENATED CASTOR OIL (UNII: 7YC686GQ8F)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM PHOSPHATE, DIBASIC, ANHYDROUS (UNII: 22ADO53M6F)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
SODIUM PHOSPHATE (UNII: SE337SVY37)	
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	BERRY (BERRY)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:59883-912-64	1890 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/21/2008	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		10/21/2008	

Labeler - Den-mat Holdings, Llc (809857704)