

**BASE LABORATORIES CORNEAL FLOW HYPERTONICITY OPHTHALMIC EYE-
sodium chloride ointment
Joonem LLC**

Base Laboratories Corneal Flow Hypertonicity Ophthalmic Eye Ointment

Drug Facts

Active ingredient

Sodium chloride 3%

Purpose

Hypertonicity agent

Uses:

Temporary relief of corneal edema

Warnings:

For external use only

Do not use

Except under the advice and supervision of a doctor

When using this product

- It may cause temporary burning and irritation
- Replace cap after use
- To avoid contamination, do not touch top of container to any surface

Stop use and ask a doctor if

- Condition worsens or persists for more than 72 hours
- You experience eye pain, changes in vision, continued redness or irritation of the eye

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Pull down the lower lid of the affected eye
- Apply a small amount (1/4 inch) of ointment to the inside of eyelid
- Apply every 3 to 4 hours or as directed by a doctor

Other information

- Store at 15-25C (59-77F)
- Keep tightly closed
- Do not Freeze
- Do not use if difficult to dispense or visible particles are seen in the product
- Side effects associated with this product may be reported to the email address below


Inactive ingredients

Water, Glycerin, Propylene Glycol, Squalane, Sodium Hyaluronate, Dimethicone, Tocopheryl Acetate, Carnosine, Butylene Glycol, Phenoxyethanol, Chlorphenesin, Ubiquinone

Questions or comments?

cs@baselaboratories.com

Package Labeling:

| | | | | | | | | | | |
|--|--|---|--|--|--|--|---|--|--|---|
|  | <p>scan to learn more To learn more about our products, ingredients, directions and suggestions please visit: www.baselaboratories.com</p>  | <table border="1"> <tr> <td data-bbox="941 871 1226 976"> Drug Facts Active ingredient Purpose Sodium chloride (3%).....Hypertonicity agent Uses: Temporary relief of corneal edema </td> <td data-bbox="1226 871 1520 976"> Directions • Pull down the lower lid of the affected eye • Apply a small amount (1/4 inch) of ointment to the inside of eyelid • Apply every 3 or 4 hours or as directed by a doctor </td> </tr> <tr> <td data-bbox="941 976 1226 1123"> Warnings: For external use only Do not use Except under the advice and supervision of a doctor When using this product • It may cause temporary burning and irritation • Replace cap after use • To avoid contamination, do not touch top of container to any surface </td> <td data-bbox="1226 976 1520 1123"> Other information • Store at 15-25C (59-77F) • Keep tightly closed • Do not Freeze • Do not use if difficult to dispense or visible particles are seen in the product • Side effects associated with this product may be reported to the email address below </td> </tr> <tr> <td data-bbox="941 1123 1226 1207"> Stop use and ask a doctor if • Condition worsens or persists for more than 72 hours • You experience eye pain, changes in vision, continued redness or irritation of the eye </td> <td data-bbox="1226 1123 1520 1207"> Inactive ingredients Water, Glycerin, Propylene Glycol, Squalane, Sodium Hyaluronate, Dimethicone, Tocopheryl Acetate, Carnosine, Butylene Glycol, Phenoxyethanol, Chlorphenesin, Ubiquinone </td> </tr> <tr> <td data-bbox="941 1207 1226 1260"> Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. </td> <td data-bbox="1226 1207 1520 1260"> Questions or comments? cs@baselaboratories.com </td> </tr> </table> <p>Extra water in the eye can cause a host of eye related problems such as hazy and poor vision, eye discomfort and tenderness when you touch your eye</p> <p>Base Laboratories Corneal Flow Eye Ointment promotes the drawing out of extra water from your swollen cornea offering corneal edema relief</p> | Drug Facts Active ingredient Purpose Sodium chloride (3%).....Hypertonicity agent Uses: Temporary relief of corneal edema | Directions • Pull down the lower lid of the affected eye • Apply a small amount (1/4 inch) of ointment to the inside of eyelid • Apply every 3 or 4 hours or as directed by a doctor | Warnings: For external use only Do not use Except under the advice and supervision of a doctor When using this product • It may cause temporary burning and irritation • Replace cap after use • To avoid contamination, do not touch top of container to any surface | Other information • Store at 15-25C (59-77F) • Keep tightly closed • Do not Freeze • Do not use if difficult to dispense or visible particles are seen in the product • Side effects associated with this product may be reported to the email address below | Stop use and ask a doctor if • Condition worsens or persists for more than 72 hours • You experience eye pain, changes in vision, continued redness or irritation of the eye | Inactive ingredients Water, Glycerin, Propylene Glycol, Squalane, Sodium Hyaluronate, Dimethicone, Tocopheryl Acetate, Carnosine, Butylene Glycol, Phenoxyethanol, Chlorphenesin, Ubiquinone | Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. | Questions or comments? cs@baselaboratories.com |
| Drug Facts Active ingredient Purpose Sodium chloride (3%).....Hypertonicity agent Uses: Temporary relief of corneal edema | Directions • Pull down the lower lid of the affected eye • Apply a small amount (1/4 inch) of ointment to the inside of eyelid • Apply every 3 or 4 hours or as directed by a doctor | | | | | | | | | |
| Warnings: For external use only Do not use Except under the advice and supervision of a doctor When using this product • It may cause temporary burning and irritation • Replace cap after use • To avoid contamination, do not touch top of container to any surface | Other information • Store at 15-25C (59-77F) • Keep tightly closed • Do not Freeze • Do not use if difficult to dispense or visible particles are seen in the product • Side effects associated with this product may be reported to the email address below | | | | | | | | | |
| Stop use and ask a doctor if • Condition worsens or persists for more than 72 hours • You experience eye pain, changes in vision, continued redness or irritation of the eye | Inactive ingredients Water, Glycerin, Propylene Glycol, Squalane, Sodium Hyaluronate, Dimethicone, Tocopheryl Acetate, Carnosine, Butylene Glycol, Phenoxyethanol, Chlorphenesin, Ubiquinone | | | | | | | | | |
| Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. | Questions or comments? cs@baselaboratories.com | | | | | | | | | |
| <h1>Corneal Flow</h1> <p>Eye Ointment with 3% Sodium Chloride Hypertonicity Ophthalmic Ointment</p> <p>For the Temporary Relief of Corneal Edema</p> <p>TRIPLE PACK 3 3 Tubes x 5g</p> <p>STERILE  NET WT 15g (1/2 Oz) TOTAL</p> | <p>scan to learn more To learn more about our products, ingredients, directions and suggestions please visit: www.baselaboratories.com</p>  | <p>KEY INGREDIENTS</p> <div style="display: flex; justify-content: space-around;"> <div data-bbox="1039 1480 1177 1564">  SODIUM CHLORIDE </div> <div data-bbox="1242 1480 1404 1564">  SODIUM HYALURONATE </div> </div>  <p>MADE IN CHINA Distributed by Jooneem LLC cs@baselaboratories.com 818-254-9459 www.baselaboratories.com</p> | | | | | | | | |

BASE LABORATORIES CORNEAL FLOW HYPERTONICITY OPTHALMIC EYE

sodium chloride ointment

Product Information

| | | | |
|---------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:80327-011 |
|---------------------|----------------|---------------------------|---------------|

Route of Administration TOPICAL

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|-------------------|--------------|
| SODIUM CHLORIDE (UNII: 451W47IQ8X) (CHLORIDE ION - UNII:Q32ZN48698) | SODIUM CHLORIDE | 30 mg in 1 g |

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| WATER (UNII: 059QF0KO0R) | |
| GLYCERIN (UNII: PDC6A3C0OX) | |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3) | |
| SQUALANE (UNII: GW89575KF9) | |
| HYALURONATE SODIUM (UNII: YSE9PPT4TH) | |
| DIMETHICONE (UNII: 92RU3N3Y1O) | |
| .ALPHA.-TOCOPHEROL ACETATE (UNII: 9E8X80D2L0) | |
| CARNOSINE (UNII: 8HO6PVN24W) | |
| BUTYLENE GLYCOL (UNII: 3XUS85K0RA) | |
| PHENOXYETHANOL (UNII: HIE492ZZ3T) | |
| CHLORPHENESIN (UNII: I670DAL4SZ) | |
| UBIDECARENONE (UNII: EJ27X76M46) | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:80327-011-00 | 3 in 1 CARTON | 01/01/2023 | |
| 1 | | 5 g in 1 TUBE; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M018 | 01/01/2023 | |

Labeler - Joonem LLC (117633878)