

**COPPERTONE COMPLETE SUNSCREEN SPF 50- avobenzone 3%, homosalate 9%, octisalate 4.5%, octocrylene 9% lotion**  
**Beiersdorf Inc**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Coppertone Complete Lotion SPF 50**

***Drug Facts***

***Active ingredients***

Avobenzone 3%, Homosalate 9%, Octisalate 4.5%, Octocrylene 9%

***Purpose***

Sunscreen

***Uses***

- helps prevent sunburn
- if used as directed with other sun protection measures (see Directions), decreases the risk of skin cancer and early skin aging caused by the sun

***Warnings***

**For external use only**

**Do not use** on damaged or broken skin

**When using this product** keep out of eyes. Rinse with water to remove.

**Stop use and ask a doctor if** rash occurs

**Keep out of reach of children.** If product is swallowed, get medical help or contact a Poison Control Center right away.

***Directions***

- apply liberally 15 minutes before sun exposure
- reapply:
  - after 80 minutes of swimming or sweating
  - immediately after towel drying
  - at least every 2 hours

■ **Sun Protection Measures.** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including:

- limit time in the sun, especially from 10 a.m. – 2 p.m.
- wear long-sleeve shirts, pants, hats, and sunglasses
- children under 6 months: Ask a doctor

### ***Other information***

- protect this product from excessive heat and direct sun
- may stain or damage some fabrics or surfaces

### ***Inactive ingredients***

water, SD alcohol 40-B, butyloctyl salicylate, diethylhexyl carbonate, glycereth-26, styrene/acrylates copolymer, butylene glycol, diisopropyl adipate, dimethicone, polyamide-8, isododecane, 1,2-hexanediol, ammonium acryloyldimethyltaurate/beheneth-25 methacrylate crosspolymer, glycerin, hydroxyacetophenone, tocopherol, ethylhexyl methoxycrylene, arachidyl alcohol, fragrance, xanthan gum, behenyl alcohol, arachidyl glucoside, disodium EDTA, sodium ascorbyl phosphate

### ***Questions?***

1-866-288-3330

Coppertone® Sunscreen Lotion

Complete 50

Proven UVA/UVB Protection

+ Moisturizes All Day\*

+ Feels Lightweight

Free of Oxybenzone, Octinoxate, Dyes & PABA

Water Resistant (80 Minutes)

Broad Spectrum SPF 50

# Coppertone<sup>®</sup>

SUNSCREEN LOTION



## COMPLETE 50

Proven UVA/UVB Protection

- + Moisturizes All Day\*
- + Feels Lightweight

Free of Oxybenzone,  
Octinoxate, Dyes & PABA

Water Resistant (80 Minutes)  
Broad Spectrum SPF 50

7 FL OZ (207 mL)

Broad Spectrum UVA/UVB protection that stops up to 98% of damaging UV rays, feels lightweight - good for the whole family.

### Drug Facts

#### Active ingredients

Avobenzene 3%, Homosalate 9%,  
Octisalate 4.5%, Octocrylene 9% ..... Sunscreen

#### Purpose

#### Use

- helps prevent sunburn

#### Warnings

For external use only

**Do not use** on damaged or broken skin

**When using this product** keep out of eyes. Rinse with water to remove.

**Stop use and ask a doctor if** rash occurs

**Keep out of reach of children.** If product is swallowed, get medical help or contact a Poison Control Center right away.

#### Directions

- apply liberally 15 minutes before sun exposure
- reapply:
  - after 80 minutes of swimming or sweating
  - immediately after towel drying
  - at least every 2 hours
- **Sun Protection Measures.** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including:
  - limit time in the sun, especially from 10 a.m. – 2 p.m.
  - wear long-sleeve shirts, pants, hats, and sunglasses
- children under 6 months: Ask a doctor

#### Other information

- protect this product from excessive heat and direct sun
- may stain or damage some fabrics or surfaces

**Inactive ingredients** water, SD alcohol 40-B, butyloctyl salicylate, diethylhexyl carbonate, glycereth-26, styrene/acrylates copolymer, butylene glycol, diisopropyl adipate, dimethicone, polyamide-8, isododecane, 1,2-hexanediol, ammonium acryloyldimethyltaurate/beheneth-25 methacrylate crosspolymer, glycerin, hydroxyacetophenone, tocopherol, ethylhexyl methoxycrylene, arachidyl alcohol, fragrance, xanthan gum, behenyl alcohol, arachidyl glucoside, disodium EDTA, sodium ascorbyl phosphate

**Questions?** 1-866-288-3330

\*For sun protection, use as directed.

www.coppertone.com

**Beiersdorf**

Beiersdorf Inc. Stamford, CT 06901  
©=reg. TM. of Beiersdorf AG,  
Germany



C00021034

## COPPERTONE COMPLETE SUNSCREEN SPF 50

avobenzene 3%, homosalate 9%, octisalate 4.5%, octocrylene 9% lotion

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:66800-0030
<b>Route of Administration</b>	TOPICAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>AVOBENZONE</b> (UNII: G63QQF2NOX) (AVOBENZONE - UNII:G63QQF2NOX)	AVOBENZONE	3 g in 100 g
<b>OCTOCRYLENE</b> (UNII: 5A68WGF6WM) (OCTOCRYLENE - UNII:5A68WGF6WM)	OCTOCRYLENE	9 g in 100 g
<b>HOMOSALATE</b> (UNII: V06SV4M95S) (HOMOSALATE - UNII:V06SV4M95S)	HOMOSALATE	9 g in 100 g
<b>OCTISALATE</b> (UNII: 4X49Y0596W) (OCTISALATE - UNII:4X49Y0596W)	OCTISALATE	4.5 g in 100 g

**Inactive Ingredients**

Ingredient Name	Strength
<b>ISODODECANE</b> (UNII: A8289P68Y2)	
<b>1,2-HEXANEDIOL</b> (UNII: TR046Y3K1G)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>HYDROXYACETOPHENONE</b> (UNII: G1L3HT4CMH)	
<b>ARACHIDYL ALCOHOL</b> (UNII: 1QR1QRA9BU)	
<b>ARACHIDYL GLUCOSIDE</b> (UNII: 6JVV35JOOJ)	
<b>SODIUM ASCORBYL PHOSPHATE</b> (UNII: 836SJG51DR)	
<b>BUTYLOCTYL SALICYLATE</b> (UNII: 2EH13UN8D3)	
<b>AMMONIUM ACRYLOYLDIMETHYLTAURATE/BEHENETH-25 METHACRYLATE CROSSPOLYMER (52000 MPA.S)</b> (UNII: LZ291VH90H)	
<b>ETHYLHEXYL METHOXYCRYLENE</b> (UNII: S3KFG6Q5X8)	
<b>BUTYLENE GLYCOL</b> (UNII: 3XUS85K0RA)	
<b>DOCOSANOL</b> (UNII: 9G1OE216XY)	
<b>DISODIUM EDTA-COPPER</b> (UNII: 6V475AX06U)	
<b>STYRENE/ACRYLAMIDE COPOLYMER (MW 500000)</b> (UNII: 5Z4DPO246A)	
<b>TOCOPHEROL</b> (UNII: R0ZB2556P8)	
<b>GLYCERETH-26</b> (UNII: NNE56F2N14)	
<b>ALCOHOL</b> (UNII: 3K9958V90M)	
<b>DIISOPROPYL ADIPATE</b> (UNII: P7E6YFV72X)	
<b>FRAGRANCE FRESH CITRUS FLORAL ORC1501495</b> (UNII: OU4GI2R2WB)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>POLYAMIDE-8 (4500 MW)</b> (UNII: 77723GV81A)	
<b>DIETHYLHEXYL CARBONATE</b> (UNII: YCD5000Z6L)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O)	

**Product Characteristics**

<b>Color</b>	white (white to off-white)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66800-0030-7	207 g in 1 TUBE; Type 0: Not a Combination Product	11/02/2021	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part352	11/02/2021	

**Labeler** - Beiersdorf Inc (001177906)

Revised: 3/2023

Beiersdorf Inc