

**BACK AND BODY EXTRA STRENGTH- aspirin, caffeine tablet, film coated  
DOLGENCORP, LLC**

-----  
**Dollar General 44-413**

***Active ingredients (in each caplet)***

Aspirin 500 mg (NSAID)\*

Caffeine 32.5 mg

\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever

Pain reliever aid

***Uses***

for the temporary relief of

- backache pain
- muscle aches and pains
- minor aches and pains of arthritis

***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- are age 60 or older
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma

### **Ask a doctor or pharmacist before use if you are**

taking a prescription drug for

- gout
- diabetes
- arthritis

### **Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - have bloody or black stools
  - vomit blood
  - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

### ***Directions***

- **do not take more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 caplets every 6 hours, not to exceed 8 caplets in 24 hours
- children under 12 years: do not use unless directed by a doctor

**Other information**

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

**Inactive ingredients**

corn starch, hypromellose, maltodextrin, microcrystalline cellulose, polyethylene glycol, sodium lauryl sulfate

**Questions or comments?**

**1-888-309-9030**

**Principal display panel**

**DG™ | health**

†Compare to the active ingredients of Bayer® Back & Body

**Extra Strength  
Back & Body**

Aspirin (**NSAID**)  
and Caffeine

**Pain Reliever/Pain Reliever Aid**

**50 Caplets**

Actual Caplet Size

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED  
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

†This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Bayer® Back & Body. 50844 REV0222B41315

DISTRIBUTED BY OLD EAST MAIN CO.  
100 MISSION RIDGE  
GOODLETTSVILLE, TN 37072

**100%  
Satisfaction  
Guaranteed!**  
(888-309-9030)

NO PRINT/NO VARNISH  
LOT NO. & EXPIRE AREA

508441315

A1061

TAMPER EVIDENT: DO NOT USE IF IMPRINTED  
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

**DG<sup>TM</sup> health**

**Extra Strength  
Back & Body**  
Aspirin (NSAID)  
and Caffeine  
**50 Caplets**  
Pain Reliever/Pain Reliever Aid

Actual Caplet Size

†Compare to the active ingredients of Bayer® Back & Body

**DG<sup>TM</sup> health**

**Extra Strength  
Back & Body**  
Aspirin (NSAID) and Caffeine  
Pain Reliever/Pain Reliever Aid

B-0315-413-15  
REV0222B41315

**Drug Facts (continued)**  
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  

- do not take more than directed
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 caplets every 6 hours, not to exceed 8 caplets in 24 hours
- children under 12 years: do not use unless directed by a doctor

**Other information**  

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

**Inactive ingredients** corn starch, hypromellose, maltodextrin, microcrystalline cellulose, polyethylene glycol, sodium lauryl sulfate

**Questions or comments?** 1-888-309-9030

\*This product is not manufactured or distributed by Bayer AG, owner of the registered trademarks Bayer® Back & Body. 50844 REV0222B41315

DISTRIBUTED BY OLD EAST MAIN CO.  
100 MISSION RIDGE  
GOODLETTSVILLE, TN 37072

**100% Satisfaction Guaranteed!**  
(888) 309-9030

NO PRINT  
NO GLUE AREA

**Drug Facts (continued)**  
Do not use:  

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if:**  

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma

**Ask a doctor or pharmacist before use if you are:**  

- taking a prescription drug for gout
- diabetic
- arthritis

**Stop use and ask a doctor if:**  

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:  
  - feel faint
  - have bloody or black stools
  - vomit blood
- have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

NO PRINT  
NO GLUE AREA

**Dollar General 44-413**

<b>BACK AND BODY EXTRA STRENGTH</b> aspirin, caffeine tablet, film coated			
<b>Product Information</b>			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:55910-134
<b>Route of Administration</b>	ORAL		
<b>Active Ingredient/Active Moiety</b>			

Ingredient Name	Basis of Strength	Strength
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	500 mg
<b>CAFFEINE</b> (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	32.5 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>MALTODEXTRIN</b> (UNII: 7CVR7L4A2D)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)	

### Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	17mm
<b>Flavor</b>		<b>Imprint Code</b>	44;413
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55910-134-15	1 in 1 CARTON	04/12/2004	
1		50 in 1 BOTTLE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	04/12/2004	

**Labeler** - DOLGENCORP, LLC (068331990)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(55910-134)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(55910-134)

### Establishment

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
LNK International, Inc.		832867894	manufacture(55910-134)

## **Establishment**

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
LNK International, Inc.		967626305	pack(55910-134)

## **Establishment**

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
LNK International, Inc.		117025878	manufacture(55910-134)

Revised: 5/2024

DOLGENCORP, LLC