

NAPROXEN SODIUM - naproxen sodium tablet, film coated
Aurohealth LLC

Naproxen Sodium Tablets, USP 220 mg

Drug Facts

Active ingredient (in each tablet)

Naproxen sodium USP 220 mg (naproxen 200 mg) (NSAID)*

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis
- muscular aches
- menstrual cramps
- toothache
- backache
- headache
- the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing

- weakness in one part or side of body
- slurred speech
- leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- drink a full glass of water with each dose

<p>Adults and children 12 years and older</p>	<ul style="list-style-type: none"> • take 1 tablet every 8 to 12 hours while symptoms last • for the first dose you may take 2 tablets within the first hour • do not exceed 2 tablets in any 8 to 12 hour period • do not exceed 3 tablets in a 24 hour period
<p>Children under 12 years</p>	<p>ask a doctor</p>

Other information

- **each tablet contains:** sodium 20 mg
- store at 20° to 25°C (68° to 77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Inactive ingredients

colloidal silicon dioxide, FD&C blue #2 aluminum lake, hypromellose, magnesium

stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, and titanium dioxide

Questions or comments?

1-855-274-4122

Distributed by: **AUROHEALTH LLC**
2572 Brunswick Pike,
Lawrenceville, NJ 08648
Made in India

Code No.: TS/DRUGS/22/2009

PACKAGE LABEL-PRINCIPAL DISPLAY PANEL - 220 mg (100 Tablets, Container Label)

Healthy Living™
NDC 58602-845-21
Naproxen Sodium
Tablets, USP
220 mg (NSAID)

- ***Pain reliever***
- ***Fever reducer***

STRENGTH TO LAST
12 HOURS

K
100 Tablets

Healthy Living™
 NDC 58602-845-21
**Naproxen Sodium
 Tablets, USP
 220 mg (NSAID)**

• Pain reliever
 • Fever reducer

**STRENGTH TO LAST
 12 HOURS**

100 Tablets



Drug Facts

Active ingredient (in each tablet)
 Naproxen sodium USP 220 mg (naproxen 200 mg)
 (NSAID)*

Purposes
 Pain reliever/fever reducer
 *nonsteroidal anti-inflammatory drug

Uses temporarily relieves minor aches and pains due to: minor pain of arthritis muscular aches backache headache the common cold temporarily reduces fever menstrual cramps toothache

Warnings
Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives facial swelling asthma (wheezing) shock skin reddening rash blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older have had stomach ulcers or bleeding problems take a blood thinning (anticoagulant) or steroid drug take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) have 3 or more alcoholic drinks every day while using this product take more or for a longer time →

Do not use if seal over bottle opening is broken or missing.

Lift Here

Distributed by: **AUROHEALTH LLC**
 2572 Brunswick Pike,
 Lawrenceville, NJ 08648
 Made in India
 Code No.: TS/DRUGS/22/2009

LM-4177
 P1426308

*

NVZ

* Lot: XXXXXXXXX
 EXP: MM/YYYY
 Prefix & Variables of Lot, EXP shall be printed online during packing.

Drug Facts (continued)

than directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Do not use** if you have ever had an allergic reaction to any other pain reliever/fever reducer right before or after heart surgery. **Ask a doctor before use if** the stomach bleeding warning applies to you you have a history of stomach problems, such as heartburn you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke you are taking a diuretic you have problems or serious side effects from taking pain relievers or fever reducers. **Ask a doctor or pharmacist before use if you are** under a doctor's care for any serious condition taking any other drug. **When using this product** take with food or milk if stomach upset occurs. **Stop use and ask a doctor if** you experience any of the following signs of stomach bleeding: red faint vomit blood have bloody or black stools have stomach pain that does not get better you have symptoms of heart problems or stroke: chest pain trouble breathing weakness in one part or side of body slurred speech leg swelling pain gets worse or lasts more than 10 days fever gets worse or lasts more than 3 days you have difficulty swallowing it feels like the pill is stuck in your throat redness or swelling is present in the painful area any new symptoms appear. **If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions do not take more than directed the smallest effective dose should be used drink a full glass of water with each dose. **Adults and children 12 years and older:** take 1 tablet every 8 to 12 hours while symptoms last for the first dose you may take 2 tablets within the first hour do not exceed 2 tablets in any 8 to 12 hour period do not exceed 3 tablets in a 24 hour period **Children under 12 years:** ask a doctor. **Other information** each tablet contains: sodium 20 mg store at 20° to 25°C (68° to 77°F). Avoid high humidity and excessive heat above 40°C (104°F). **Inactive ingredients** colloidal silicon dioxide, FD&C blue #2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, and titanium dioxide.

Questions or comments? 1-855-274-4122

LM-4177 P1426308

PACKAGE LABEL-PRINCIPAL DISPLAY PANEL - 220 mg (100 Tablets Carton Label)

Healthy Living™

NDC 58602-845-21

**Compare to Aleve®

active ingredient

**Naproxen Sodium Tablets, USP
 220 mg (NSAID)**

- Pain reliever
- Fever reducer

K

STRENGTH TO LAST 12 HOURS

100 Tablets



NAPROXEN SODIUM

naproxen sodium tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58602-845
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:5Y7Y6R9ATQ)	NAPROXEN SODIUM	220 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	

HYPROMELLOSE 2910 (5 MPA.S) (UNII: R75537T0T4)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL 8000 (UNII: Q662QK8M3B)	
POVIDONE K30 (UNII: U725QWY32X)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	BLUE (light blue)	Score	no score
Shape	OVAL	Size	12mm
Flavor		Imprint Code	K;45
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58602-845-21	1 in 1 CARTON	05/16/2020	
1		100 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA205497	05/16/2020	

Labeler - Aurohealth LLC (078728447)

Establishment

Name	Address	ID/FEI	Business Operations
Aurobindo Pharma Limited		650381903	ANALYSIS(58602-845) , MANUFACTURE(58602-845)

Revised: 12/2023

Aurohealth LLC