

# GABAPENTIN- gabapentin tablet

## Epic Pharma, LLC

-----

### HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use GABAPENTIN TABLETS safely and effectively. See full prescribing information for GABAPENTIN TABLETS.

**GABAPENTIN tablets (once-daily), for oral use**  
**Initial U.S. Approval: 1993**

### INDICATIONS AND USAGE

Gabapentin tablets are indicated for the management of Postherpetic Neuralgia (PHN).

**Important Limitation: Once-daily gabapentin tablets are not interchangeable with other gabapentin products because of differing pharmacokinetic profiles that affect the frequency of administration**(See Warnings and Precautions)

### DOSAGE AND ADMINISTRATION

- Gabapentin tablets should be titrated to an 1,800 mg dose taken orally, once-daily, with the evening meal. Gabapentin tablets should be swallowed whole. Do not crush, split, or chew the tablets. (2.1)
- If gabapentin tablets dose is reduced, discontinued, or substituted with an alternative medication, this should be done gradually over a minimum of 1 week or longer (at the discretion of the prescriber). (2.1)
- Renal impairment: Dose should be adjusted in patients with reduced renal function. Gabapentin tablets should not be used in patients with CrCl less than 30 or in patients on hemodialysis. (2.2)

### DOSAGE FORMS AND STRENGTHS

300 and 600 mg tablets (3)

### CONTRAINDICATIONS

Gabapentin tablets are contraindicated in patients who have demonstrated hypersensitivity to the drug or its ingredients. (4)

### WARNINGS AND PRECAUTIONS

- Once-daily gabapentin tablets are not interchangeable with other gabapentin products.
- Antiepileptic drugs, including gabapentin, the active ingredient in gabapentin tablets, increase the risk of suicidal thoughts or behavior. (5.1)
- Respiratory depression may occur with gabapentin tablets when used with concomitant CNS depressants or in the setting of underlying respiratory impairment. Monitor patients and adjust dosage as appropriate. (5.2)
- Increased seizure frequency may occur in patients with seizure disorders if gabapentin tablets are rapidly discontinued. Withdraw gabapentin tablets gradually over a minimum of 1 week. (5.3)

### ADVERSE REACTIONS

The most common adverse reaction (greater than or equal to 5% and twice placebo) is dizziness. (6.1)

**To report SUSPECTED ADVERSE REACTIONS, contact Epic Pharma, LLC at 1-888-374-2791 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

### DRUG INTERACTIONS

- An increase in gabapentin AUC values have been reported when administered with hydrocodone. (7.6)
- An increase in gabapentin AUC values have been reported when administered with morphine. (7.7)
- An antacid containing aluminum hydroxide and magnesium hydroxide reduced the bioavailability of gabapentin immediate release by about approximately 20%, but by only 5% when gabapentin was taken 2 hours after antacids. It is recommended that gabapentin tablets be taken at least 2 hours following antacid administration. (7.10)

### USE IN SPECIFIC POPULATIONS

- Elderly: Reductions in gabapentin tablets dose should be made in patients with age-related compromised renal function. (8.5)
- Renal impairment: Dosage adjustment is necessary for patients with impaired renal function. (8.7)

**FULL PRESCRIBING INFORMATION: CONTENTS\***

**1 INDICATIONS AND USAGE**

**2 DOSAGE AND ADMINISTRATION**

- 2.1 Postherpetic Neuralgia
- 2.2 Patients with Renal Impairment

**3 DOSAGE FORMS AND STRENGTHS**

**4 CONTRAINDICATIONS**

**5 WARNINGS AND PRECAUTIONS**

- 5.1 Suicidal Behavior and Ideation
- 5.2 Respiratory Depression
- 5.3 Withdrawal of Gabapentin
- 5.4 Tumorigenic Potential
- 5.5 Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS)/Multiorgan Hypersensitivity
- 5.6 Laboratory Tests

**6 ADVERSE REACTIONS**

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing and Other Experience with other Formulations of Gabapentin

**7 DRUG INTERACTIONS**

- 7.1 Phenytoin
- 7.2 Carbamazepine
- 7.3 Valproic Acid
- 7.4 Phenobarbital
- 7.5 Naproxen
- 7.6 Hydrocodone
- 7.7 Morphine
- 7.8 Cimetidine
- 7.9 Oral Contraceptives
- 7.10 Antacid (containing aluminum hydroxide and magnesium hydroxide)
- 7.11 Probenecid
- 7.12 Drug/Laboratory Test Interactions

**8 USE IN SPECIFIC POPULATIONS**

- 8.1 Pregnancy
- 8.2 Lactation
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Hepatic Impairment
- 8.7 Renal Impairment

**9 DRUG ABUSE AND DEPENDENCE**

**10 OVERDOSAGE**

**11 DESCRIPTION**

**12 CLINICAL PHARMACOLOGY**

- 12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

12.4 Special Populations

### **13 NONCLINICAL TOXICOLOGY**

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

### **14 CLINICAL STUDIES**

### **16 HOW SUPPLIED/STORAGE AND HANDLING**

### **17 PATIENT COUNSELING INFORMATION**

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## **FULL PRESCRIBING INFORMATION**

### **1 INDICATIONS AND USAGE**

Gabapentin tablets are indicated for the management of postherpetic neuralgia.

Once-daily gabapentin tablets are not interchangeable with other gabapentin products because of differing pharmacokinetic profiles that affect the frequency of administration.

### **2 DOSAGE AND ADMINISTRATION**

#### **2.1 Postherpetic Neuralgia**

Do not use once-daily gabapentin tablets interchangeably with other gabapentin products.

Titrate gabapentin tablets to an 1,800 mg dose taken orally once daily with the evening meal. Gabapentin tablets should be swallowed whole. Do not split, crush, or chew the tablets.

If gabapentin tablets dosing is reduced, discontinued, or substituted with an alternative medication, this should be done gradually over a minimum of one week or longer (at the discretion of the prescriber).

In adults with postherpetic neuralgia, gabapentin tablets therapy should be initiated and titrated as follows:

**Table 1: Gabapentin Tablets Recommended Titration Schedule**

	<b>Day 1</b>	<b>Day 2</b>	<b>Days 3 to 6</b>	<b>Days 7 to 10</b>	<b>Days 11 to 14</b>	<b>Day 15</b>
Daily Dose	300 mg	600 mg	900 mg	1,200 mg	1,500 mg	1,800 mg

#### **2.2 Patients with Renal Impairment**

In patients with stable renal function, creatinine clearance ( $C_{Cr}$ ) can be reasonably well estimated using the equation of Cockcroft and Gault:

For females  $C_{Cr}=(0.85)(140-\text{age})(\text{weight})/[(72)(S_{Cr})]$

For males  $C_{Cr}=(140-\text{age})(\text{weight})/[(72)(S_{Cr})]$

where age is in years, weight is in kilograms and  $S_{Cr}$  is serum creatinine in mg/dL.

The dose of gabapentin tablets should be adjusted in patients with reduced renal function, according to Table 2. Patients with reduced renal function must initiate gabapentin tablets at a daily dose of 300 mg. Gabapentin tablets should be titrated following the schedule outlined in Table 1. Daily dosing in patients with reduced renal function must be individualized based on tolerability and desired clinical benefit.

**Table 2: Gabapentin Tablets Dosage Based on Renal Function**

<b>Once-daily dosing</b>	
Creatinine Clearance (mL/min)	Gabapentin Tablets Dose (once daily with evening meal)
≥ 60	1,800 mg
30 to 60	600 mg to 1,800 mg
< 30	Gabapentin tablets should not be administered
patients receiving hemodialysis	Gabapentin tablets should not be administered

### **3 DOSAGE FORMS AND STRENGTHS**

Tablets:

300 mg: white, oval-shaped, film-coated tablets debossed with “300” above “ ” on one side and plain on the other side.

600 mg: orange, oval-shaped, film-coated tablets debossed with “600” above “ ” on one side and plain on the other side.

### **4 CONTRAINDICATIONS**

Gabapentin tablets are contraindicated in patients with demonstrated hypersensitivity to the drug or their ingredients.

### **5 WARNINGS AND PRECAUTIONS**

Once-daily gabapentin tablets are not interchangeable with other gabapentin products because of differing pharmacokinetic profiles that affect the frequency of administration.

The safety and effectiveness of gabapentin tablets in patients with epilepsy has not been studied.

#### **5.1 Suicidal Behavior and Ideation**

Antiepileptic drugs (AEDs), including gabapentin, the active ingredient in gabapentin tablets, increase the risk of suicidal thoughts or behavior in patients taking these drugs for any indication. Patients treated with any AED for any indication should be monitored for the emergence or worsening of depression, suicidal thoughts or behavior, and/or any unusual changes in mood or behavior.

Pooled analyses of 199 placebo-controlled clinical trials (mono- and adjunctive therapy) of 11 different AEDs showed that patients randomized to one of the AEDs had approximately twice the risk (adjusted Relative Risk 1.8, 95% CI:1.2, 2.7) of suicidal thinking or behavior compared to patients randomized to placebo. In these trials, which had a median treatment duration of 12 weeks, the estimated incidence rate of suicidal behavior or ideation among 27,863 AED-treated patients was 0.43%, compared to 0.24% among 16,029 placebo-treated patients, representing an increase of approximately one case of suicidal thinking or behavior for every 530 patients treated. There were four suicides in drug-treated patients in the trials and none in placebo-treated patients, but the number is too small to allow any conclusion about drug effect on suicide.

The increased risk of suicidal thoughts or behavior with AEDs was observed as early as one week after starting drug treatment with AEDs and persisted for the duration of treatment assessed. Because most trials included in the analysis did not extend beyond 24 weeks, the risk of suicidal thoughts or behavior beyond 24 weeks could not be assessed.

The risk of suicidal thoughts or behavior was generally consistent among drugs in the data analyzed. The finding of increased risk with AEDs of varying mechanisms of action and across a range of indications suggests that the risk applies to all AEDs used for any indication. The risk did not vary substantially by age (5-100 years) in the clinical trials analyzed. Table 3 shows absolute and relative risk by indication for all evaluated AEDs.

**Table 3: Risk by Indication for Antiepileptic Drugs (including gabapentin, the active ingredient in gabapentin tablets) in the Pooled Analysis**

<b>Indication</b>	<b>Placebo Patients with Events Per 1,000 Patients</b>	<b>Drug Patients with Events Per 1,000 Patients</b>	<b>Relative Risk: Incidence of Events in Drug Patients/Incidence in Placebo Patients</b>	<b>Risk Difference: Additional Drug Patients with Events Per 1,000 Patients</b>
Epilepsy	1.0	3.4	3.5	2.4
Psychiatric	5.7	8.5	1.5	2.9
Other	1.0	1.8	1.9	0.9
Total	2.4	4.3	1.8	1.9

The relative risk for suicidal thoughts or behavior was higher in clinical trials for epilepsy than in clinical trials for psychiatric or other conditions, but the absolute risk differences were similar for the epilepsy and psychiatric indications.

Anyone considering prescribing gabapentin tablets must balance the risk of suicidal thoughts or behavior with the risk of untreated illness. Epilepsy and many other illnesses for which products containing active components that are AEDs (such as gabapentin, the active component in gabapentin tablets) are prescribed are themselves associated with morbidity and mortality and an increased risk of suicidal thoughts and behavior. Should suicidal thoughts and behavior emerge during treatment, the prescriber needs to consider whether the emergence of these symptoms in any given patient may be related to the illness being treated.

Patients, their caregivers, and families should be informed that gabapentin tablets

contain gabapentin which are also used to treat epilepsy and that AEDs increase the risk of suicidal thoughts and behavior and should be advised of the need to be alert for the emergence or worsening of the signs and symptoms of depression, any unusual changes in mood or behavior, or the emergence of suicidal thoughts, behavior, or thoughts about self-harm. Behaviors of concern should be reported immediately to healthcare providers.

## **5.2 Respiratory Depression**

There is evidence from case reports, human studies, and animal studies associating gabapentin with serious, life-threatening, or fatal respiratory depression when co-administrated with central nervous system (CNS) depressants, including opioids, or in the setting of underlying respiratory impairment. When the decision is made to co-prescribe gabapentin tablets with another CNS depressant, particularly an opioid, or to prescribe gabapentin tablets to patients with underlying respiratory impairment, monitor patients for symptoms of respiratory depression and sedation, and consider initiating gabapentin tablets at a low dose. The management of respiratory depression may include close observation, supportive measures, and reduction or withdrawal of CNS depressants (including gabapentin tablets).

## **5.3 Withdrawal of Gabapentin**

Gabapentin should be withdrawn gradually. If gabapentin tablets are discontinued, this should be done gradually over a minimum of 1 week or longer (at the discretion of the prescriber).

## **5.4 Tumorigenic Potential**

In standard preclinical *in vivo* lifetime carcinogenicity studies, an unexpectedly high incidence of pancreatic acinar adenocarcinomas was identified in male, but not female, rats. The clinical significance of this finding is unknown.

In clinical trials of gabapentin therapy in epilepsy comprising 2,085 patient-years of exposure in patients over 12 years of age, new tumors were reported in 10 patients, and pre-existing tumors worsened in 11 patients, during or within 2 years after discontinuing the drug. However, no similar patient population untreated with gabapentin was available to provide background tumor incidence and recurrence information for comparison. Therefore, the effect of gabapentin therapy on the incidence of new tumors in humans or on the worsening or recurrence of previously diagnosed tumors is unknown.

## **5.5 Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS)/Multiorgan Hypersensitivity**

Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS), also known as Multiorgan Hypersensitivity, has been reported in patients taking antiepileptic drugs, including gabapentin tablets. Some of these events have been fatal or life-threatening. DRESS typically, although not exclusively, presents with fever, rash, and/or lymphadenopathy in association with other organ system involvement, such as hepatitis, nephritis, hematological abnormalities, myocarditis, or myositis sometimes resembling an acute viral infection. Eosinophilia is often present. Because this disorder is variable in its expression, other organ systems not noted here may be involved.

It is important to note that early manifestations of hypersensitivity, such as fever or lymphadenopathy, may be present even though rash is not evident. If such signs or symptoms are present, the patient should be evaluated immediately. Gabapentin tablets should be discontinued if an alternative etiology for the signs or symptoms cannot be established.

## 5.6 Laboratory Tests

Clinical trial data do not indicate that routine monitoring of clinical laboratory procedures is necessary for the safe use of gabapentin tablets. The value of monitoring gabapentin blood concentrations has not been established.

## 6 ADVERSE REACTIONS

### 6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

A total of 359 patients with neuropathic pain associated with postherpetic neuralgia have received gabapentin tablets at doses up to 1,800 mg daily during placebo-controlled clinical studies. In clinical trials in patients with postherpetic neuralgia, 9.7% of the 359 patients treated with gabapentin tablets and 6.9% of 364 patients treated with placebo discontinued prematurely due to adverse reactions. In the gabapentin tablets treatment group, the most common reason for discontinuation due to adverse reactions was dizziness. Of gabapentin tablets-treated patients who experienced adverse reactions in clinical studies, the majority of those adverse reactions were either "mild" or "moderate".

Table 4 lists all adverse reactions, regardless of causality, occurring in at least 1% of patients with neuropathic pain associated with postherpetic neuralgia in the gabapentin tablets group for which the incidence was greater than in the placebo group.

**Table 4: Treatment-Emergent Adverse Reaction Incidence in Controlled Trials in Neuropathic Pain Associated with Postherpetic Neuralgia (Events in at Least 1% of all Gabapentin Tablets-Treated Patients and More Frequent Than in the Placebo Group)**

<b>Body System - Preferred Term</b>	<b>Gabapentin Tablets N = 359 %</b>	<b>Placebo N = 364 %</b>
<b>Ear and Labyrinth Disorders</b>		
Vertigo	1.4	0.5
<b>Gastrointestinal Disorders</b>		
Diarrhea	3.3	2.7
Dry mouth		
Constipation	2.8	1.4

Dyspepsia	1.4 1.4	0.3 0.8
<b>General Disorders</b>		
Peripheral edema Pain	3.9 1.1	0.3 0.5
<b>Infections and Infestations</b>		
Nasopharyngitis Urinary tract infection	2.5 1.7	2.2 0.5
<b>Investigations</b>		
Weight increased	1.9	0.5
<b>Musculoskeletal and Connective Tissue Disorders</b>		
Pain in extremity Back pain	1.9 1.7	0.5 1.1
<b>Nervous System Disorders</b>		
Dizziness Somnolence Headache Lethargy	10.9 4.5 4.2 1.1	2.2 2.7 4.1 0.3

In addition to the adverse reactions reported in Table 4 above, the following adverse reactions with an uncertain relationship to gabapentin tablets were reported during the clinical development for the treatment of postherpetic neuralgia. Events in more than 1% of patients but equally or more frequently in the gabapentin tablets-treated patients than in the placebo group included blood pressure increase, confusional state, gastroenteritis viral, herpes zoster, hypertension, joint swelling, memory impairment, nausea, pneumonia, pyrexia, rash, seasonal allergy, and upper respiratory infection.

## 6.2 Postmarketing and Other Experience with other Formulations of Gabapentin

In addition to the adverse experiences reported during clinical testing of gabapentin, the following adverse experiences have been reported in patients receiving other formulations of marketed gabapentin. These adverse experiences have not been listed above and data are insufficient to support an estimate of their incidence or to establish causation. The listing is alphabetized: angioedema, blood glucose fluctuation, breast enlargement, bullous pemphigoid, elevated creatine kinase, elevated liver function tests, erythema multiforme, fever, hyponatremia, jaundice, movement disorder, Stevens-Johnson syndrome.

Adverse events following the abrupt discontinuation of gabapentin immediate release have also been reported. The most frequently reported events were anxiety, insomnia,

nausea, pain and sweating.

There are postmarketing reports of life-threatening or fatal respiratory depression in patients taking gabapentin with opioids or other central nervous system (CNS) depressants, or in the setting of underlying respiratory impairment.

## **7 DRUG INTERACTIONS**

*In vitro* studies were conducted to investigate the potential of gabapentin to inhibit the major cytochrome P450 enzymes (CYP1A2, CYP2A6, CYP2C9, CYP2C19, CYP2D6, CYP2E1, and CYP3A4) that mediate drug and xenobiotic metabolism using isoform selective marker substrates and human liver microsomal preparations. Only at the highest concentration tested (171 mcg/mL; 1mM) was a slight degree of inhibition (14% to 30%) of isoform CYP2A6 observed. No inhibition of any of the other isoforms tested was observed at gabapentin concentrations up to 171 mcg/mL (approximately 15 times the  $C_{max}$  at 3,600 mg/day).

Gabapentin is not appreciably metabolized nor does it interfere with the metabolism of commonly coadministered antiepileptic drugs.

The drug interaction data described in this section were obtained from studies involving healthy adults and adult patients with epilepsy.

### **7.1 Phenytoin**

In a single (400 mg) and multiple dose (400 mg three times daily) study of gabapentin immediate release in epileptic patients (N=8) maintained on phenytoin monotherapy for at least 2 months, gabapentin had no effect on the steady-state trough plasma concentrations of phenytoin and phenytoin had no effect on gabapentin pharmacokinetics.

### **7.2 Carbamazepine**

Steady-state trough plasma carbamazepine and carbamazepine 10, 11 epoxide concentrations were not affected by concomitant gabapentin immediate release (400 mg three times daily; N=12) administration. Likewise, gabapentin pharmacokinetics were unaltered by carbamazepine administration.

### **7.3 Valproic Acid**

The mean steady-state trough serum valproic acid concentrations prior to and during concomitant gabapentin immediate release administration (400 mg three times daily; N=17) were not different and neither were gabapentin pharmacokinetic parameters affected by valproic acid.

### **7.4 Phenobarbital**

Estimates of steady-state pharmacokinetic parameters for phenobarbital or gabapentin immediate release (300 mg three times daily; N=12) are identical whether the drugs are administered alone or together.

### **7.5 Naproxen**

Coadministration of single doses of naproxen (250 mg) and gabapentin immediate release (125 mg) to 18 volunteers increased gabapentin absorption by 12% to 15%. Gabapentin immediate release had no effect on naproxen pharmacokinetics. The doses are lower than the therapeutic doses for both drugs. The effect of coadministration of these drugs at therapeutic doses is not known.

## **7.6 Hydrocodone**

Coadministration of gabapentin immediate release (125 mg and 500 mg) and hydrocodone (10 mg) reduced hydrocodone C<sub>max</sub> by 3% and 21%, respectively, and AUC by 4% and 22%, respectively. The mechanism of this interaction is unknown. Gabapentin AUC values were increased by 14%; the magnitude of the interaction at other doses is not known.

## **7.7 Morphine**

When a single dose (60 mg) of controlled-release morphine capsule was administered 2 hours prior to a single dose (600 mg) of gabapentin immediate release in 12 volunteers, mean gabapentin AUC values increased by 44% compared to gabapentin immediate release administered without morphine. The pharmacokinetics of morphine were not affected by administration of gabapentin immediate release 2 hours after morphine. The magnitude of this interaction at other doses is not known.

## **7.8 Cimetidine**

Cimetidine 300 mg decreased the apparent oral clearance of gabapentin by 14% and creatinine clearance by 10%. The effect of gabapentin immediate release on cimetidine was not evaluated. This decrease is not expected to be clinically significant.

## **7.9 Oral Contraceptives**

Gabapentin immediate release (400 mg three times daily) had no effect on the pharmacokinetics of norethindrone (2.5 mg) or ethinyl estradiol (50 mcg) administered as a single tablet, except that the C<sub>max</sub> of norethindrone was increased by 13%. This interaction is not considered to be clinically significant.

## **7.10 Antacid (containing aluminum hydroxide and magnesium hydroxide)**

An antacid containing aluminum hydroxide and magnesium hydroxide reduced the bioavailability of gabapentin immediate release by about approximately 20%, but by only 5% when gabapentin immediate release was taken 2 hours after the antacid. It is recommended that gabapentin tablets be taken at least 2 hours following the antacid (containing aluminum hydroxide and magnesium hydroxide) administration.

## **7.11 Probenecid**

Gabapentin immediate release pharmacokinetic parameters were comparable with and without probenecid, indicating that gabapentin does not undergo renal tubular secretion by the pathway that is blocked by probenecid.

## **7.12 Drug/Laboratory Test Interactions**

False positive readings were reported with the Ames-N-Multistix SG<sup>®</sup> dipstick test for

urine protein when gabapentin was added to other antiepileptic drugs; therefore, the more specific sulfosalicylic acid precipitation procedure is recommended to determine the presence of urine protein.

## **8 USE IN SPECIFIC POPULATIONS**

### **8.1 Pregnancy**

#### Risk Summary

Available data from published prospective and retrospective cohort studies, and case reports over decades of use with gabapentin during pregnancy have not identified a drug-associated risk of major birth defects. The available data are insufficient to evaluate a drug-associated risk of miscarriage and other maternal or fetal outcomes. In nonclinical studies in mice, rats, and rabbits, gabapentin was developmentally toxic (increased fetal skeletal and visceral abnormalities, and increased embryofetal mortality) when administered to pregnant animals at doses similar to those used clinically (see Data).

The estimated background risk of major birth defects and miscarriage for the indicated populations is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively.

#### Data

##### *Animal Data*

When pregnant mice received oral doses of gabapentin (1,000 or 3,000 mg/kg/day, approximately 3 to 8 times the maximum recommended dose of 1,800 mg on a mg/m<sup>2</sup> basis) during the period of organogenesis, embryofetal toxicity (increased incidences of skeletal variations) was observed. The no effect level was 500 mg/kg/day, representing approximately the maximum recommended human dose [MRHD] on a mg/m<sup>2</sup> basis.

When rats were dosed prior to and during mating, and throughout gestation, pups from all dose groups (500, 1,000 and 2,000 mg/kg/day) were affected. These doses are equivalent to approximately 3 to 11 times the MRHD on a mg/m<sup>2</sup> basis. There was an increased incidence of hydroureter and/or hydronephrosis in rats in a study of fertility and general reproductive performance at 2,000 mg/kg/day with no effect at 1,000 mg/kg/day, in a teratology study at 1,500 mg/kg/day with no effect at 300 mg/kg/day, and in a perinatal and postnatal study at all doses studied (500, 1,000 and 2,000 mg/kg/day). The doses at which the effects occurred are approximately 3 to 11 times the maximum recommended dose of 1,800 mg on a mg/m<sup>2</sup> basis; the no-effect doses were approximately 5 times (Fertility and General Reproductive Performance study) and approximately equal to (Teratogenicity study) the MRHD on a mg/m<sup>2</sup> basis. Other than hydroureter and hydronephrosis, the etiologies of which are unclear, the incidence of malformations was not increased compared to controls in offspring of mice, rats, or rabbits given doses up to 8 times (mice), 10 times (rats), or 16 times (rabbits) the human daily dose on a mg/m<sup>2</sup> basis.

When pregnant rabbits were treated with gabapentin during the period of

organogenesis, an increase in embryofetal mortality was observed at 60, 300, and 1,500 mg/kg/day (0.6 to 16 times the MRHD on a mg/m<sup>2</sup> basis).

In a published study, gabapentin (400 mg/kg/day) was administered by intraperitoneal injection to neonatal mice during the first postnatal week, a period of synaptogenesis in rodents (corresponding to the last trimester of pregnancy in humans). Gabapentin caused a marked decrease in neuronal synapse formation in brains of intact mice and abnormal neuronal synapse formation in a mouse model of synaptic repair. Gabapentin has been shown in vitro to interfere with activity of the  $\alpha 2\delta$  subunit of voltage-activated calcium channels, a receptor involved in neuronal synaptogenesis. The clinical significance of these findings is unknown.

## **8.2 Lactation**

### Risk Summary

Gabapentin is present in human milk following oral administration. Adverse effects on the breastfed infant have not been reported. There are no data on the effects of the drug on milk production. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for gabapentin tablets and any potential adverse effects on the breastfed infant from gabapentin tablets or from the underlying maternal condition

## **8.4 Pediatric Use**

The safety and effectiveness of gabapentin tablets in the management of postherpetic neuralgia in patients less than 18 years of age has not been studied.

## **8.5 Geriatric Use**

The total number of patients treated with gabapentin tablets in controlled clinical trials in patients with postherpetic neuralgia was 359, of which 63% were 65 years of age or older. The types and incidence of adverse events were similar across age groups except for peripheral edema, which tended to increase in incidence with age.

Gabapentin tablets are known to be substantially excreted by the kidney. Reductions in gabapentin tablets dose should be made in patients with age-related compromised renal function [see *Dosage and Administration (2.2)*].

## **8.6 Hepatic Impairment**

Because gabapentin is not metabolized, studies have not been conducted in patients with hepatic impairment.

## **8.7 Renal Impairment**

Gabapentin tablets are known to be substantially excreted by the kidney. Dosage adjustment is necessary in patients with impaired renal function. Gabapentin tablets should not be administered in patients with CrCL between 15 and 30 or in patients undergoing hemodialysis [see *Dosage and Administration (2.2)*].

## **9 DRUG ABUSE AND DEPENDENCE**

The abuse and dependence potential of gabapentin tablets has not been evaluated in human studies.

## 10 OVERDOSAGE

Signs of acute toxicity in animals included ataxia, labored breathing, ptosis, sedation, hypoactivity, or excitation.


Acute oral overdoses of gabapentin have been reported. Symptoms include double-vision, tremor, slurred speech, drowsiness, altered mental status, dizziness, lethargy, and diarrhea. Fatal respiratory depression has been reported with gabapentin overdose, alone and in combination with other central nervous system (CNS) depressants.

Gabapentin can be removed by hemodialysis. Hemodialysis has been performed in overdose cases reported, and it may be indicated by the patient's clinical state or in patients with significant renal impairment.

## 11 DESCRIPTION

Gabapentin tablets contain gabapentin, a gamma-aminobutyric acid (GABA) analogue, as the active pharmaceutical ingredient. Gabapentin's chemical name is 1-(aminomethyl)cyclohexaneacetic acid; with a molecular formula of  $C_9H_{17}NO_2$  and a molecular weight of 171.24 g/mol.

Gabapentin chemical structural formula is:



Gabapentin is a white to off-white crystalline solid with a pKa1 of 3.7 and a pKa2 of 10.7. It is freely soluble in water and acidic and basic solutions. The log of the partition coefficient (n-octanol/ 0.05M phosphate buffer) at pH 7.4 is -1.25.

Gabapentin is intended for oral administration and is supplied as tablets containing 300 mg or 600 mg of gabapentin.

Each 300 mg tablet contains the inactive ingredients copovidone, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene oxide, and Opadry® II white. Opadry® II white contains polyvinyl alcohol, polyethylene glycol, titanium dioxide, and talc.

Each 600 mg tablet contains the inactive ingredients copovidone, hypromellose,

magnesium stearate, polyethylene oxide, and Opadry® II orange. Opadry® II orange contains D&C yellow #10 aluminum lake, FD&C red #40 aluminum lake, polyvinyl alcohol, polyethylene glycol, titanium dioxide, and talc.

## **12 CLINICAL PHARMACOLOGY**

### **12.1 Mechanism of Action**

The mechanism of action by which gabapentin exerts its analgesic action is unknown but in animal models of analgesia, gabapentin prevents allodynia (pain-related behavior in response to a normally innocuous stimulus) and hyperalgesia (exaggerated response to painful stimuli). Gabapentin prevents pain-related responses in several models of neuropathic pain in rats and mice (e.g., spinal nerve ligation models, spinal cord injury model, acute herpes zoster infection model). Gabapentin also decreases pain-related responses after peripheral inflammation (carrageenan footpad test, late phase of formalin test), but does not alter immediate pain-related behaviors (rat tail flick test, formalin footpad acute phase). The relevance of these models to human pain is not known.

Gabapentin is structurally related to the neurotransmitter GABA (gamma-aminobutyric acid), but it does not modify GABAA or GABAB radioligand binding, it is not converted metabolically into GABA or a GABA agonist, and it is not an inhibitor of GABA uptake or degradation. In radioligand binding assays at concentrations up to 100 µM, gabapentin did not exhibit affinity for a number of other receptor sites, including benzodiazepine, glutamate, Nmethyl-D-aspartate (NMDA), quisqualate, kainate, strychnine-insensitive or strychnine-sensitive glycine; alpha 1, alpha 2, or beta adrenergic; adenosine A1 or A2; cholinergic, muscarinic, or nicotinic; dopamine D1 or D2; histamine H1; serotonin S1 or S2; opiate mu, delta, or kappa; cannabinoid 1; voltage-sensitive calcium channel sites labeled with nitrendipine or diltiazem; or at voltage-sensitive sodium channel sites labeled with batrachotoxinin A20-alpha-benzoate. Gabapentin did not alter the cellular uptake of dopamine, noradrenaline, or serotonin.

In vitro studies with radiolabeled gabapentin have revealed a gabapentin binding site in areas of rat brain including neocortex and hippocampus. A high-affinity binding protein in animal brain tissue has been identified as an auxiliary subunit of voltage-activated calcium channels. However, functional correlates of gabapentin binding, if any, remain to be elucidated. It is hypothesized that gabapentin antagonizes thrombospondin binding to  $\alpha 2\delta$ -1 as a receptor involved in excitatory synapse formation and suggested that gabapentin may function therapeutically by blocking new synapse formation.

### **12.2 Pharmacodynamics**

No pharmacodynamic studies have been conducted with gabapentin tablets.

### **12.3 Pharmacokinetics**

#### Absorption

Gabapentin is absorbed from the proximal small bowel by a saturable L-amino transport system. Gabapentin bioavailability is not dose proportional; as the dose is increased, bioavailability decreases.

When gabapentin tablets (1,800 mg once daily) and gabapentin immediate release (600 mg three times a day) were administered with high fat meals (50% of calories from fat), gabapentin tablets have a higher  $C_{max}$  and lower AUC at steady state compared to gabapentin immediate release (Table 5). Time to reach maximum plasma concentration ( $T_{max}$ ) for gabapentin tablets is 8 hours, which is about 4 to 6 hours longer compared to gabapentin immediate release.

**Table 5: Mean  $\pm$  SD Steady-State Pharmacokinetics for Gabapentin Tablets and Gabapentin Immediate Release in Healthy Subjects under high-fat high calorie fed state (Day 5, n = 21)**

<b>Pharmacokinetic Parameter (Mean <math>\pm</math> SD)</b>	<b>Gabapentin Tablets 1,800 mg QD (3 X 600 mg)</b>	<b>Gabapentin Immediate Release 600 mg TID</b>
<b>AUC<sub>0-24</sub> (mcg <math>\bullet</math> hr/mL)</b>	132.8 $\pm$ 34.7	141.3 $\pm$ 29.8
<b>C<sub>max</sub>(mcg/mL)</b>	9.59 $\pm$ 2.33	8.54 $\pm$ 1.72
<b>C<sub>min</sub>(mcg/mL)</b>	1.84 $\pm$ 0.65	2.6 $\pm$ 0.78
<b>T<sub>max</sub> (hr)<sup>1</sup></b>	8 (3-12)	2 (1-5) <sup>*</sup>

<sup>1</sup>T<sub>max</sub> is presented as median (range); \*relative to most recent dose

Do not use once-daily gabapentin tablets interchangeably with other gabapentin products because of differing pharmacokinetic profiles that affect frequency of administration.

Gabapentin tablets should be taken with evening meals. If it is taken on an empty stomach, the bioavailability will be substantially lower.

Administration of gabapentin tablets with food increases the rate and extent of absorption of gabapentin compared to the fasted state.  $C_{max}$  of gabapentin increases 33-84% and AUC of gabapentin increases 33-118% with food depending on the fat content of the meal. Gabapentin tablets should be taken with food.

### Distribution

Gabapentin is less than 3% bound to plasma proteins. After 150 mg intravenous administration, the mean  $\pm$  SD volume of distribution is 58  $\pm$  6 L.

### Elimination

Gabapentin is eliminated by renal excretion as unchanged drug.

In patients with normal renal function given gabapentin immediate release 1,200 to 3,000 mg/day, the drug elimination half-life ( $t_{1/2}$ ) was 5 to 7 hours. Elimination kinetics do not change with dose level or multiple doses.

### *Metabolism*

Gabapentin is not appreciably metabolized in humans.

### *Excretion*

Gabapentin elimination rate constant, plasma clearance, and renal clearance are directly

proportional to creatinine clearance. In elderly patients and patients with impaired renal function, plasma clearance is reduced. Gabapentin can be removed from plasma by hemodialysis.

Dosage adjustment in patients with compromised renal function is necessary. In patients undergoing hemodialysis, Gabapentin tablets should not be administered [see *Dosage and Administration (2.2)*].

## **12.4 Special Populations**

### ***Renal Insufficiency***

As renal function decreases, renal and plasma clearances and the apparent elimination rate constant decrease, while  $C_{max}$  and  $t_{1/2}$  increase.

In patients (N=60) with creatinine clearance of at least 60, 30 to 59, or less than 30 mL/min, the median renal clearance rates for a 400 mg single dose of gabapentin immediate release were 79, 36, and 11 mL/min, respectively, and the median  $t_{1/2}$  values were 9.2, 14, and 40 hours, respectively.

Dosage adjustment is necessary in patients with impaired renal function [see *Dosage and Administration (2.2)*].

### ***Hemodialysis***

In a study in anuric adult subjects (N=11), the apparent elimination half-life of gabapentin on nondialysis days was about 132 hours; during dialysis the apparent half-life of gabapentin was reduced to 3.8 hours. Hemodialysis thus has a significant effect on gabapentin elimination in anuric subjects. Gabapentin tablets should not be administered in patients undergoing hemodialysis. Alternative formulations of gabapentin products should be considered in patients undergoing hemodialysis.

### ***Elderly***

Apparent oral and renal clearances of gabapentin decrease with increasing age, although this may be related to the decline in renal function with age. Reductions in gabapentin dose should be made in patients with age-related compromised renal function [see *Dosage and Administration (2.2)*].

### ***Hepatic Impairment***

Because gabapentin is not metabolized, studies have not been conducted in patients with hepatic impairment.

### ***Pediatrics***

The pharmacokinetics of gabapentin tablets have not been studied in patients less than 18 years of age.

### ***Gender***

Although no formal study has been conducted to compare the pharmacokinetics of gabapentin in men and women, it appears that the pharmacokinetic parameters for males and females are similar and there are no significant gender differences.

## **Race**

Pharmacokinetic differences due to race have not been studied. Because gabapentin is primarily renally excreted and there are no important racial differences in creatinine clearance, pharmacokinetic differences due to race are not expected.

## **13 NONCLINICAL TOXICOLOGY**

### **13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility**

#### Carcinogenesis

Gabapentin was given in the diet to mice at 200, 600, and 2,000 mg/kg/day and to rats at 250, 1,000, and 2,000 mg/kg/day for 2 years. A statistically significant increase in the incidence of pancreatic acinar cell adenoma and carcinomas was found in male rats receiving the high dose; the no-effect dose for the occurrence of carcinomas was 1,000 mg/kg/day. Peak plasma concentrations of gabapentin in rats receiving the high dose of 2,000 mg/kg/day were more than 10 times higher than plasma concentrations in humans receiving 1,800 mg per day and in rats receiving 1,000 mg/kg/day peak plasma concentrations were more than 6.5 times higher than in humans receiving 1,800 mg/day. The pancreatic acinar cell carcinomas did not affect survival, did not metastasize and were not locally invasive. The relevance of this finding to carcinogenic risk in humans is unclear.

Studies designed to investigate the mechanism of gabapentin-induced pancreatic carcinogenesis in rats indicate that gabapentin stimulates DNA synthesis in rat pancreatic acinar cells *in vitro* and, thus, may be acting as a tumor promoter by enhancing mitogenic activity. It is not known whether gabapentin has the ability to increase cell proliferation in other cell types or in other species, including humans.

#### Mutagenesis

Gabapentin did not demonstrate mutagenic or genotoxic potential in 3 *in vitro* and 4 *in vivo* assays. It was negative in the Ames test and the *in vitro* HGPRT forward mutation assay in Chinese hamster lung cells; it did not produce significant increases in chromosomal aberrations in the *in vitro* Chinese hamster lung cell assay; it was negative in the *in vivo* chromosomal aberration assay and in the *in vivo* micronucleus test in Chinese hamster bone marrow; it was negative in the *in vivo* mouse micronucleus assay; and it did not induce unscheduled DNA synthesis in hepatocytes from rats given gabapentin.

#### Impairment of Fertility

No adverse effects on fertility or reproduction were observed in rats at doses up to 2,000 mg/kg (approximately 11 times the maximum recommended human dose on an mg/m<sup>2</sup> basis).

## **14 CLINICAL STUDIES**

The efficacy of gabapentin tablets for the management of postherpetic neuralgia was established in a double-blind, placebo-controlled, multicenter study. This study enrolled

patients between the age of 21 to 89 with postherpetic neuralgia persisting for at least 6 months following healing of herpes zoster rash and a minimum baseline pain intensity score of at least 4 on an 11-point numerical pain rating scale ranging from 0 (no pain) to 10 (worst possible pain).

This 11-week study compared gabapentin tablets 1,800 mg once daily with placebo. A total of 221 and 231 patients were treated with gabapentin tablets or placebo, respectively. The study treatment including titration for all patients comprised a 10-week treatment period followed by 1-week of dose tapering. Double-blind treatment began with titration starting at 300 mg/day and titrated up to a total daily dose of 1,800 mg over 2 weeks, followed by 8 weeks fixed dosing at 1,800 mg once daily, and then 1 week of dose tapering. During the 8-week stable dosing period, patients took 3 active or placebo tablets each night with the evening meal. During baseline and treatment, patients recorded their pain in a daily diary using an 11-point numeric pain rating scale. The mean baseline pain score was 6.6 and 6.5 for gabapentin tablets and placebo-treated patients, respectively.

Treatment with gabapentin tablets statistically significantly improved the endpoint mean pain score from baseline. For various degrees of improvement in pain from baseline to study endpoint, Figure 1 shows the fraction of patients achieving that degree of improvement. The figure is cumulative, so that patients whose change from baseline is, for example, 50%, are also included at every level of improvement below 50%. Patients who did not complete the study were assigned 0% improvement.




Figure 1: Percent of Patients Achieving Various Levels of Pain Relief

## 16 HOW SUPPLIED/STORAGE AND HANDLING

Once-daily Gabapentin Tablets are supplied as follows:

300 mg tablets:

Gabapentin 300 mg tablets are white, oval film-coated tablet debossed with “300” above “ ” on one side and plain on the other side.

NDC 42806-656-09 (Bottle of 90)

600 mg tablets:

Gabapentin 600 mg tablets are orange, oval film-coated tablet debossed with “600” above “ ” on one side and plain on the other side.

NDC 42806-657-09 (Bottle of 90)

**Storage**

Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature].

Keep out of reach of children.

**17 PATIENT COUNSELING INFORMATION**

Advise patients of the availability of a Medication Guide, and instruct them to read the Medication Guide prior to taking gabapentin tablets.

- Advise patients that once-daily gabapentin tablets are not interchangeable with other formulations of gabapentin.
- Advise patients to take gabapentin tablets only as prescribed. Gabapentin tablets may cause dizziness, somnolence, and other signs and symptoms of CNS depression.
- Advise patients not to drive or operate other complex machinery until they have gained sufficient experience on gabapentin tablets to gauge whether or not it adversely affects their mental and/or motor performance. Advise patients who require concomitant treatment with morphine to tell their prescriber if they develop signs of CNS depression such as somnolence. If this occurs the dose of gabapentin tablets or morphine should be reduced accordingly.
- Advise patients that if they miss a dose of gabapentin tablets to take it with food as soon as they remember. If it is almost time for the next dose, just skip the missed dose and take the next dose at the regular time. Do not take two doses at the same time.
- Advise patients that if they take too much gabapentin tablets, to call their healthcare provider or poison control center, or go to the nearest emergency room right away.

Suicidal Thoughts and Behavior

Advise patients, their caregivers, and families that AEDs, including gabapentin, the active ingredient in gabapentin tablets, may increase the risk of suicidal thoughts and behavior and should be advised of the need to be alert for the emergence or worsening of symptoms of depression, any unusual changes in mood or behavior, or the emergence of suicidal thoughts, behavior, or thoughts about self-harm. Behaviors of concern should be reported immediately to healthcare providers [see *Warnings and Precautions*

(5.1)].

### Respiratory Depression

Inform patients about the risk of respiratory depression. Include information that the risk is greatest for those using concomitant central nervous system (CNS) depressants (such as opioid analgesics) or in those with underlying respiratory impairment. Teach patients how to recognize respiratory depression and advise them to seek medical attention immediately if it occurs [see *Warnings and Precautions (5.2)*].

### Dosing and Administration

Once-daily gabapentin tablets are not interchangeable with other gabapentin products because of differing pharmacokinetic profiles that affect the frequency of administration.

The safety and effectiveness of gabapentin tablets in patients with epilepsy has not been studied.

Advise patients that gabapentin tablets should be taken orally once daily with the evening meal. Gabapentin tablets should be swallowed whole. Do not split, crush, or chew the tablets [see *Dosage and Administration (2.1)*].

### Pregnancy

Advise patients to notify their healthcare provider if they become pregnant or intend to become pregnant during treatment with gabapentin tablets [see *Use in Specific Populations (8.1)*].

Dispense with Medication Guide available at: [www.epic-pharma.com/medguide/Gabapentin-Tablets.pdf](http://www.epic-pharma.com/medguide/Gabapentin-Tablets.pdf)

Distributed by:

**Epic pharma, LLC**

Laurelton, NY 11413

LN0006

Rev. 09-2023-02

All Product/Brand names are the trademarks of their respective owners.

**Dispense with Medication Guide available at: [www.epic-pharma.com/medguide/Gabapentin-Tablets.pdf](http://www.epic-pharma.com/medguide/Gabapentin-Tablets.pdf)**

## **Medication Guide**

### **Gabapentin**

**(gab" a pen' tin)**

### **Tablets (Once-Daily)**

Read this Medication Guide before you start taking gabapentin tablets and each time you get a refill. There may be new information. This information does not take the place of talking to your healthcare provider about your medical condition or treatment. If you have any questions about gabapentin tablets, ask your healthcare provider or pharmacist.

## **What is the most important information I should know about gabapentin tablets?**

**Do not stop taking gabapentin tablets without first talking with your healthcare provider.** Stopping gabapentin tablets suddenly can cause serious problems.

Like other antiepileptic drugs, gabapentin, the active ingredient in gabapentin tablets, may cause suicidal thoughts or actions in a very small number of people, about 1 in 500. However, it is not known if gabapentin tablets are safe and effective in people with seizure problems (epilepsy). Therefore, once-daily gabapentin tablets should not be used in place of other gabapentin products.

**Call a healthcare provider right away if you have any of these symptoms, especially if they are new, worse, or worry you:**

- thoughts about suicide or dying
- attempts to commit suicide
- serious breathing problems
- new or worse depression
- new or worse anxiety
- feeling agitated or restless
- panic attacks
- trouble sleeping (insomnia)
- new or worse irritability
- acting aggressive, being angry, or violent
- acting on dangerous impulses
- an extreme increase in activity and talking (mania)
- other unusual changes in behavior or mood

**How can I watch for early symptoms of suicidal thoughts and actions?**

- Pay attention to any changes, especially sudden changes, in mood, behaviors, thoughts, or feelings.
- Keep all follow-up visits with your healthcare provider as scheduled.
- Call your healthcare provider between visits as needed, especially if you are worried about symptoms.

## **Serious breathing problems**

- Serious breathing problems can occur when gabapentin tablets are taken with other medicines that can cause severe sleepiness or decreased awareness, or when it is taken by someone who already has breathing problems. Watch for increased sleepiness or decreased breathing when starting gabapentin tablets or when the dose is increased. Get help right away if breathing problems occur.

**Do not stop taking gabapentin tablets without first talking with your healthcare provider.**

- Stopping gabapentin tablets suddenly can cause serious problems.

## **What are gabapentin tablets?**

Gabapentin tablets are prescription medicine used in adults, 18 years and older, to treat:

- pain from damaged nerves (neuropathic pain) that follows healing of shingles (a painful rash that comes after a herpes zoster infection).

It is not known if gabapentin tablets are safe and effective in people with seizure problems (epilepsy).

It is not known if gabapentin tablets are safe and effective in children under 18 years of age with postherpetic pain.

Once-daily gabapentin tablets are not interchangeable with other gabapentin products.

### **Who should not take gabapentin tablets?**

Do not take gabapentin tablets if you are allergic to gabapentin or any of the ingredients in gabapentin tablets. See the end of this Medication Guide for a complete list of ingredients in gabapentin tablets.

### **What should I tell my healthcare provider before taking gabapentin tablets?**

Before taking gabapentin tablets, tell your healthcare provider if you:

- have or have had depression, mood problems or suicidal thoughts or behavior
- have breathing problems
- have seizures
- have kidney problems or get kidney dialysis
- are pregnant or plan to become pregnant. It is not known if gabapentin tablets can harm your unborn baby. Tell your healthcare provider right away if you become pregnant while taking gabapentin tablets. You and your healthcare provider will decide if you should take gabapentin tablets while you are pregnant.
- are breastfeeding or plan to breastfeed. Gabapentin passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby during treatment with gabapentin tablets.

Tell your healthcare provider about all the medicines you take including prescription and nonprescription medicines, vitamins or herbal supplements. Especially tell your healthcare provider if you take any opioid pain medicine (such as oxycodone), or medicines for anxiety (such as lorazepam) or insomnia (such as zolpidem). You may have a higher chance for dizziness, sleepiness, or serious breathing problems if these medicines are taken with gabapentin tablets.

Taking gabapentin tablets with certain other medicines can cause side effects or affect how well they work. Do not start or stop other medicines without talking to your healthcare provider.

Know the medicines you take. Keep a list of them and show it to your healthcare provider and pharmacist when you get a new medicine.

### **How should I take gabapentin tablets?**

- Take gabapentin tablets exactly as prescribed. Your healthcare provider will tell you how much gabapentin tablets to take and when to take it. Take gabapentin tablets at the same time each day.
- **Do not change your dose or stop taking gabapentin tablets without talking with your healthcare provider.** If you stop taking gabapentin tablets suddenly, you may experience side effects. Talk with your healthcare provider

about how to stop gabapentin tablets slowly.

- Take gabapentin tablets with food one time each day with your evening meal.
- Take gabapentin tablets whole. Do not split, crush, or chew gabapentin tablets before swallowing.
- Your healthcare provider may change your dose of gabapentin tablets. Do not change your dose of gabapentin tablets without talking to your healthcare provider.
- If you miss a dose, take it as soon as you remember with food. If it is almost time for your next dose, just skip the missed dose. Take the next dose at your regular time. **Do not take two doses at the same time.**
- If you take too much gabapentin tablets, call your healthcare provider or poison control center, or go to the nearest emergency room right away.
- If you are taking an antacid containing aluminum hydroxide and magnesium hydroxide, it is recommended that gabapentin tablets be taken at least 2 hours following administration of the antacid.

### **What should I avoid while taking gabapentin tablets?**

- Do not drink alcohol or take other medicines that make you sleepy or dizzy while taking gabapentin tablets without first talking to your healthcare provider. Taking gabapentin tablets with alcohol or medicines that cause sleepiness or dizziness may make your sleepiness or dizziness worse.
- Do not operate heavy machines or do other dangerous activities until you know how gabapentin tablets affect you. Gabapentin tablets can slow your thinking and motor skills.

### **What are the possible side effects of gabapentin tablets?**

The most common side effect of gabapentin tablets are:

- dizziness

Tell your healthcare provider about any side effect that bothers you or that does not go away.

These are not all the possible side effects of gabapentin tablets. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

### **How should I store gabapentin tablets?**

Store gabapentin tablets at 59°F to 86°F (15°C to 30°C).

- **Keep gabapentin tablets and all medicines out of the reach of children.**

### **General information about the safe and effective use of gabapentin tablets**

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use gabapentin tablets for a condition for which it was not prescribed. Do not give gabapentin tablets to other people, even if they have the same symptoms you have. It may harm them.

This Medication Guide summarizes the most important information about gabapentin

tablets. If you would like more information, talk with your healthcare provider. You can ask your healthcare provider or pharmacist for information about gabapentin tablets that is written for health professionals.

For more information about gabapentin tablets, call 1-888-374-2791.

### What are the ingredients in gabapentin tablets?

Active ingredient: gabapentin

Inactive ingredients:

300 mg tablet: copovidone, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene oxide, and Opadry® II white. Opadry® II white contains polyvinyl alcohol, polyethylene glycol, titanium dioxide, and talc.

600 mg tablet: copovidone, hypromellose, magnesium stearate, polyethylene oxide, and Opadry® II orange. Opadry® II orange contains D&C yellow #10 aluminum lake, FD&C red #40 aluminum lake, polyvinyl alcohol, polyethylene glycol, titanium dioxide, and talc.

Distributed by:

**Epic Pharma, LLC**

Laurelton NY, 11413

LG0005

Rev. 09-2023-02

This Medication Guide has been approved by the U.S. Food and Drug Administration.

All Product/Brand names are the trademarks of their respective owners.

### PACKAGE/LABEL PRINCIPAL DISPLAY PANEL

Gabapentin Tablets, USP

300mg 90 Tablets

**Rx Only**

**Dosage and Administration:** See Prescribing Information.

**Take with evening meal. Swallow tablet whole. Do not crush, split or chew.**

Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature].

LE0061  
Rev. 09-2023-00

NDC 42806-656-09

**ONCE-DAILY  
Gabapentin  
Tablets  
300 mg**

Do not use Once-Daily Gabapentin Tablets interchangeably with other gabapentin products.

**PHARMACIST:** Dispense the accompanying Medication Guide to each patient.

**Rx Only  
90 Tablets**

**Each Tablet Contains:**  
300 mg of gabapentin.

You are encouraged to report side effects of prescription drugs to the FDA.

Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch), or call FDA at 1-800-FDA-1088.

**Keep out of reach of children.**

Print Medication Guide at: [www.epic-pharma.com/medguide/Gabapentin-Tablets.pdf](http://www.epic-pharma.com/medguide/Gabapentin-Tablets.pdf)

Distributed by:  
**Epic Pharma, LLC**  
Laurelton, NY 11413

# PACKAGE/LABEL PRINCIPAL DISPLAY PANEL

Gabapentin Tablets, USP

600 mg 90 Tablets

**Rx Only**

**Dosage and Administration:** See Prescribing Information.  
**Take with evening meal. Swallow tablet whole. Do not crush, split or chew.**  
Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature].  
LE0062  
Rev. 09-2023-00

**NDC 42806-657-09**  
**ONCE-DAILY**  
**Gabapentin**  
**Tablets**  
**600 mg**

Do not use Once-Daily Gabapentin Tablets interchangeably with other gabapentin products.  
**PHARMACIST:** Dispense the accompanying Medication Guide to each patient.  
**Rx Only**  
**90 Tablets**

**Each Tablet Contains:**  
600 mg of gabapentin.

You are encouraged to report side effects of prescription drugs to the FDA.  
Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch), or call FDA at 1-800-FDA-1088.  
**Keep out of reach of children.**  
Print Medication Guide at: [www.epic-pharma.com/medguide/Gabapentin-Tablets.pdf](http://www.epic-pharma.com/medguide/Gabapentin-Tablets.pdf)

Distributed by:  
**Epic Pharma, LLC**  
Laurelton, NY 11413

**EPIC PHARMA**

GTIN XXXXXXXXXXXXXXXX  
SN XXXXXXXXXXXXXXXX  
EXP XXXXX  
LOT XXXXXXXXX

3 42806-657-09 6

## GABAPENTIN

gabapentin tablet

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:42806-656
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>GABAPENTIN</b> (UNII: 6CW7F3G59X) (GABAPENTIN - UNII:6CW7F3G59X)	GABAPENTIN	300 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>COPOVIDONE K25-31</b> (UNII: D9C330MD8B)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ05DW1A)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	

### Product Characteristics

<b>Color</b>	WHITE	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	16mm
<b>Flavor</b>		<b>Imprint Code</b>	300
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:42806-656-09	90 in 1 BOTTLE; Type 0: Not a Combination Product	02/26/2024	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA216252	02/26/2024	

## GABAPENTIN

gabapentin tablet

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:42806-657
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>GABAPENTIN</b> (UNII: 6CW7F3G59X) (GABAPENTIN - UNII:6CW7F3G59X)	GABAPENTIN	600 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>COPOVIDONE K25-31</b> (UNII: D9C330MD8B)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	

### Product Characteristics

<b>Color</b>	ORANGE	<b>Score</b>	no score
--------------	--------	--------------	----------

<b>Shape</b>	OVAL	<b>Size</b>	19mm
<b>Flavor</b>		<b>Imprint Code</b>	600
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:42806-657-09	90 in 1 BOTTLE; Type 0: Not a Combination Product	02/26/2024	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA216252	02/26/2024	

**Labeler** - Epic Pharma, LLC (827915443)

**Registrant** - Epic Pharma, LLC (827915443)

**Establishment**

Name	Address	ID/FEI	Business Operations
Epic Pharma, LLC		827915443	MANUFACTURE(42806-656, 42806-657)

Revised: 2/2024

Epic Pharma, LLC