ARNICARE BRUISE- arnica montana tablet Boiron

Disclaimer: This homeopathic product has not been evaluated by the Food and Drug Administration for safety or efficacy. FDA is not aware of scientific evidence to support homeopathy as effective.

Arnicare Bruise Tablets

Active ingredient** (in each tablet)

Arnica montana 9C HPUS (2.5 mg)

The letters "HPUS" indicate that the components in this product are officially monographed in the Homeopathic Pharmacopoeia of the United States.

Purpose*

Arnica montana 9C HPUS (2.5 mg) ... Relieves discoloration, pain, and swelling associated with bruising

Uses*

reduces symptoms of bruising such as: discoloration, pain, swelling

Ask a doctor before use if

- bruising limits the range of motion of your eyes, limbs, head
- pain is severe
- bruising is frequent.

Stop use and ask a doctor if symptoms persist for more than 7 days or worsen, new symptoms occur, or if redness or swelling is present. These could be signs of a serious condition.

If pregnant or breastfeeding, ask a health profession before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

Adults and children 6 years of age and older - As soon as bruising occurs, dissolve 2 tablets in the mouth and repeat every hour for 2 more hours. Then, dissolve 2 tablets in the mouth every 6 hours until bruising has faded.

Children 2 to 5 years of age - Dissolve 2 tablets in 1 tablespoon of water for easier intake. Follow directions above.

Children under 2 years of age - Ask a doctor.

- do not use if glued carton end flaps are open or if the blister seal is broken
- store below 86° F (30° C)

60 meltaway tablets

Bruise relief*

Discoloration, Pain, Swelling*

No Known Drug Interactions

Non-Drowsy

Gentle on the stomach

Convenient & Painless

Meltaway Tablets Require No Food or Water

*CLAIMS BASED ON TRADITIONAL HOMEOPATHIC PRACTICE, NOT ACCEPTED MEDICAL EVIDENCE. NOT FDA EVALUATED.

**C,K,CK, and X are homeopathic dilutions: see BoironUSA.com/info for details.

croscarmellose sodium, lactose, magnesium stearate

Arnicare.com

BoironUSA.com

Info@Boiron.com

1-800-BOIRON-1

(1-800-264-7661)

Made in France

Distributed by Boiron Inc.

Newtown Square, PA 19073



Adults and children 6 years of age and older: As soon as bruising occurs, dissolve 2 tablets in the mouth and repeat every hour for 2 more hours. Then, dissolve 2 tablets in the mouth every 6 hours until bruising has faded. Children 2 to 5 years of age: Dissolve 2 tablets in 1 tablespoon of water for easier intake. Follow directions above. Children under 2 years of age: Ask a doctor.







ARNICARE BRUISE

arnica montana tablet

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0220-9083
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name Basis of Strength Strength

ARNICA MONTANA (UNII: 080TY208ZW) (ARNICA MONTANA - UNII:080TY208ZW) | ARNICA MONTANA | 9 [hp_C]

Inactive Ingredients Ingredient Name Strength CROSCARMELLOSE SODIUM (UNII: M28OL1HH48) MAGNESIUM STEARATE (UNII: 70097M6I30) LACTOSE, UNSPECIFIED FORM (UNII: J2B2A4N98G)

Product Characteristics			
Color	white	Score	no score
Shape	ROUND (Boiron)	Size	9mm
Flavor		Imprint Code	;
Contains			

Packaging					
	# Item Code Package Description		Marketing Start Marketing B Date Date		
	1	NDC:0220- 9083-04	60 in 1 BLISTER PACK; Type 0: Not a Combination Product	01/18/2022	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved homeopathic		01/18/2022	
·			

Labeler - Boiron (282560473)

Registrant - Boiron Inc. (014892269)

Establishment				
Name	Address	ID/FEI	Business Operations	
Boiron		282560473	manufacture(0220-9083)	

Revised: 10/2023 Boiron