# TOPLAST PAIN RELIEF LIDOCAINE 4PERCENT PLUS MENTHOL 1PERCENTlidocaine, menthol patch Icure Pharmaceutical Inc, Wanju Factory

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

-----

# **Drug Facts**

# **ACTIVE INGREDIENTS (in each patch)**

Lidocaine 4% Menthol 1%

### **INACTIVE INGREDIENTS**

Glycerin, D-Sorbitol Solution, Propylene Glycol, Sodium Polyacrylate, Polyacrylic Acid, Urea, Carboxymethylcellulose Sodium, Gelatin, Kaolin, Titanium Oxide, Tartaric Acid, Dihydroxyaluminum Aminoacetate, Methylparaben, Polysorbate 80, Sorbitan Oleate, Disodium Edetate Hydrate, Propylparaben, Water

### **PURPOSE**

Topical Anesthetic Topical Analgesic

#### WARNINGS

# For external use only

Safety Warning: Prior to using the product, conduct a skin patch test on your forearm to assess sensitivity and/or allergic reactions

------

### Do not use

 $\blacksquare$  More than 1 patch on your body at a time  $\blacksquare$  On the cut, irritated, swollen or puncture wounds  $\blacksquare$  For more than one week without consulting a doctor

\_\_\_\_\_

**Ask a doctor or pharmacist before use** if you are allergic to any active or inactive ingredients

\_\_\_\_\_\_

# When using this product

■ Use only as directed ■ Read and follow all directions and warnings on this label ■ Do not allow contact with the eyes and mucous membranes ■ Do not bandage tightly or apply local heat (such as heating pads) to the area of use ■ Do not use at the same time as other topical analgesics ■ Make sure to dispose the used patch away from children and pets, as used ones still contain the drug that can cause serious adverse effects on them if they chew or ingest the patch

\_\_\_\_\_\_

# Stop use and ask a doctor if

■ Condition worsens ■ Redness is present ■ Irritation develops ■ Symptoms persist for more than 7 days or recur ■ Skin swells or blisters where the product was applied

-----

If pregnant or breast-feeding, ask a health professional before use Keep out of reach of children and pets. If swallowed, get medical help or contact a Poison Control Center right away

# Keep out of reach of children and pets

If swallowed, get medical help or contact a Poison Control Center right away

### Uses

For temporary relief of minor pain

#### **Directions**

# Adult and children 12 years of age and over :

■ Clean and dry affected area
 ■ Peel film from patch and apply to the skin (see illustration)
 ■ Use 1 patch at a time and do not use for more than 12 hours
 Children under 12 years of age: consult a physician

### Other Information

■ Store at room temperature 68° to 77°F (20° to 25°C) ■ Store in a clean, dry place out of direct sunlight ■ Protect from excessive moisture

#### **Ouestions or Comments**

■ Visit us at : icure.co.kr/en ■ Contact us : icurepharm@icure.co.kr


#### PACKAGE LABEL.PRINCIPAL DISPLAY PANEL



Lidocaine 4% plus Menthol 1%







Drug Facts

Active Ingredients (in each patch)
Lidocaine 4%
Menthol 1%

Topical An Topical Analgesi

Uses For temporary relief of minor pair

### Warnings For external use only

Safety Warning: Prior to using the product, conduct a skin patch test on your forearm to assess sensitivity and/or allergic reactions

Do not use

• More than 1 patch on your body at a time • On the cut, irritated, swollen or puncture wounds • For more than one week without consulting a doctor

Ask a doctor or pharmacist before use if you are allergic to any active or inactive ingredients

Ask a doctor or pharmacist before use if you are allergic to any active or inactive ingredients.

\*When using this product

\*Use only as directed \*Read and follow all directions and warnings on this label \*Po
not allow contact with the eyes and mucous membranes \*Po not bandage tightly or
apply local heat (such as heating pads) to the area of use \*Po not use at the same time
as other topical analgesics \*Nakes sure to dispose the used patch away from children
and pets, as used ones still contain the drug that can cause serious adverse effects
on them if they other or ingest the patch

\*Stop use and ask a doctor if

\*Condition vorsens \*Pedness is present \*Irritation develops \*Symptoms persist for more
than 7 days or recur \*Skin swells or bilisters where the product was applied

\*Irrennant or breast-faceling ask a health professional before use

If pregnant or breast-feeding, ask a health professional before use
Keep out of reach of children and pets. If swallowed, get medical help or contact
a Poison Control Center right away

Directions
Adult and children 12 years of age and over:

-Clean and dry affected area -Peel film from patch and apply to the skin (see illustration)

-Use 1 patch at a time and do not use for more than 12 hours

Children under 12 years of age: consult a physician

Other Information

\*Store at room temperature 68° to 77°F ( 20° to 25°C) \*Store in a clean, dry place out of direct sunlight \* Protect from excessive moisture

Questions or Comments?

•Visit us at: icure.co.kr/en

•Contact us: icurepharm@icure.co.kr

Inactive Ingredients Glycerin, D-Sorbitol Solution, Propylene Glycol, Sodium Polyacrylate, Polyacrylic Acid, Urea, Carboxymethylcellulose Sodium, Gelatin, Kaolini Titanium Oxide, Tarlaric Acid, Dihydroxyaluminum Aminoacetate, Methylparaben, Polysorbate 80, Sorbitan Oleate, Disodium Edelate Hydrate, Propylparaben, Water

Manufactured By

icure ICURE PHARM. INC.
248 Wanjusandan 5-ro.
Bongdong-eup, Wanju-gun,
Jeollabuk-do (55322),

52 x 17.5 (mm)



(Toplast')

# TOPLAST PAIN RELIEF LIDOCAINE 4PERCENT PLUS MENTHOL 1PERCENT

lidocaine, menthol patch

# **Product Information**

**HUMAN OTC DRUG** NDC:73279-0004 **Product Type** Item Code (Source)

**TOPICAL Route of Administration** 

# **Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
Lidocaine (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	Lidocaine	240 mg
Menthol (UNII: L7T10EIP3A) (MENTHOL - UNII:L7T10EIP3A)	Menthol	60 mg

<b>Inactive Ingredients</b>
-----------------------------

ı		
	Ingredient Name	Strength
ı		

Glycerin (UNII: PDC6A3C0OX)

SORBITOL (UNII: 506T60A25R)	
Propylene Glycol (UNII: 6DC9Q167V3)	
Urea (UNII: 8W8T17847W)	
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED (UNII: K6790BS311)	
Gelatin (UNII: 2G86QN327L)	
Kaolin (UNII: 24H4NWX5CO)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
Tartaric Acid (UNII: W4888I119H)	
Dihydroxyaluminum Aminoacetate (UNII: DO250MG0W6)	
Methylparaben (UNII: A2I8C7HI9T)	
Polysorbate 80 (UNII: 6OZP39ZG8H)	
SORBITAN MONOOLEATE (UNII: 06XEA2VD56)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
Propylparaben (UNII: Z8IX2SC1OH)	
Water (UNII: 059QF0KO0R)	

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73279- 0004-2	5 in 1 CARTON	07/01/2020	
1	NDC:73279- 0004-1	1 in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph not final	part348	07/01/2020		

# Labeler - Icure Pharmaceutical Inc, Wanju Factory (695687612)

# Registrant - Icure Pharmaceutical Inc, Wanju Factory (695687612)

Establishment				
Na me	Address	ID/FEI	Business Operations	
Icure Pharmaceutical Inc, Wanju Factory		695687612	manufacture(73279-0004)	

Revised: 4/2021 Icure Pharmaceutical Inc, Wanju Factory