

**ASCURIN- aspirin capsule**  
**Healthdoc LLC**

-----  
**Ascurin (Aspirin Capsules 81mg)**

***Drug Facts***

***Active ingredient (per capsule)***

Aspirin (**NSAID\***) 81mg

***Purpose***

pain reliever-fever reducer \*nonsteroidal anti-inflammatory drug

***Use***

For the temporary relief of minor aches and pains associated with a cold, headache, toothache, muscular aches, backache, the premenstrual and menstrual periods, for the minor pain from arthritis and to reduce fever

***Warnings***

**Reye's syndrome:**Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- Hives
- Facial swelling
- Asthma (wheezing)
- Shock

**Stomach bleeding warning:**This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

***Do not use***

if you have ever had an allergic reaction to any other pain reliever, fever reducer

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

**Ask a doctor or pharmacist before use if you are**

- taking a prescription drug for diabetes, gout, or arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
  - pain gets worse or lasts more than 10 days
  - fever gets worse or lasts more than 3 days
  - redness or swelling is present
  - any new symptoms occur
  - ringing in the ears or loss of hearing occurs

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

***Directions***

- Adults: Drink a full glass of water with each dose.
- Adults and children 12 years and over: Take 5-8 capsules every 4 hours not to exceed 49 capsules in 24 hours unless directed by a doctor.
- Children under 12 years of age: consult a doctor.
- See your doctor for other uses of aspirin, but do not use for more than 10 days without consulting your doctor because serious side effects may occur.

***Other information***

Do not use the product if the bottle seal inside the bottle lid is broken.

***Inactive ingredients***

Curcumin, Hydroxypropyl Methylcellulose, Orange Oil, Polyvinylpyrrolidone.

## Questions or Comments?

Call (609) 902-7128 9:00 AM – 7:00 PM E.T. Monday-Friday

## Package Labeling:

**HealthDoc**  
**AsCurin**  
Adult's Aspirin (NSAID) Capsules  
Pain Reliever-Fever Reducer  
81 mg per capsule

**Drug Facts**

Active ingredient (per capsule)	Purpose
Aspirin (NSAID)* 81mg	pain reliever- fever reducer
*nonsteroidal anti-inflammatory drug	

**Use**  
For the temporary relief of minor aches and pains associated with a cold, headache, toothache, muscular aches, backache, the premenstrual and menstrual periods, for the minor pain from arthritis and to reduce fever.

**Warnings**  
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
Allergy alert: Aspirin may cause a severe allergic reaction which may include:  
■ Hives  
■ Facial swelling  
■ Asthma (wheezing)  
■ Shock  
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you  
■ are age 60 or older  
■ have had stomach ulcers or bleeding problems  
■ take a blood thinning (anticoagulant) or steroid drug  
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
■ have 3 or more alcoholic drinks every day while using this product  
■ take more or for a longer time than directed  
Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer  
Ask a doctor before use if  
■ stomach bleeding warning applies to you  
■ you have a history of stomach problems, such as heartburn  
■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease  
■ you are taking a diuretic  
■ you have asthma  
Ask a doctor or pharmacist before use if you are  
■ taking a prescription drug for diabetes, gout, or arthritis  
Stop use and ask a doctor if  
■ an allergic reaction occurs. Seek medical help right away.  
■ you experience any of the following signs of stomach bleeding:  
■ feel faint  
■ vomit blood  
■ have bloody or black stools  
■ have stomach pain that does not get better  
■ pain gets worse or lasts more than 10 days  
■ fever gets worse or lasts more than 3 days  
■ redness or swelling is present  
■ any new symptoms occur  
■ ringing in the ears or loss of hearing occurs  
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless expressly directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.  
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

**Drug Facts (continued)**

**Directions**  
■ Adults: Drink a full glass of water with each dose.  
■ Adults and children 12 years and over: Take 5-8 capsules every 4 hours not to exceed 49 capsules in 24 hours unless directed by a doctor.  
■ Children under 12 years of age: consult a doctor.  
■ See your doctor for other uses of aspirin, but do not use for more than 10 days without consulting your doctor because serious side effects may occur.

**Other Information**  
Do not use the product if the bottle seal inside the bottle lid and outside of the lid is broken.


**Inactive Ingredients** Curcumin, Hydroxypropyl Methylcellulose, Orange Oil, Polyvinylpyrrolidone.

**Questions or Comments?** Call (609) 902-7128 9:00 AM – 7:00 PM E.T. Monday-Friday

**Distributed by:** HealthDoc LLC  
1004 Industrial Dr. Ste 3  
Berlin Township, NJ 08091  
United States  
www.ascurin.com

**Consult your Doctor before use**


**HealthDoc**  
**AsCurin**  
Adult's Aspirin (NSAID) Capsules  
Pain Reliever-Fever Reducer  
81 mg per capsule



**30 Capsules      81 mg per capsule**

Other Information - Save carton for full directions and warnings - Store at 15-30°C (59-86°F) Avoid excessive heat above 40°C (104°F).

Lot #  
Exp.

  
8 55921 00857 0

**HealthDoc**  
**AsCurin**  
Adult's Aspirin (NSAID) Capsules  
Pain Reliever-Fever Reducer  
81 mg per capsule

**30 Capsules      81 mg per capsule**

**THIS UNIT NOT LABELED FOR RETAIL SALE**

**See carton for complete warnings and information**

**ACTIVE INGREDIENT (PER CAPSULE)**  
ASPIRIN (NSAID)\* 81MG  
\*NONSTEROIDAL ANTI-INFLAMMATORY DRUG

**DIRECTIONS**  
ADULTS: DRINK A FULL GLASS OF WATER WITH EACH DOSE. ADULTS AND CHILDREN 12 YEARS AND OVER: TAKE 5-8 CAPSULES EVERY 4 HOURS NOT TO EXCEED 49 CAPSULES IN 24 HOURS UNLESS DIRECTED BY A DOCTOR. SEE YOUR DOCTOR FOR OTHER USES OF ASPIRIN, BUT DO NOT USE FOR MORE THAN 10 DAYS WITHOUT CONSULTING YOUR DOCTOR BECAUSE SERIOUS SIDE EFFECTS MAY OCCUR.

**WARNINGS**  
**STOMACH BLEEDING WARNING:** THIS PRODUCT CONTAINS AN NSAID, WHICH MAY CAUSE SEVERE STOMACH BLEEDING. THE CHANCE IS HIGHER IF YOU ■ ARE AGE 60 OR OLDER ■ HAVE HAD STOMACH ULCERS OR BLEEDING PROBLEMS ■ TAKE A BLOOD THINNING (ANTICOAGULANT) OR STEROID DRUG ■ TAKE OTHER DRUGS CONTAINING PRESCRIPTION OR

NONPRESCRIPTION NSAIDS (ASPIRIN, IBUPROFEN, NAPROXEN, OR OTHERS)  
■ HAVE 3 OR MORE ALCOHOLIC DRINKS EVERY DAY WHILE USING THIS PRODUCT  
■ TAKE MORE OR FOR A LONGER TIME THAN DIRECTED  
**ALLERGY ALERT:** ASPIRIN MAY CAUSE A SEVERE ALLERGIC REACTION WHICH MAY INCLUDE HIVES, FACIAL SWELLING, SHOCK AND ASTHMA (WHEEZING).  
**REYE'S SYNDROME:** CHILDREN AND TEENAGERS WHO HAVE OR ARE RECOVERING FROM CHICKEN POX OR FLU-LIKE SYMPTOMS SHOULD NOT USE THIS PRODUCT. WHEN USING THIS PRODUCT, IF CHANGES IN BEHAVIOR WITH NAUSEA AND VOMITING OCCUR, CONSULT A DOCTOR BECAUSE THESE SYMPTOMS COULD BE AN EARLY SIGN OF REYE'S SYNDROME, A RARE BUT SERIOUS ILLNESS.

**DISTRIBUTED BY:** HEALTHDOC LLC  
1004 INDUSTRIAL DR. STE 3  
BERLIN TOWNSHIP, NJ 08091  
UNITED STATES  
WWW.ASCURIN.COM (609) 902-7128

Lot #  
Exp.

**ASCURIN**

aspirin capsule

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:83607-002
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>CURCUMIN</b> (UNII: IT942ZTH98)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>ORANGE OIL, COLD PRESSED</b> (UNII: AKN3KSD11B)	

## Product Characteristics

<b>Color</b>	white (Colorless)	<b>Score</b>	no score
<b>Shape</b>	CAPSULE	<b>Size</b>	25mm
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
<b>1</b>	NDC:83607-002-00	1 in 1 BOX	12/10/2023	
<b>1</b>		30 in 1 BOTTLE; Type 0: Not a Combination Product		

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M013	12/10/2023	

**Labeler** - Healthdoc LLC (013084196)