

**KIRKLAND SIGNATURE NAPROXEN SODIUM- naproxen sodium tablet, film coated**  
**Costco Wholesale Company**

-----  
**Costco Wholesale Corp. Naproxen Sodium Tablets USP, 220 mg Drug Facts**

**Active ingredient (in each caplet)**

Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)\*

\*nonsteroidal anti-inflammatory drug

**Purposes**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis
- muscular aches
- backache
- menstrual cramps
- headache
- toothache
- the common cold
- temporarily reduces fever

**Warnings**

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen,

or others)

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

### **Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

### **Ask a doctor before use if**

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have asthma

### **Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking any other drug

### **When using this product**

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
  - pain gets worse or lasts more than 10 days
  - fever gets worse or lasts more than 3 days
  - you have difficulty swallowing
  - it feels like the pill is stuck in your throat
  - redness or swelling is present in the painful area
  - any new symptoms appear

### **If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

### Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose

Adults and children 12 years and older	<ul style="list-style-type: none"><li>• take 1 caplet every 8 to 12 hours while symptoms last</li><li>• for the first dose you may take 2 caplets within the first hour</li><li>• do not exceed 2 caplets in any 8- to 12-hour period</li><li>• do not exceed 3 caplets in a 24-hour period</li></ul>
Children under 12 years	<ul style="list-style-type: none"><li>• ask a doctor</li></ul>

### Other information

- each caplet contains: sodium 21 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

### Inactive ingredients

FD&C blue no. 2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

### Questions or comments?

1-800-719-9260

### Principal Display Panel

COMPARE TO ALEVE® active ingredient

NAPROXEN SODIUM Tablets USP, 220 mg

Pain Reliever/Fever Reducer

(NSAID)

Strength to Last 12 HOURS

Actual Size

400 CAPLETS†

†capsule shaped tablets

**Drug Facts** (continued)

**Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding:  
 ■ feel faint ■ vomit blood ■ have bloody or black stools  
 ■ have stomach pain that does not get better  
 ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days  
 ■ you have difficulty swallowing ■ it feels like the pill is stuck in your throat  
 ■ redness or swelling is present in the painful area ■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions** ■ do not take more than directed  
 ■ the smallest effective dose should be used ■ drink a full glass of water with each dose  
 Adults and children 12 years and older ■ take 1 caplet every 8 to 12 hours while symptoms last  
 ■ for the first dose you may take 2 caplets within the first hour  
 ■ do not exceed 2 caplets in any 8- to 12-hour period  
 ■ do not exceed 3 caplets in a 24-hour period  
 Children under 12 years ■ ask a doctor

**Other information** ■ each caplet contains: sodium 21 mg  
 ■ store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

**Inactive ingredients** FD&C blue no. 2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

**Questions or comments?** 1-800-719-9260  
 15V0367a

**Actual Size**

**Questions or comments?** 1-800-719-9260  
 \*This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Aleve®.  
**GLUTEN FREE**  
 Distributed by: Perrigo®; 615 Eastern Ave. Allegan, MI 49010 USA 1-800-719-9260  
 For: Costco Wholesale Corporation P.O. Box 34535 Seattle, WA 98124-1535 USA 1-800-774-2678 www.costco.com

**KIRKLAND Signature**

**COMPARE TO ALEVE®** active ingredient\* NDC 63981-368-79 ITM./ART. 197197

**NAPROXEN SODIUM**  
 Tablets USP, 220 mg

**Pain Reliever/Fever Reducer**

**(NSAID)**  
**Strength to Last 12 HOURS**

**400 CAPLETS†**  
 †capsule shaped tablets

**DO NOT USE IF PRINTED FOIL UNDER CAP IS BROKEN OR MISSING**

**Drug Facts**

**Active Ingredient (in each caplet)** Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)\* .....Pain reliever/fever reducer  
 \*nonsteroidal anti-inflammatory drug

**Purposes**

**Uses** ■ temporarily relieves minor aches and pains due to:  
 ■ minor pain of arthritis ■ muscular aches ■ backache ■ menstrual cramps  
 ■ headache ■ toothache ■ the common cold  
 ■ temporarily reduces fever

**Warnings**

**Allergy alert:** Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing)  
 ■ shock ■ skin reddening ■ rash ■ blisters  
 If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  
 ■ are age 60 or older ■ have had stomach ulcers or bleeding problems  
 ■ take a blood thinning (anticoagulant) or steroid drug  
 ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product  
 ■ take more or for a longer time than directed

**Do not use** ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer  
 ■ right before or after heart surgery

**Ask a doctor before use if** ■ the stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have asthma

**Ask a doctor or pharmacist before use if you are** ■ under a doctor's care for any serious condition ■ taking any other drug

**When using this product** ■ take with food or milk if stomach upset occurs ■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

3679 SN F10

**KIRKLAND SIGNATURE NAPROXEN SODIUM**

naproxen sodium tablet, film coated

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:63981-368
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
NAPROXEN SODIUM (UNII: 9 TN8 7S3A3C) (NAPROXEN - UNII:57Y76 R9 ATQ)	NAPROXEN SODIUM	220 mg

**Inactive Ingredients**

Ingredient Name	Strength
HYPROMELLOSES (UNII: 3NXW29 V3WO)	
MAGNESIUM STEARATE (UNII: 70 09 7M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D6 1U)	
POLYETHYLENE GLYCOLS (UNII: 3WJQ0 SDW1A)	
POVIDONES (UNII: FZ989 GH9 4E)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9 V2JP)	
FD&C BLUE NO. 2 (UNII: L06K8 R7DQK)	

**Product Characteristics**

<b>Color</b>	BLUE (Light Blue)	<b>Score</b>	no score
<b>Shape</b>	CAPSULE (oval)	<b>Size</b>	12mm
<b>Flavor</b>		<b>Imprint Code</b>	L368
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63981-368-79	400 in 1 BOTTLE; Type 0: Not a Combination Product	06/04/2007	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074661	06/04/2007	

**Labeler** - Costco Wholesale Company (103391843)

Revised: 12/2016

Costco Wholesale Company