

BASIC CARE LOPERAMIDE HYDROCHLORIDE- loperamide hcl solution
Amazon.com Services LLC

Amazon Loperamide Hydrochloride Drug Facts

Active ingredient (in each 7.5 mL)

Loperamide HCl 1 mg

Purpose

Anti-diarrheal

Use

controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Heart alert: Taking more than directed can cause serious heart problems or death

Do not use

if you have bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product

tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- shake well before using
- use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.
- mL = milliliter

adults and children 12 years and over	30 mL after the first loose stool; 15 mL after each subsequent loose stool; but no more than 60 mL in 24 hours
children 9-11 years (60-95 lbs)	15 mL after the first loose stool; 7.5 mL after each subsequent loose stool; but no more than 45 mL in 24 hours
children 6-8 years (48-59 lbs)	15 mL after the first loose stool; 7.5 mL after each subsequent loose stool; but no more than 30 mL in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

Other information

- **each 30 mL contains:** sodium 15 mg
- store between 20-25°C (68-77°F)

Inactive ingredients

anhydrous citric acid, carboxymethylcellulose sodium, D&C yellow no. 10, FD&C blue no. 1, glycerin, microcrystalline cellulose, natural and artificial mint flavor, propylene glycol, purified water, simethicone, sodium benzoate, sucralose, titanium dioxide, xanthan gum

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

Anti-Diarrheal Oral Solution

Compare to Imodium[®] A-D active ingredient

Loperamide Hydrochloride

Oral Solution, 1 mg per 7.5 mL

Anti-Diarrheal

Controls the Symptoms of Diarrhea

Mint Flavor

See New Warning and Directions

8 FL OZ (240 mL)

Drug Facts

Active ingredient Purpose
(in each 7.5 mL)

Loperamide HCl 1 mg.....Anti-diarrheal

Use controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Heart alert: Taking more than directed can cause serious heart problems or death

Do not use if you have bloody or black stool

Peel open to read complete directions and warnings before purchase

amazon

basic⁺care

Anti-Diarrheal Oral Solution

NDC 72288-645-34

Loperamide Hydrochloride
Oral Solution, 1 mg per 7.5 mL
Anti-Diarrheal

Controls the Symptoms of Diarrhea
Mint Flavor

See New Warning and Directions
8 FL OZ (240 mL)

Compare to
Imodium[®] A-D
active
ingredient*

Do not use if
printed plastic
neckband is
broken or missing.

DISTRIBUTED BY:
AMAZON.COM
SERVICES LLC
410 TERRY AVENUE N.
SEATTLE, WA 98109

© 2020 AMAZON.COM, INC.
OR ITS AFFILIATES. ALL
RIGHTS RESERVED. BASIC
CARE AND ALL RELATED
LOGOS ARE TRADEMARKS
OF AMAZON.COM, INC. OR
ITS AFFILIATES.

GLUTEN FREE

*This product is not
manufactured or
distributed by
Johnson & Johnson
Consumer Inc.,
distributor of
Imodium[®] A-D.



8

: 64534 BN F1

Drug Facts (continued)

Ask a doctor before use if you have

- fever ■ mucus in the stool ■ a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are
taking a prescription drug. Loperamide may interact with
certain prescription drugs.

When using this product tiredness, drowsiness or
dizziness may occur. Be careful when driving or operating
machinery.

Stop use and ask a doctor if ■ symptoms get worse
■ diarrhea lasts for more than 2 days
■ you get abdominal swelling or bulging.
These may be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional
before use.

Keep out of reach of children. In case of overdose, get
medical help or contact a Poison Control Center right
away. (1-800-222-1222)

Drug Facts (continued)

Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- shake well before using
- use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.
- mL = milliliter

adults and children
12 years and over

30 mL after the first loose stool;
15 mL after each subsequent
loose stool; but no more than
60 mL in 24 hours

children 9-11 years
(60-95 lbs)

15 mL after the first loose stool;
7.5 mL after each subsequent
loose stool; but no more than
45 mL in 24 hours

Drug Facts (continued)

children 6-8 years
(48-59 lbs)

15 mL after the first loose stool;
7.5 mL after each subsequent
loose stool; but no more than
30 mL in 24 hours

children 2-5 years
(34 to 47 lbs)

ask a doctor

children under
2 years
(up to 33 lbs)

do not use

Other information

- each 30 mL contains: sodium 15 mg
- store between 20-25°C (68-77°F)

Drug Facts (continued)

Inactive ingredients anhydrous citric acid,
carboxymethylcellulose sodium, D&C yellow no. 10,
FD&C blue no. 1, glycerin, microcrystalline cellulose,
natural and artificial mint flavor, propylene glycol,
purified water, simethicone, sodium benzoate,
sucralose, titanium dioxide, xanthan gum

Questions or comments?

1-800-719-9260

BASIC CARE LOPERAMIDE HYDROCHLORIDE

loperamide hcl solution

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:72288-645
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	1 mg in 7.5 mL

Inactive Ingredients	
Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED FORM (UNII: K679OBS311)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)	
GLYCERIN (UNII: PDC6A3C0OX)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
XANTHAN GUM (UNII: TTV12P4NEE)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	

Product Characteristics			
Color	GREEN (opaque, viscous)	Score	
Shape		Size	
Flavor	MINT	Imprint Code	
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72288-645-34	240 mL in 1 BOTTLE; Type 0: Not a Combination Product	01/21/2021	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date

ANDA	ANDA091292	01/21/2021	
------	------------	------------	--

Labeler - Amazon.com Services LLC (128990418)

Revised: 1/2021

Amazon.com Services LLC