

**NIX- permethrin rinse  
Insight Pharmaceuticals LLC**

---

**Nix Creme Rinse - 63736-030**

**Drug Facts**

**Active ingredient (in each fluid ounce)**

Permethrin 280 mg (1%)

**Purpose**

Lice treatment

**Use**

treats head lice

**Warnings**

**For external use only**

**Do not use**

- on children under 2 months of age
- near the eyes
- inside the nose, ear, mouth, or vagina
- on lice in eyebrows or eyelashes. See your doctor.

**Ask a doctor before use if you are**

allergic to ragweed. May cause breathing difficulty or an asthmatic episode.

**When using this product**

- keep eyes tightly closed and protect eyes with a washcloth or towel
- if product gets into the eyes, immediately flush with large amounts of water
- scalp itching or redness may occur

**Stop use and see a doctor if**

- breathing difficulty occurs
- eye irritation occurs
- skin or scalp irritation continues or infection occurs

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**

**Important: Read warnings before use**

**■ Adults and children 2 months of age and over:**

**Inspect**

- all household members should be checked by another person for lice and/or nits (eggs)
- use a magnifying glass in bright light to help you see the lice and nits (eggs)
- use a tool, such as a comb or two unsharpened pencils to lift and part the hair
- look for tiny nits near the scalp, beginning at the back of the neck and behind the ears
- small sections of hair (1-2 inches wide) should be examined at a time
- unlike dandruff, nits stick to the hair. Dandruff should move when lightly touched.
- if either lice or nits (eggs) are found, treat with Nix® Creme Rinse

**Treat**

- wash hair with a shampoo without conditioner. Do not use a shampoo that contains a conditioner or a conditioner alone since this may decrease the activity of Nix®. Rinse with water.
- towel dry hair so it is damp but not wet
- shake the bottle of Nix® well
- completely saturate the hair and scalp with Nix®. Begin to apply Nix® behind the ears and at the back of the neck.
- keep Nix® out of the eyes. Protect the eyes with a washcloth or towel.
- leave Nix® on the hair for 10 minutes, but no longer
- rinse with warm water
- towel dry hair and comb out tangles
- if live lice are seen seven days or more after the first treatment, a second treatment should be given

**Remove Lice/Nits**

- remove nits by combing the hair with the special small tooth comb provided. Remaining nits may be removed by hand (using a throw-away glove), or cutting the nits out.
- use the nit comb provided and make sure the hair remains slightly damp while removing nits
- if the hair dries during combing, dampen it slightly with water
- part the hair into 4 sections. Work on one section at a time. Longer hair may take more time (1-2 hours).
- start at the top of the head on the section you have picked
- with one hand, lift a 1-2 inch wide strand of hair. Get the teeth of the comb as close to the scalp as possible and comb with a firm, even motion away from the scalp to the end of the hair.

- use clips to pin back each strand of hair after you have combed out the nits
- clean the comb completely as you go. Wipe the nits from the comb with a tissue and throw away the tissue in a sealed plastic bag to prevent the lice from coming back.
- after combing, recheck the entire head for nits and repeat combing if necessary
- check the affected head daily to remove any nits that you might have missed

### **Other information**

- read all the directions in the Consumer Information Insert and warnings before use. Keep the carton. It contains important information.
- store at 20° to 25°C (68° to 77°F)

### **Inactive ingredients**

canada balsam, cetyl alcohol, citric acid, FD&C yellow no. 6, hydrolyzed animal protein, hydroxyethylcellulose, isopropyl alcohol, methylparaben, polyoxyethylene 10 cetyl ether, propylene glycol, propylparaben, stearalkonium chloride, water

### **Questions or comments?**

Call 1-888-LICE LINE (1-888-542-3546).

Monday to Friday, 9AM-5PM EST

### **PRINCIPAL DISPLAY PANEL**

**Nix®  
Lice Killing Creme Rinse**

Includes 1 bottle

Plus 1 nit comb

Spanish instructions included

2 FL OZ (59 mL)

<b>Drug Facts (continued)</b>
<b>Directions</b>
Important: Read warnings before use.
• Adults and children 2 months of age and over:
<b>Inspect</b>
▪ all household members should be checked by another person for lice and/or nits (eggs)
▪ use a magnifying glass in bright light to help you see the lice and nits (eggs)
▪ use a comb, such as a comb or two unsharpened pencils to lift and part the hair
▪ look for live nits (1-2 inches wide) should be examined at a time
▪ unlike dandruff, nits stick to the hair; Dandruff should move when lightly touched.
▪ if other lice or nits (eggs) are found, treat with Nix® Creme Rinse
<b>Test</b>
▪ wash hair with a shampoo without conditioner. Do not use a shampoo that contains a conditioner or a conditioner alone since this may decrease the activity of Nix®. Rinse with water.
▪ towel dry hair so it is damp but not wet
▪ shake the bottle of Nix® well
▪ completely saturate the hair and scalp with Nix®. Begin to apply Nix® behind the ears and at the back of the neck.
▪ keep Nix® out of the eyes. Protect the eyes with a washcloth or towel.
▪ leave Nix® on the hair for 10 minutes, but no longer
▪ rinse with warm water
▪ towel dry hair and comb out tangles
▪ if live lice are seen seven days or more after the first treatment, a second treatment should be given.
<b>Remove Lice/Nits</b>
▪ remove nits by combing the hair with the special small tooth comb provided. Remaining nits may be removed by hand using a throw-away glove, or cutting the nits out.
▪ use the nit comb provided and make sure the hair remains slightly damp while removing nits
▪ if the hair dries during combing, dampen it slightly with water
▪ part the hair into 4 sections. Work on one section at a time. Longer hair may take more time (1-2 hours).
▪ start at the top of the head on the section you have picked
▪ with one hand, lift a 1-2 inch wide strand of hair. Get the teeth of the comb as close to the scalp as possible and comb with a firm, even motion away from the scalp to the end of the hair.
▪ use clips to pin back each strand of hair after you have combed out the nits.
▪ clean the comb completely with warm water. Wipe the comb on a tissue and throw away the feces in a sealed plastic bag to prevent the lice from coming back.
▪ after combing, recheck the entire head for nits and repeat combing if necessary
▪ check the affected head daily to remove any nits that you might have missed

<b>Drug Facts (continued)</b>
<b>Other information</b>
▪ Read all the directions in the Consumer Information Insert and warnings before use. Keep the carton. It contains important information.
▪ store at 20° to 25°C (68° to 77°F)
<b>Inactive ingredients</b>
canada balsam, cetyl alcohol, citric acid, FD&C yellow no. 6, hydrolyzed animal protein, hydroxyethylcellulose, isopropyl alcohol, methylparaben, polyoxyethylene 10 cetyl ether, propylene glycol, propylparaben, stearalkonium chloride, water
<b>Questions or comments?</b>
call 1-800-LICE LINE (1-800-542-3546), Monday to Friday, 9 AM - 5 PM EST
<b>LIMITED CONSUMER WARRANTY</b>
If you are not satisfied with Nix®, we will refund your purchase price. Send consumer information insert, UPC #09464, and reason for dissatisfaction to: INSIGHT Pharmaceuticals LLC., 660 White Plains Rd., Tarrytown, NY 10591
nixlice.com
Dist. by INSIGHT Pharmaceuticals LLC 660 White Plains Road Tarrytown, NY 10591 A Johnson & Johnson Healthcare Company © 2021 Trade dress is owned by Insight Pharmaceuticals LLC. All rights reserved. Creme Rinse made in India. Comb made in China.
NXU5003102

**Nix**

PERMETHRIN

Creme Rinse is Safe for use on children as young as 2 months old.

## Lice Killing Creme Rinse

- ✓ Kills lice and eggs in one application
- ✓ Prevents reinestation\*
- ✓ Maximum Strength

**Includes 1 bottle**  
Plus 1 nit comb

Spanish Instructions included

For up to 14 days

2 FL OZ (59 mL)

#1 BRAND

PEDIATRICIAN RECOMMENDED

Have questions?  
Call our expert hotline  
888.542.3546

**Drug Facts**

<b>Active ingredient</b> (in each fluid ounce) Permethrin 200 mg (1%)	<b>Purpose</b> Lice treatment
<b>Use</b> treat head lice	
<b>Warnings</b> For external use only	
<b>Do not use</b>	
• on children under 2 months of age	
• near the eyes	
• inside the nose, ear, mouth, or vagina	
• on lice in eyebrows or eyelashes.	
See your doctor.	
Ask a doctor before use if you are allergic to ragweed. May cause breathing difficulty or an asthma-like episode.	
<b>When using this product</b>	
• keep eyes closed and protect eyes with a washcloth or towel	
• If product gets into the eyes, immediately flush with large amounts of water	
• scalp itching or redness may occur	
<b>Stop use and see a doctor if</b>	
• breathing difficulty occurs	
• eye irritation occurs	
• skin or scalp irritation continues or infection occurs	
<b>If pregnant or breast-feeding,</b> ask a health professional before use.	
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.	

# NIX

permethrin rinse

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:63736-030
<b>Route of Administration</b>	TOPICAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>PERMETHRIN</b> (UNII: 509F88P9SZ) (PERMETHRIN - UNII:509F88P9SZ)	PERMETHRIN	1 g in 100 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>CANADA BALSAM</b> (UNII: K304V198I3)	
<b>CETYL ALCOHOL</b> (UNII: 936JST6JCN)	
<b>CITRIC ACID MONOHYDRATE</b> (UNII: 2968PHW8QP)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>STEARALKONIUM CHLORIDE</b> (UNII: 0OUO26BB88)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>ISOPROPYL ALCOHOL</b> (UNII: ND2M416302)	
<b>METHYLPARABEN</b> (UNII: A2I8C7HI9T)	
<b>PROPYLPARABEN</b> (UNII: Z8IX2SC1OH)	

## Product Characteristics

<b>Color</b>	orange (Light Orange)	<b>Score</b>
<b>Shape</b>		<b>Size</b>
<b>Flavor</b>		<b>Imprint Code</b>

**Contains****Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63736-030-01	1 in 1 CARTON	11/15/2013	
1		59 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:63736-030-02	2 in 1 CARTON	11/15/2013	
2		59 mL in 1 BOTTLE; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA019918	11/15/2013	

**Labeler -** Insight Pharmaceuticals LLC (055665422)

Revised: 12/2021

Insight Pharmaceuticals LLC