

**MUCINEX FAST-MAX CONGESTION AND HEADACHE MAXIMUM STRENGTH-
acetaminophen, dextromethorphan hydrobromide, and phenylephrine
hydrochloride capsule, liquid filled
RB Health (US) LLC**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

**Mucinex® Fast-Max® Congestion & Headache
Maximum Strength**

Drug Facts

<i>Active ingredients (in each liquid gel)</i>	<i>Purposes</i>
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - nasal congestion
 - cough
 - headache
 - minor aches and pains
 - sinus congestion and pressure
- temporarily reduces fever
- temporarily promotes nasal and/or sinus drainage

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 liquid gels in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's Disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see Overdose warning)**
- do not take more than 12 liquid gels in any 24-hour period
- adults and children 12 years of age and over: take 2 liquid gels every 4 hours
- children under 12 years of age: do not use

Other information

- store at 20-25°C (68-77°F)
- avoid excessive heat

Inactive ingredients

FD&C red no. 40, FD&C yellow no. 6, gelatin, glycerin, hypromellose, isopropyl alcohol, lecithin, light mineral oil, polyethylene glycol, povidone, propylene glycol, sorbitol sorbitan solution, titanium dioxide, water

Questions?

1-866-MUCINEX (1-866-682-4639)

You may also report side effects to this phone number.

Dist. by: RB Health (US)
Parsippany, NJ 07054-0224

PRINCIPAL DISPLAY PANEL - 16 Capsule Blister Pack Carton

Fast Dissolving Liquid Gels!

NDC 72854-128-16

MAXIMUM STRENGTH

Mucinex®
FAST-MAX®

CONGESTION &
HEADACHE

Acetaminophen - Pain Reliever/Fever Reducer
Dextromethorphan HBr - Cough Suppressant
Phenylephrine HCl - Nasal Decongestant

- ✓ Relieves Headache & Nasal Congestion
- ✓ Controls Cough
- ✓ Reduces Fever

Actual Size

16
LIQUID GELS
(Liquid Filled Capsules)

DAY TIME
FOR AGES 12+

MAXIMUM STRENGTH



CONGESTION & HEADACHE

Tamper evident: Do not use if carton is damaged or if printed seal on blister is broken or missing.

16 LIQUID GELS

Fast Dissolving Liquid Gels!

MAXIMUM STRENGTH

NDC 72854-128-16



CONGESTION & HEADACHE

Acetaminophen – Pain Reliever/Fever Reducer
Dextromethorphan HBr – Cough Suppressant
Phenylephrine HCl – Nasal Decongestant

- ✓ Relieves Headache & Nasal Congestion
- ✓ Controls Cough
- ✓ Reduces Fever

Actual Size



16 LIQUID GELS
(Liquid Filled Capsules)

DAY TIME
FOR AGES 12+

For Cold & Flu Multi-Symptom relief
for both Day and Night, try our
Maximum Strength Mucinex®
FAST-MAX® Day/Night
liquid gel packs.

Dist. by: RB Health (US)
 Parsippany, NJ 07054-0224

©2021 RB Health

022721 3158958



HEALTH • HYGIENE • HOME

www.mucinex.com

Patents: www.rb.com/patents



6382458716 5

LOT:

EXP.:

3158958



Drug Facts

Active ingredients (in each liquid gel)

Purposes

Acetaminophen 325 mg.....Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....Cough suppressant
Phenylephrine HCl 5 mg.....Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - nasal congestion
 - cough
 - headache
 - minor aches and pains
 - sinus congestion and pressure
- temporarily reduces fever
- temporarily promotes nasal and/or sinus drainage

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 liquid gels in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's Disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus) ▶

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 12 liquid gels in any 24-hour period
- adults and children 12 years of age and over: take 2 liquid gels every 4 hours
- children under 12 years of age: do not use

Other information

- store at 20-25°C (68-77°F)
- avoid excessive heat

Inactive ingredients

FD&C red no. 40, FD&C yellow no. 6, gelatin, glycerin, hypromellose, isopropyl alcohol, lecithin, light mineral oil, polyethylene glycol, povidone, propylene glycol, sorbitol sorbitan solution, titanium dioxide, water

Questions? 1-866-MUCINEX (1-866-682-4639)

You may also report side effects to this phone number.



CONGESTION &
HEADACHE

MAXIMUM STRENGTH

Maximum Strength per 4-hour dose
Do not take more than a total of
12 liquid gels in a 24-hour period.
Take only as directed.
Keep carton for full information.

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

MUCINEX FAST-MAX CONGESTION AND HEADACHE MAXIMUM STRENGTH

acetaminophen, dextromethorphan hydrobromide, and phenylephrine hydrochloride capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:72854-128
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C red no. 40 (UNII: WZB9127XOA)	
FD&C yellow no. 6 (UNII: H77VEI93A8)	
gelatin, unspecified (UNII: 2G86QN327L)	
glycerin (UNII: PDC6A3C0OX)	
hypromellose, unspecified (UNII: 3NXW29V3WO)	
isopropyl alcohol (UNII: ND2M416302)	
light mineral oil (UNII: N6K5787QVP)	
polyethylene glycol, unspecified (UNII: 3WJQ0SDW1A)	
povidone, unspecified (UNII: FZ989GH94E)	
propylene glycol (UNII: 6DC9Q167V3)	
titanium dioxide (UNII: 15FIX9V2JP)	
water (UNII: 059QF0KOOR)	

Product Characteristics

Color	RED	Score	no score
Shape	OVAL	Size	24mm
Flavor		Imprint Code	AR04
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72854-128-16	2 in 1 CARTON	07/26/2021	

1	8 in 1 BLISTER PACK; Type O: Not a Combination Product		
Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part341	07/26/2021	

Labeler - RB Health (US) LLC (081049410)

Revised: 7/2021

RB Health (US) LLC