

**ASPIRIN 325 MG- aspirin tablet  
WALGREENS**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**331R Walgreens 0363-6453 Aspirin 325 mg 500s**

**DRUG FACTS**

***Active ingredient (in each tablet)***

Aspirin 325 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever

***Uses***

- For the temporary relief of minor aches and pains due to:
  - o headache
  - o toothache
  - o muscle pain
  - o minor arthritis pain
  - o menstrual pain
  - o colds
- or as recommended by a doctor

***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinner (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDS (aspirin, ibuprofen, naproxen, or others)

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use:**

- If you have ever had an allergic reaction to any other pain reliever/fever reducer.

**Ask a doctor before use if:**

- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

**Ask a doctor or pharmacist before use if you are**

- taking a prescription drug for diabetes, gout, or arthritis.

**Stop use and ask a doctor if:**

you experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- an allergic reaction occurs

**Seek medical help right away if:**

- new symptoms occur
- ringing in the ears or loss of hearing occurs
- redness or swelling is present
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**

- drink a full glass of water with each dose.
- adults and children 12 years and over: take 1 to 2 tablets every 4 hours while symptoms last. Do not take more than 12 tablets in 24 hours unless directed by a doctor.
- children under 12 years: ask a doctor.

**Other information**

- **each tablet contains:** sodium 1 mg (VERY LOW SODIUM)
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F).
- use by expiration date on package

**Inactive ingredients** corn starch, croscarmellose sodium, D&C yellow #10 aluminum lake, FD & C yellow #6 aluminum lake, hypromellose, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, mineral oil, polysorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, talc, titanium dioxide, triethyl citrate

**Questions or comments?** Call **1-877-290-4008**

**VALUE SIZE**      NDC 0363-6453-53

**LOT / EXP AREA**

**Drug Facts (continued)**

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- hives   ■ facial swelling   ■ asthma (wheezing)   ■ shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinner (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Walgreens**

Compare to the active ingredient in Ecotrin®

**Aspirin 325**

325 mg / PAIN RELIEVER (NSAID)

Regular Strength    Enteric-Coated

**500 TABLETS**

ACTUAL SIZE

**HEART HEALTH**

\*\*Talk to your doctor or other healthcare provider before using this product for your heart.

**Drug Facts**

Active ingredient (in each tablet)	Purpose
Aspirin 325 mg (NSAID)*	Pain reliever
*nonsteroidal anti-inflammatory drug	

**Uses** ■ For the temporary relief of minor aches and pains due to:

- headache   ■ toothache
- muscle pain   ■ minor arthritis pain
- menstrual pain   ■ colds

■ or as recommended by a doctor

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.

This product is not manufactured or distributed by Medtech Products Inc., owner of the registered trademark Ecotrin®.

DISTRIBUTED BY: WALGREEN CO.  
200 WILMOT RD., DEERFIELD, IL 60015

ITEM 284618    W00000-0000-0

100% SATISFACTION GUARANTEED

walgreens.com    ©2022 Walgreen Co.

MADE IN U.S.A. OF U.S.A. AND IMPORTED INGREDIENTS

PEEL HERE FOR MORE DRUG FACTS

W3ORG0422-F    331R 0222

**Drug Facts (continued)**

**Do not use:** ■ If you have ever had an allergic reaction to any other pain reliever/fever reducer.

**Ask a doctor before use if:**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

**Ask a doctor or pharmacist before use if you are**

- taking a prescription drug for diabetes, gout, or arthritis.

**Stop use and ask a doctor if:**

- you experience any of the following signs of stomach bleeding:
  - feel faint   ■ vomit blood   ■ have bloody or black stools
  - have stomach pain that does not get better
  - an allergic reaction occurs

**Seek medical help right away if:**

- new symptoms occur
- ringing in the ears or loss of hearing occurs
- redness or swelling is present
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days

**Drug Facts (continued)**

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**

- drink a full glass of water with each dose.
- adults and children 12 years and over: take 1 to 2 tablets every 4 hours while symptoms last. Do not take more than 12 tablets in 24 hours unless directed by a doctor.
- children under 12 years: ask a doctor.

**Other information**

- each tablet contains: sodium 1 mg (VERY LOW SODIUM)
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F).
- use by expiration date on package

**Inactive ingredients** corn starch, croscarmellose sodium, D&C yellow #10 aluminum lake, FD & C yellow #6 aluminum lake, hypromellose, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, mineral oil, polysorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, talc, titanium dioxide, triethyl citrate

**Questions or comments?** Call 1-877-290-4008

PERMANENT  
ADHESIVE  
AREA

<b>ASPIRIN 325 MG</b>			
aspirin tablet			
<b>Product Information</b>			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0363-6453
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

**Inactive Ingredients**

Ingredient Name	Strength
<b>D&amp;C YELLOW NO. 10 ALUMINUM LAKE</b> (UNII: CQ3XH3DET6)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>MINERAL OIL</b> (UNII: T5L8T28FGP)	
<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)	
<b>CROSCARMELOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O)	
<b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>SODIUM HYDROXIDE</b> (UNII: 55X04QC32I)	
<b>TRIETHYL CITRATE</b> (UNII: 8Z96QXD6UM)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>METHACRYLIC ACID AND ETHYL ACRYLATE COPOLYMER</b> (UNII: NX76LV5T8J)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	

**Product Characteristics**

<b>Color</b>	orange	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	T
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-6453-53	500 in 1 BOTTLE; Type 0: Not a Combination Product	06/15/2022	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	06/15/2022	

**Labeler** - WALGREENS (008965063)**Registrant** - TIME CAP LABORATORIES, INC. (037052099)

## Establishment

Name	Address	ID/FEI	Business Operations
TIME CAP LABORATORIES, INC.		037052099	manufacture(0363-6453)

Revised: 5/2022

WALGREENS