

ASPIRIN 81- aspirin tablet
WALGREENS

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

482R Walgreens 0363-6452 Aspirin 81 mg 500s

DRUG FACTS

Active ingredient (in each tablet)

Aspirin 81 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

- temporarily relieves minor aches and pains
- ask your doctor about other uses for enteric coated 81 mg Aspirin.

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: •hives • facial swelling • asthma (wheezing) • shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if

- stomach bleeding warning applies to you

- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for gout, diabetes or arthritis

Stop use and ask a doctor if

an allergic reaction occurs. Seek medical help right away.
you experience any of the following signs of stomach bleeding:

- feel faint ● vomit blood ● have bloody or black stools
- have stomach pain that does not get better

pain gets worse or lasts more than 10 days
ringing in the ears or a loss of hearing occurs

redness or swelling is present
new symptoms occur. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- Store at 20-25°C (68-77°F)
- close cap tightly after use

Inactive ingredients

anhydrous lactose, carnauba wax, colloidal silicon dioxide, croscarmellose sodium, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, iron oxide yellow, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polysorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, triethyl citrate

Questions or comments? Call 1-877-290-4008

VALUE SIZE NDC 0363-6452-50

Walgreens

Compare to the active ingredient in St. Joseph® Low Dose 81 mg Aspirin*

EASY-OPEN

Aspirin 81

81 mg / PAIN RELIEVER (NSAID)

Low Dose

Enteric-Coated



500
TABLETS

ACTUAL SIZE

**Talk to your doctor or other healthcare provider before using this product for your heart.

NOT FOR HOUSEHOLDS WITH YOUNG CHILDREN

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.

Drug Facts

Active ingredient (in each tablet)
Aspirin 81 mg (NSAID) Pain reliever
nonsteroidal anti-inflammatory drug

Purpose
Pain reliever

Uses ■ temporarily relieves minor aches and pain
■ ask your doctor about other uses for enteric coated 81 mg Aspirin.

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:
■ hives ■ facial swelling ■ asthma (wheezing) ■ shock
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you
■ are age 60 or older

■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
■ have 3 or more alcoholic drinks every day while using this product
■ take more or for a longer time than directed

Do not use

■ if you are allergic to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if

■ stomach bleeding warning applies to you
■ you have a history of stomach problems, such as heartburn
■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
■ you are taking a diuretic
■ you have asthma

Walgreens Pharmacist Recommended. Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands. *This product is not manufactured or distributed by Foundation Consumer Healthcare LLC, owner of the registered trademark St. Joseph® Low Dose 81 mg Aspirin.

81 mg Aspirin. Lot No.: W30RG0422-F
DISTRIBUTED BY: WALGREENS CO. ITEM 285292 W00000-0000-0
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
MADE IN U.S.A. OF U.S. AND IMPORTED INGREDIENTS
3 11917 13650 9

PEEL HERE

Varnish Omit Area

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking a prescription drug for gout, diabetes or arthritis

Stop use and ask a doctor if
■ an allergic reaction occurs. Seek medical help right away.
■ you experience any of the following signs of stomach bleeding:
● feel faint ● vomit blood ● have bloody or black stools
■ have stomach pain that does not get better
■ pain gets worse or lasts more than 10 days
■ ringing in the ears or a loss of hearing occurs
■ redness or swelling is present
■ new symptoms occur. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

■ drink a full glass of water with each dose
■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
■ children under 12 years: do not use unless directed by a doctor

Other information

■ Store at 20-25°C (68-77°F) ■ close cap tightly after use

Inactive ingredients

anhydrous lactose, carnauba wax, colloidal silicon dioxide, croscarmellose sodium, FD&C Red #40 aluminum lake, FD&C Yellow #6 aluminum lake, iron oxide yellow, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, poly sorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, triethyl citrate

Questions or comments? Call 1-877-290-4008

STOP PEELING

ASPIRIN 81
aspirin tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-6452
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
STARCH, CORN (UNII: O8232NY3SJ)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
FD&C RED NO. 4 (UNII: X3W0AM1JLX)	
METHACRYLIC ACID AND ETHYL ACRYLATE COPOLYMER (UNII: NX76LV5T8J)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
TALC (UNII: 7SEV7J4R1U)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	

Product Characteristics

Color	pink (Peach Colored tablets)	Score	no score
Shape	ROUND	Size	7mm
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-6452-50	500 in 1 BOTTLE; Type 0: Not a Combination Product	06/15/2022	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not		06/15/2022	

final	part 543	06/15/2022	
-------	----------	------------	--

Labeler - WALGREENS (008965063)

Registrant - TIME CAP LABORATORIES, INC. (037052099)

Establishment			
Name	Address	ID/FEI	Business Operations
TIME CAP LABORATORIES, INC.		037052099	manufacture(0363-6452)

Revised: 5/2022

WALGREENS