

SODIUM SULFACETAMIDE 10% AND SULFUR 2% CLEANSER- sulfacetamide sodium, sulfur liquid

Method Pharmaceuticals

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Sodium Sulfacetamide 10% - Sulfur 2% Cleanser


Rx Only

DESCRIPTION

Each gram of sodium sulfacetamide 10% and sulfur 2% cleanser contains 100 mg of sodium sulfacetamide and 20 mg of sulfur in a cleanser containing Aloe Vera leaf Extract, Butylated Hydroxytoluene, Cetyl Alcohol, Citric Acid, Cocamidopropyl Betaine, Disodium EDTA, Glycerin, Glyceryl Stearate SE, PEG-100 Stearate, Phenoxyethanol, Purified Water, Sodium laureth sulfate, Sodium Thiosulfate, Stearyl alcohol, Triacetin, Xanthan Gum.

Sodium sulfacetamide is a sulfonamide with antibacterial activity while sulfur acts as a keratolytic agent. Sodium sulfacetamide is $C_8H_9N_2NaO_3 \cdot H_2O$ with molecular weight of 254.24. Chemically, it is N-[(4-aminophenyl)sulfonyl]-acetamide, monosodium salt, monohydrate.

The structural formula is:



Clinical Pharmacology

Sodium sulfacetamide exerts a bacteriostatic effect against sulfonamide sensitive Gram-positive and Gram-negative microorganisms commonly isolated from secondary cutaneous pyogenic infections. It acts by restricting the synthesis of folic acid required by bacteria for growth, by its competition with para-aminobenzoic acid. There is no clinical data available on the degree and rate of systemic absorption of this product when applied to the skin or scalp. However, significant absorption of sodium

sulfacetamide through the skin has been reported.

The following in vitro data is available but the clinical significance is unknown. Organisms that show susceptibility to sodium sulfacetamide are: Streptococci, Staphylococci, E. coli, Klebsiella pneumoniae, Pseudomonas pyocyanea, Salmonella species, Proteus vulgaris, Nocardia and Actinomyces.

The exact mode of action of sulfur in the treatment of acne is unknown, but it has been reported that it inhibits the growth of Propionibacterium acnes and the formation of free fatty acids.

INDICATIONS

SODIUM SULFACETAMIDE 10% - SULFUR 2% CLEANSER is indicated for use in the topical control of acne vulgaris, acne rosacea and seborrheic dermatitis.

CONTRAINDICATIONS:

SODIUM SULFACETAMIDE 10% - SULFUR 2% CLEANSER is contraindicated for use by patients having known hypersensitivity to sulfonamides, sulfur or any other component of this preparation. SODIUM SULFACETAMIDE 10% - SULFUR 2% CLEANSER is not to be used by patients with kidney disease.

WARNINGS

Although it is rare, sensitivity to sodium sulfacetamide may occur. Therefore, caution and careful supervision should be observed when prescribing this drug for patients who may be prone to hypersensitivity to topical sulfonamides. Systemic toxic reactions such as agranulocytosis, acute hemolytic anemia, purpura hemorrhagica, drug fever, jaundice, and contact dermatitis indicate hypersensitivity to sulfonamides. Particular caution should be employed if areas of denuded or abraded skin are involved.

FOR EXTERNAL USE ONLY. NOT FOR INTRAVAGINAL OR OPHTHALMIC USE. (KEEP AWAY FROM EYES).

KEEP OUT OF REACH OF CHILDREN.

In case of accidental ingestion contact a poison control center immediately. Keep container tightly closed.

Precaution

General: If irritation develops, use of the product should be discontinued and appropriate therapy instituted. Patients should be carefully observed for possible local irritation or sensitization during long-term therapy. The object of this therapy is to achieve desquamation without irritation, but sodium sulfacetamide and sulfur can cause reddening and scaling of the epidermis. These side effects are not unusual in the treatment of acne vulgaris, but patients should be cautioned about the possibility.

Information for Patients: Avoid contact with eyes, eyelids, lips and mucous membranes. If accidental contact occurs, rinse with water. If excessive irritation develops, discontinue use and consult your physician.

Carcinogenesis, Mutagenesis and Impairment of Fertility: Long-term studies in animals have not been performed to evaluate carcinogenic potential.

Pregnancy: Category C. Animal reproduction studies have not been conducted with Sodium Sulfacetamide 10% & Sulfur 2% Cleanser. It is also not known whether Sodium Sulfacetamide 10% & Sulfur 2% Cleanser can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Sodium Sulfacetamide 10% & Sulfur 2% Cleanser should be given to a pregnant woman only if clearly needed.

Nursing Mothers: It is not known whether sodium sulfacetamide is excreted in the human milk following topical use of Sodium Sulfacetamide 10% & Sulfur 2% Cleanser. However, small amounts of orally administered sulfonamides have milk. In view of this and because many drugs are excreted in human milk, caution should be exercised when Sodium Sulfacetamide 10% & Sulfur 2% Cleanser is administered to a nursing woman.

Pediatric Use: Safety and effectiveness in children under the age of 12 have not been established.

Adverse Reaction

Although rare, sodium sulfacetamide may cause local irritation.

Call your doctor for medical advice about side effects.

To report a serious adverse event or obtain product information, call 1-877-250-3427.

DOSAGE AND ADMINISTRATION

Wash affected area once or twice daily, or as directed by your physician. Avoid contact with eyes or mucous membranes. Wet skin and liberally apply to areas to be cleansed, massage gently into skin for 10-20 seconds working into a full lather, rinse thoroughly and pat dry. If drying occurs, it may be controlled by rinsing cleanser off sooner or using less often.

See label booklet for Full Prescribing Information.

HOW SUPPLIED

Store at 20°C to 25°C (68°F to 77°F), excursions permitted between 15°C and 30°C (between 59°F and 86°F). Brief exposure to temperatures up to 40°C (104°F) may be tolerated provided the mean kinetic temperature does not exceed 25°C (77°F); however, such exposure should be minimized.

NOTICE: Protect from freezing and excessive heat. The product may tend to darken

slightly on storage. Slight discoloration does not impair the efficacy or safety of the product. Keep bottle tightly closed. Occasionally, a slight discoloration of fabric may occur when an excessive amount of the product is used and comes in contact with white fabrics. This discoloration, however, presents no problem, as it is readily removed by ordinary laundering without bleaches.

This bottle is not filled to the top but does contain 8 fl oz of product as identified on the front panel of the bottle.

Method Pharmaceuticals
NDC 58657-471-08

Sodium Sulfacetamide & Sulfur

(Sodium Sulfacetamide 10% & Sulfur 2%)

10% / 2% Cleanser

For External Use Only

Rx Only
Net Wt. 8 oz (227 g)

INDICATIONS: Sodium Sulfacetamide 10% & Sulfur 2% Cleanser is indicated for the topical control of acne vulgaris, acne rosacea and seborrheic dermatitis.

DIRECTIONS FOR USE: Wash affected area once or twice daily, or as directed by your physician. Avoid contact with eyes or mucous membranes. Wet skin and liberally apply to areas to be cleansed, massage gently into skin for 10-20 seconds working into a full lather, rinse thoroughly and pat dry. If drying occurs, it may be controlled by rinsing cleanser off sooner or using less often. **See label booklet for Full Prescribing Information.**

FOR EXTERNAL USE ONLY. NOT FOR INTRAVAGINAL OR OPHTHALMIC USE. (KEEP AWAY FROM EYES).

KEEP OUT OF REACH OF CHILDREN.

In case of accidental ingestion contact a poison control center immediately. Keep container tightly closed.

CONTRAINDICATIONS: Sodium Sulfacetamide 10% & Sulfur 2% Cleanser is contraindicated in persons with know or suspected hypersensitivity to sulfonamides, sulfur or any other component of this preparation. Sodium Sulfacetamide 10% & Sulfur 2% Cleanser is not to be used by patients with kidney disease.

CAUTION: If redness or irritation occurs, discontinue use.

Each gram of sodium sulfacetamide 10% and sulfur 2% cleanser contains 100 mg of sodium sulfacetamide and 20 mg of sulfur in a cleanser containing Aloe Vera leaf Extract, Butylated Hydroxytoluene, Cetyl Alcohol, Citric Acid, Cocamidopropyl Betaine, Disodium EDTA, Glycerin, Glyceryl Stearate SE, PEG-100 Stearate, Phenoxyethanol, Purified Water, Sodium laureth sulfate, Sodium Thiosulfate, Stearyl alcohol, Triacetin, Xanthan Gum.

STORAGE: Store at 20°C to 25°C (68°F to 77°F), excursions permitted between 15°C and 30°C (between 59°F and 86°F). Brief exposure to temperatures up to 40°C (104°F) may be tolerated provided the mean kinetic temperature does not exceed 25°C (77°F); however, such exposure should be minimized.

NOTICE: Protect from freezing and excessive heat. The product may tend to darken slightly on storage. Slight discoloration does not impair the efficacy or safety of the product. Keep bottle tightly closed.

Occasionally, a slight discoloration of fabric may occur when an excessive amount of the product is used and comes in contact with white fabrics. This discoloration, however, presents no problem, as it is readily removed by ordinary laundering without bleaches.

This bottle is not filled to the top but does contain 8 oz of product as identified on the front panel of the bottle.

Manufactured in the U.S.A. for Method Pharmaceuticals, LLC Southlake, Texas 76092

Sodium Sulfacetamide 10% & Sulfur 2% Cleanser

Rx Only

3 58657 47108 8

SODIUM SULFACETAMIDE 10% AND SULFUR 2% CLEANSER

sulfacetamide sodium, sulfur liquid

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:58657-471
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SULFACETAMIDE SODIUM (UNII: 4NRT660KJQ) (SULFACETAMIDE - UNII:4965G3J0F5)	SULFACETAMIDE SODIUM	100 mg in 1 g
SULFUR (UNII: 70FD1KFU70) (SULFUR - UNII:70FD1KFU70)	SULFUR	20 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
TRIACETIN (UNII: XHX3C3X673)	
ALOE VERA LEAF (UNII: ZY81Z83H0X)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
COCAMIDOPROPYL BETAINE (UNII: 5OCF3O11KX)	
CETYL ALCOHOL (UNII: 936JST6JCN)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
GLYCERYL STEARATE SE (UNII: FCZ5MH785I)	
PEG-100 STEARATE (UNII: YD01N1999R)	
WATER (UNII: 059QF0KO0R)	
SODIUM THIOSULFATE (UNII: HX1032V43M)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
STEARYL ALCOHOL (UNII: 2KR89I4H1Y)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58657-471-08	227 g in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	02/01/2022	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		02/01/2022	

Labeler - Method Pharmaceuticals (060216698)

Revised: 2/2022

Method Pharmaceuticals