ANTACID - aluminum hydroxide, magnesium hydroxide, simethicone liquid Chain Drug Marketing Association Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

QC Regular Strength Antacid

ACTIVE INGREDIENTS (in each 10 mL)

Aluminum hydroxide (equiv. to dried gel, USP) 400 mg Magnesium hydroxide 400 mg Simethicone 40 mg

PURPOSE

Antacid Antacid Antigas

USE(S)

relieves:

- acid indigestion
- heartburn
- sour stomach
- upset stomach and gas associated with these symptoms

WARNINGS

Do not take more than 80 mL in a 24-hour period, or use the maximum dosage of this product for more than 2 weeks, except under the advice and supervision of a doctor.

ASK A DOCTOR BEFORE USE IF YOU HAVE

- kidney disease
- a magnesium-restricted diet

ASK A DOCTOR OR PHARMACIST BEFORE USE IF YOU ARE

• presently taking a prescription drug. Antacids may interact with certain prescription drugs.

STOP USE AND ASK DOCTOR

if symptoms last more than two weeks

KEEP OUT OF REACH OF CHILDREN

.

DIRECTIONS

- shake well before use
- mL = milliliter
- adults and children 12 years and older: take 10 mL to 20 mL four times a day, or as directed by a doctor
- children under 12 years: consult a doctor

OTHER INFORMATION

- each 10 mL contains: magnesium 170 mg, sodium 5 mg
- store at controlled room temperature 20°C-25°C (68°-77°F)
- do not freeze

INACTIVE INGREDIENTS

ethyl alcohol, flavor, glycerin, hydroxyethyl cellulose, methylparaben, propylene glycol, propylparaben, purified water, saccharin sodium, simethicone emulsion, sorbitol

PRINCIPAL DISPLAY PANEL

NDC 63868-173-12 **QC**

QUALITY CHOICE

Regular strength Antacid Liquid Antacid & Antigas


Aluminum Hydroxide 400 mg Magnesium Hydroxide 400 mg Simethicone 40 mg


Relieves:

- Acid Indigestion
- Heartburn
- Sour Stomach
- Pressure & Bloating

Original Flavor Alcohol content 0.2% v/v

12 FL OZ (355 mL)





ANTACID

aluminum hydroxide, magnesium hydroxide, simethicone liquid

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63868-173
Route of Administration	ORAL		

Active	Ingred	ient/A	ctive	Moiet	У

Basis of

_. ..

ingredient Name	Strength	Strength
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0) (ALUMINUM HYDROXIDE - UNII:5QB0T2IUN0)	ALUMINUM HYDROXIDE	400 mg in 10 mL
MAGNESIUM HYDROXIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838)	MAGNESIUM HYDROXIDE	400 mg in 10 mL
DIMETHICONE (UNII: 92RU3N3Y10) (DIMETHICONE - UNII:92RU3N3Y10)	DIMETHICONE	40 mg in 10 mL

Inactive Ingredients			
Ingredient Name	Strength		
ALCOHOL (UNII: 3K9958V90M)			
GLYCERIN (UNII: PDC6A3C0OX)			
HYDROXYETHYL CELLULOSE (2000 MPA.S AT 1%) (UNII: S38J6RZ N16)			
METHYLPARABEN (UNII: A218C7HI9T)			
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)			
PROPYLPARABEN (UNII: Z8IX2SC10H)			
WATER (UNII: 059QF0KO0R)			
SACCHARIN SODIUM (UNII: SB8ZUX40TY)			
SORBITOL (UNII: 506T60A25R)			

Product Characteristics			
Color	WHITE	Score	
Shape		Size	
Flavor	LEMON (Mint)	Imprint Code	
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63868-173- 12	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	08/15/2022	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part331	08/15/2022	
FINAL			

Labeler - Chain Drug Marketing Association Inc. (011920774)