

SOUND BODY NAPROXEN SODIUM- naproxen sodium tablet, film coated
Big Lots Stores, Inc.

Big Lots, Inc Naproxen Sodium Drug Facts

Active ingredient (in each caplet)

Naproxen sodium 220 mg
(naproxen 200 mg) (NSAID)*

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis
- muscular aches
- backache
- menstrual cramps
- headache
- toothache
- the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
- chest pain
- trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling
- pain gets worse or lasts more than 10 days

- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- drink a full glass of water with each dose

Adults and children 12 years and older	<ul style="list-style-type: none"> • take 1 caplet every 8 to 12 hours while symptoms last • for the first dose you may take 2 caplets within the first hour • do not exceed 2 caplets in any 8- to 12-hour period • do not exceed 3 caplets in a 24-hour period
Children under 12 years	<ul style="list-style-type: none"> • ask a doctor

Other information

- **each caplet contains:** sodium 22 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Inactive ingredients

FD&C blue no. 2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

Questions?

1-800-719-9260

Principal Display Panel

SOUNDBODY™

Compare to the Active Ingredient in Aleve® Caplets

Naproxen Sodium Tablets, 220 mg

EASY OPEN ARTHRITIS CAP

Pain Reliever / Fever Reducer (NSAID)

STRENGTH TO LAST 12 HOURS

PACKAGE NOT CHILD-RESISTANT

225 CAPLETS†

†CAPSULE-SHAPED TABLETS

Actual Size



Naproxen Sodium
Tablets, 220 mg
Pain Reliever / Fever Reducer (NSAID)
STRENGTH TO LAST 12 HOURS
PACKAGE NOT CHILD-RESISTANT
225 CAPLETS*
*CAPSULE-SHAPED TABLETS

EASY OPEN ARTHRITIS CAP

Actual Size

Compare to the Active Ingredient in Aleve® Caplets*

DO NOT USE IF PRINTED POLYMER CAP IS BROKEN OR MISSING

Drug Facts
Active ingredient
(in each caplet)
Naproxen sodium 220 mg Pain reliever/fever reducer
Naproxen 220 mg (NSAID)*...fever reducer
*Nonsteroidal anti-inflammatory drug

Purpose
Pain reliever/fever reducer

Uses
Naproxen sodium relieves minor aches and pains due to:

- muscle pain or arthritis
- menstrual cramps
- backache
- toothache
- temporarily reduces fever

Warnings
After 8 1/2 etc. Naproxen sodium may cause severe allergic reaction, especially in people allergic to aspirin. Symptoms may include hives, difficulty breathing, and swelling of the face, lips, tongue, or throat. Stop taking this product immediately if you experience these symptoms. If you have had stomach ulcers or bleeding problems, take a blood thinning (anticoagulant) or steroid drug, take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or other NSAIDs), or are taking or have taken aspirin, ibuprofen, or other NSAIDs, tell your doctor or pharmacist you are taking this product. Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Directions
If you have ever had an allergic reaction to naproxen sodium, do not take this product. If you have ever had heart surgery, ask your doctor before taking this product.

Other information

- Keep this product and this information out of the reach of children.
- Do not take more than directed.
- Do not take this product if you are pregnant, planning to get pregnant, or are breastfeeding.
- Do not take this product if you are taking or have taken aspirin, ibuprofen, or other NSAIDs, or are taking or have taken blood thinning (anticoagulant) or steroid drugs.
- Do not take this product if you are taking or have taken aspirin, ibuprofen, or other NSAIDs, or are taking or have taken blood thinning (anticoagulant) or steroid drugs.
- Do not take this product if you are taking or have taken aspirin, ibuprofen, or other NSAIDs, or are taking or have taken blood thinning (anticoagulant) or steroid drugs.

Drug Facts (continued)

Other information

- Keep this product and this information out of the reach of children.
- Do not take more than directed.
- Do not take this product if you are pregnant, planning to get pregnant, or are breastfeeding.
- Do not take this product if you are taking or have taken aspirin, ibuprofen, or other NSAIDs, or are taking or have taken blood thinning (anticoagulant) or steroid drugs.
- Do not take this product if you are taking or have taken aspirin, ibuprofen, or other NSAIDs, or are taking or have taken blood thinning (anticoagulant) or steroid drugs.
- Do not take this product if you are taking or have taken aspirin, ibuprofen, or other NSAIDs, or are taking or have taken blood thinning (anticoagulant) or steroid drugs.

Compare to the Active Ingredient in Aleve® Caplets*

Other information

- Keep this product and this information out of the reach of children.
- Do not take more than directed.
- Do not take this product if you are pregnant, planning to get pregnant, or are breastfeeding.
- Do not take this product if you are taking or have taken aspirin, ibuprofen, or other NSAIDs, or are taking or have taken blood thinning (anticoagulant) or steroid drugs.
- Do not take this product if you are taking or have taken aspirin, ibuprofen, or other NSAIDs, or are taking or have taken blood thinning (anticoagulant) or steroid drugs.
- Do not take this product if you are taking or have taken aspirin, ibuprofen, or other NSAIDs, or are taking or have taken blood thinning (anticoagulant) or steroid drugs.

Drug Facts (continued)

Other information

- Keep this product and this information out of the reach of children.
- Do not take more than directed.
- Do not take this product if you are pregnant, planning to get pregnant, or are breastfeeding.
- Do not take this product if you are taking or have taken aspirin, ibuprofen, or other NSAIDs, or are taking or have taken blood thinning (anticoagulant) or steroid drugs.
- Do not take this product if you are taking or have taken aspirin, ibuprofen, or other NSAIDs, or are taking or have taken blood thinning (anticoagulant) or steroid drugs.
- Do not take this product if you are taking or have taken aspirin, ibuprofen, or other NSAIDs, or are taking or have taken blood thinning (anticoagulant) or steroid drugs.

Drug Facts (continued)

Other information

- Keep this product and this information out of the reach of children.
- Do not take more than directed.
- Do not take this product if you are pregnant, planning to get pregnant, or are breastfeeding.
- Do not take this product if you are taking or have taken aspirin, ibuprofen, or other NSAIDs, or are taking or have taken blood thinning (anticoagulant) or steroid drugs.
- Do not take this product if you are taking or have taken aspirin, ibuprofen, or other NSAIDs, or are taking or have taken blood thinning (anticoagulant) or steroid drugs.
- Do not take this product if you are taking or have taken aspirin, ibuprofen, or other NSAIDs, or are taking or have taken blood thinning (anticoagulant) or steroid drugs.

Drug Facts (continued)

Other information

- Keep this product and this information out of the reach of children.
- Do not take more than directed.
- Do not take this product if you are pregnant, planning to get pregnant, or are breastfeeding.
- Do not take this product if you are taking or have taken aspirin, ibuprofen, or other NSAIDs, or are taking or have taken blood thinning (anticoagulant) or steroid drugs.
- Do not take this product if you are taking or have taken aspirin, ibuprofen, or other NSAIDs, or are taking or have taken blood thinning (anticoagulant) or steroid drugs.
- Do not take this product if you are taking or have taken aspirin, ibuprofen, or other NSAIDs, or are taking or have taken blood thinning (anticoagulant) or steroid drugs.

SOUND BODY NAPROXEN SODIUM			
naproxen sodium tablet, film coated			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50594-140
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)	NAPROXEN SODIUM	220 mg	

Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ05DW1A)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	BLUE (light blue)	Score	no score
Shape	OVAL	Size	12mm
Flavor		Imprint Code	L368
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50594-140-83	225 in 1 BOTTLE; Type 0: Not a Combination Product	05/07/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074661	05/07/2020	

Labeler - Big Lots Stores, Inc. (017885351)

Revised: 3/2024

Big Lots Stores, Inc.