NICOTINE- nicotine polacrilex gum, chewing Walgreens

Nicotine Polacrilex Gum Coated Ice Mint 4 mg ANDA204833 - Equate

Drug Facts

Active ingredient (in each chewing piece)

Nicotine Polacrilex 4 mg (nicotine)

Purpose

Stop smoking aid

Use

 reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings

If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

Stop use and ask a doctor if

- mouth, teeth, or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness, and rapid heartbeat
- oral blistering occurs
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets.

Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum according to the following 12 week schedule:

Weeks 1	Weeks 7	Weeks 10
to 6	to 9	to 12
1 piece	1 piece	1 piece
every	every	every
1 to 2	2 to 4	4 to 8
hours	hours	hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

Other information

- each piece contains: calcium 115 mg, sodium 11 mg
- store at 20 25°C (68 77°F)
- protect from light

Inactive ingredients

acacia, acesulfame potassium, butylated hydroxytoluene, carnauba wax, FD&C red #40, gum base, hydroxypropyl cellulose, hypromellose, natural and artificial, cinnamon flavors, sodium bicarbonate, sodium carbonate, talc, titanium dioxide and xylitol.

Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal Display Panel

Compare to the active ingredient in Nicorrette® Gum††

COATED

Nicotine Gum

NICOTINE POLACRILEX GUM USP, 4 mg (NICOTINE)

STOP SMOKING AID

SUGAR FREE

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE WITHIN 30 MINUTES OF WAKING UP

If you smoke your first cigarette $\underline{\mathsf{MORE\ THAN}}\ 30\ \mathsf{MINUTES}\ \mathsf{after\ waking\ up},\ \mathsf{use\ Nicotine}\ \mathsf{Polacrilex\ gum},\ 2\ \mathsf{mg}$

Cinnamon Flavor

TO INCREASE YOUR SUCCESS IN QUITTING:

- 1. You must be motivate to quit
- 2. Use Enough Chew at least 9 pieces of Nicotine Polacrilex Gum per day during the first six weeks
- 3. Use long enough Use Nicotine Polacrilex Gum for the full 12 weeks
- 4. Use with a Support Program as directed in the enclosed User's Guide

To remove the gum, tear off single unit

Peel off backing starting at corner with loose edge

Push gum through foil

- NOT FOR SALE TO THOSE UNDER 18 YEARS OF AGE.
- PROOF OF AGE.
- NOT FOR SALE IN VENDING MACHINES OR FROM SOURCE WHERE PROOF OF AGE CANNOT BE VERIFIED

††This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare, L.P., owner of the registered trademark Nicorette® .

BLISTER PACKAGED FOR YOUR PROTECTION. DO NOT USE IF INDIVIDUAL SEALS ARE OPEN OR TORN.

DISTRIBUTED BY: WALGREEN CO.

200 WILMOT RD., DEERFIELD, IL 60015

Package Label

TO INCREASE YOUR SUCCESS IN QUITTING:

- 1. You must be motivated to quit.
- 2. Use Enough Chew at least 9 pieces of Nicotine Polacrilex Gum per day during the first
- 3. Use Long Enough -Use Nicotine Polacrilex Gum for the full 12 weeks.
- 4. Use with a support program as directed in the enclosed User's Guide.









For tools, resources and motivation, please visit Walgreens.com/QuitSmoking

NDC 0363-3911-10

Walgreens to the active



COATED

Nicotine

NICOTINE POLACRILEX GUM 4 mg (NICOTINE) STOP SMOKING AID

Sugar Free







ACTUAL SIZE

Cinnamon

4 mg EACH

POR THOSE WHO SMOKE THEIR FIRST CIGARETTE WITHIN 30 MINUTES OF WAXING UP.
If you smoke your first cigarette MORE THAN 30 MINUTES after waking up, use Nicotine Polacrifer Gum, 2 mg.

THEFT SURVEILLANCE TAG AREA

YART XOB ЬУРЕЯ МИГЛ-ГАУЕВ

CRI 1-811-123-3832 Warqal-Luqai avw-zuw EZI (Inestious ol comuleurs)

celulose, hypromellose, natural and artificial cirmamon flavors, sodium bicarbonato, sodium carbonate, sucralose, taic, thanium dioxide, vylitol Inactive ingredients acase, aceantane posses indrognopy hydroxylaluse, canasta wax, FD&C ned #40, gum base, hydroxypropy

- bogin using the gum on your quik day
 iii you emake your thest cigarette more than 30 minutes efter we
- Directions

 If you are notes 18 years of age, ask a doctor before use
 If you are notes 18 years of age, ask a doctor before use of the recessed User's Baids for complete
 three-baids and other interesting the recessed User's Baids for complete
 three-baids and the second of the secon

Neep out of rescul et children and gets. Pieces of nicotive gun may here eccept in other active the second in the mass of matching and eccept and the way.

Contact a Peleon Control Centre (1-600-5222-1522) right sawy.

- Stop use and ask a doctor if m mouth, when two perplations occur as mouther they two problems occur as mouth sets we do incolor conditions and mouth they doctored demand betaking occurs and mouth teachers are difficulty breathing or as and betaking occurs. In one betaking occurs of an aleight mession (such as difficulty breathing or seals)

does may need to be adjusted.

PRODUCT OF IRELAND M3OHG01SS-F HEV-0922 PLD-C620A FC008332




ILEM 892169 W10024-0922-L

walgreens.com ©2021 Walgreen Co.

100% GUARANTEED

200 WILMOT RD., DEERFIELD, IL 60015



WALGREENS Coated Nicotine Gum Cinnamon Flavor

NICOTINE

nicotine polacrilex gum, chewing

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:0363-3911

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name Basis of Strength Strength

NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)

NICOTINE

4 mg

Inactive Ingredients		
Ingredient Name	Strength	
ACACIA (UNII: 5C5403N260)		
ACESULFAME POTASSIUM (UNII: 230V73Q5G9)		
CARNAUBA WAX (UNII: R12CBM0EIZ)		
FD&C RED NO. 40 (UNII: WZB9127XOA)		
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)		
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)		
SODIUM BICARBONATE (UNII: 8MDF5V39QO)		
SODIUM CARBONATE (UNII: 45P3261C7T)		
TALC (UNII: 7SEV7J4R1U)		
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)		
XYLITOL (UNII: VCQ006KQ1E)		
SUCRALOSE (UNII: 96K6UQ3ZD4)		
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)		

Product Characteristics			
Color	pink (Dark)	Score	no score
Shape	SQUARE	Size	14mm

Flavor	CINNAMON	Imprint Code	
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-3911- 16	160 in 1 CARTON	10/30/2020	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:0363-3911- 10	100 in 1 CARTON	10/30/2020	
2		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079219	10/30/2020	

Labeler - Walgreens (008965063)

Revised: 10/2022 Walgreens