GRAVOL- dimenhydrinate tablet, coated Church & Dwight Co., Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Gravol™ Motion Sickness Relief - Coated Tablets

Active ingredient (in each tablet)

Dimenhydrinate 50 mg

Purpose

Antiemetic

Use

For prevention and treatment of these symptoms associated with motion sickness:

- nausea
- vomiting
- dizziness

Warnings

Do not use

for children under 2 years of age unless directed by a doctor.

Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are taking

sedatives or tranquilizers

When using this product

- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase dizziness
- be careful when driving a motor vehicle or operating machinery or any activity that requires alertness

If pregnant or breast-feeding,

ask a doctor before use.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- to prevent motion sickness, the first dose should be taken ½ to 1 hour before starting activity
- to prevent or treat motion sickness, see below:

and over:

- take 1 to 2 tablets every 4-6 hours
- adults and children 12 years do not take more than 8 tablets in 24 hours, or as directed by a doctor
 - give 1/2 to 1 tablet every 6-8 hours

children 6 to under 12 years:

- do not give more than 3 tablets in 24 hours, or as directed by a doctor
- give 1/4 to 1/2 tablet every 6-8 hours

children 2 to under 6 years:

• do not give more than 1-1/2 tablets in 24 hours, or as directed by a doctor

Other information

- store at room temperature 59º-77ºF (15º-25ºC)
- do not use if packet is torn

Inactive ingredients

Corn starch, croscarmellose sodium, FD&C yellow 6 aluminum lake, iron oxide yellow, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, silicon dioxide, talc, titanium dioxide

Questions or comments?

1-800-952-5080 M-F 9am to 5pm ET

PRINCIPAL DISPLAY PANEL

OVER 70 YEARS OF

QUALITY & TRUST†

Prevents and Treats:

Nausea & Vomiting

Dizziness*

*Prevention and treatment of nausea,

vomiting and dizziness due to motion sickness

Gravol™ COATED TABLETS

Gravol™

MOTION SICKNESS RELIEF

Dimenhydrinate 50 mg Tablets - Antiemetic

10

COATED TABLETS

TASTE FREE

EASY TO SWALLOW





Gravo COATED TABLETS

T3 mg2 at ma9 7-M 0802-586-008-F fs sulls3 Sanoiteaup

Inactive ingredients to a state creatments as property of a property of

outstagmet moon is state- **notisemnohni** 19400 met al teoloog til seu tan ab+ $(0.25, ^{\circ}2), 1.77, ^{\circ}62$

от ав фільськой by а фостог 6-8 hours • do not give more than 1-15 tablets in 24 hours, Januar Para and properties in 24 hours are the Para and P

every 4-6 hours -do not bake more than 8 bablets in 24 hours or as directive dy a doctor echibran 6 to under 12 years -give is to 1 tablet every *adults and children 12 years and over: *take 1 to 2 tablets

Directions - to present motion sickness, the first does should be taken it to it now before starting activity - to prevent or treat motion alongs, see the section of the property of the property of the property of the pro-teat of deliberant by sense and seem dates it to Stablets

overtices, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Roop out at reach of children, in case at accidental the professor or beauting, ask a doctor before use.

seurobei sessi Aprison (n. en) estrato de popular. usok juduses glowajuese «pe cerupin nijiku gujikub s uotou When using this product *marked drowsiness may occur exold alcoholic drinks *alcohol, sedatives, and tranquilizers satismbuez jo savgepos

As a cock to the properties of the process of the p

Do not use for children under 2 years of age unless directed

SSOURTROM BURBLION ROSSESIA TRADUNOR HODDEL BUM beforese anothers sent to tremtest bus naturated art to

Active ingredient (in each tablet) Purpose

2501,51915

Church & Dwight Co., Inc., Ewing, NJ 08628 Graval is a trademark of Church & Dwight Canada Corp. Made in Canada





CRPC-00162-01 72022776

GRAVOL

dimenhydrinate tablet, coated

Decade		Inform	:
Prod	IICT.	Intorm	ATION
FIGU	ull		ıatıvı

HUMAN OTC DRUG NDC:10237-941 Item Code (Source) **Product Type**

ORAL **Route of Administration**

Active Ingredient/Active Moiety

Ingredient Nan	ne
----------------	----

Basis of Strength Strength

DIMENHYDRINATE (UNII: JB937PER5C) (DIPHENHYDRAMINE - UNII:8GTS82S83M)

DIMENHYDRINATE

50 mg

Inactive Ingredients		
Ingredient Name	Strength	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)		
MAGNESIUM STEARATE (UNII: 70097M6I30)		
SILICON DIOXIDE (UNII: ETJ7Z 6XBU4)		
TALC (UNII: 7SEV7J4R1U)		
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)		
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)		
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)		
CROSCARMELLOSE (UNII: 029TFK992N)		
STARCH, CORN (UNII: O8232NY3SJ)		
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)		
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)		
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)		

Product Characteristics				
Color	orange	Score	4 pieces	
Shape	ROUND	Size	8mm	
Flavor		Imprint Code		
Contains				

F	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:10237- 941-10	1 in 1 CARTON	03/02/2022		
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product			

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part336	03/02/2022	

Labeler - Church & Dwight Co., Inc. (001211952)

Establishment

Name	Address	ID/FEI	Business Operations
Church & Dwight Canada Corp.		253933600	manufacture(10237-941)

Revised: 3/2022 Church & Dwight Co., Inc.