

**IBUPROFEN- ibuprofen suspension**  
**CVS Pharmacy**

-----  
**CVS Pharmacy, Inc. Ibuprofen Oral Suspension Drug Facts**

**Active ingredient (in each 5 mL)**

Ibuprofen 100 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

**Purposes**

Pain reliever/fever reducer

**Uses**

temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more

than directed or for longer than directed.

**Sore throat warning:** Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

### **Do not use**

- if the child has ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

### **Ask a doctor before use if**

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

### **Ask a doctor or pharmacist before use if the child is**

- under a doctor's care for any serious condition
- taking any other drug

### **When using this product**

- give with food or milk if stomach upset occurs

### **Stop use and ask a doctor if**

- child experiences any of the following signs of stomach bleeding:
- feels faint
- vomits blood
- has bloody or black stools
- has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
- chest pain
- trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

## Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

## Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.
- if needed, repeat dose every 6-8 hours
- do not use more than 4 times a day
- replace original bottle cap to maintain child resistance
- wash dosage cup after each use

---

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL)**
under 24 lbs	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

\*\* or as directed by a doctor

## Other information

- **each 5 mL contains:** sodium 2 mg
- do not use if printed neckband is broken or missing
- store at 20-25°C (68-77°F)

- do not freeze

**Inactive ingredients**

anhydrous citric acid, FD&C red #40, glycerin, high fructose corn syrup, hypromellose, natural and artificial bubblegum flavors, polysorbate 80, purified water, sodium benzoate, sorbitol solution, xanthan gum

**Questions or comments?**

**1-800-719-9260**

**Principal Display Panel**

CVS Health®

Compare to the active ingredient in Children's Motrin®

Suspension Liquid

For Ages 2 to 11 Years

Children's Ibuprofen

Oral Suspension

100 mg per 5 mL

Pain reliever;

Fever reducer (NSAID)

Gluten free

Alcohol free

SEE NEW WARNINGS

LASTS UP TO 8 HOURS

Bubble Gum Flavor

4 FL OZ (120 mL)



Compare to the active ingredient in Children's Motrin®

Suspension Liquid

NDC 59779-166-26

For Ages 2 to 11 Years  
Children's

**Ibuprofen**

Oral Suspension  
100 mg per 5 mL

Pain reliever;  
Fever reducer (NSAID)

- Gluten free
- Alcohol free

**SEE NEW WARNINGS**



Bubble Gum Flavor

4 FL OZ (120 mL)

Important: Read all product information before using. Keep this box for important information. This product is intended for use in children.

### Drug Facts

**Active ingredient** (in each 5 mL) **Purposes**  
Ibuprofen 100 mg (NSAID)\* .....Pain reliever/  
fever reducer  
\*nonsteroidal anti-inflammatory drug

### Uses

- temporarily:
  - relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
  - reduces fever

### Warnings

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug

■ takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
■ takes more or for a longer time than directed  
**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Sore throat warning:** Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

### Do not use

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer

Compare to the active ingredient in Children's Motrin®  
**CVS Health**  
For Ages 2 to 11 Years  
**Ibuprofen**  
Oral Suspension  
100 mg per 5 mL  
Pain reliever;  
Fever reducer (NSAID)  
Suspension Liquid

### Drug Facts (continued)

- right before or after heart surgery

### Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

### Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

### When using this product

- give with food or milk if stomach upset occurs

### Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding:
  - feels faint
  - vomits blood
  - has bloody or black stools
  - has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

### Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed

### Drug Facts (continued)

- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.
- if needed, repeat dose every 6-8 hours
- do not use more than 4 times a day
- replace original bottle cap to maintain child resistance
- wash dosage cup after each use

### Dosing Chart

Weight (lb)	Age (yr)	Dose (mL)**
under 24 lbs	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

\*\*or as directed by a doctor

### Other information

- each 5 mL contains: sodium 2 mg
- do not use if printed neckband is broken or missing
- store at 20-25°C (68-77°F)
- do not freeze

**Inactive ingredients** anhydrous citric acid, FD&C red #40, glycerin, high fructose corn syrup, hypromellose, natural and artificial bubblegum flavors, polysorbate 80, purified water, sodium benzoate, sorbitol solution, xanthan gum

### Questions or comments?

1-800-719-9260

This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Children's Motrin®.

**Distributed by: CVS Pharmacy, Inc.**  
One CVS Drive, Woonsocket, RI 02895

© 2017 CVS/pharmacy  
CVS.com® 1-800-SHOP CVS

V-17160

**CVS Quality**  
Money Back Guarantee



#191238



16626 17 C9

# Ibuprofen

Children's  
For ages 2 to 11 Years

Suspension Liquid

Compare to the  
active ingredient in  
Children's Motrin®

CVS  
Health™

Oral Suspension  
100 mg per 5 mL  
Pain reliever;  
Fever reducer (NSAID)

## Drug Facts (continued)

- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, or kidney disease
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- give with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding
  - feels faint
  - vomits blood
  - has bloody or black stools
  - has stomach pain that does not get better
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

## Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- shake well before using
- mL = milliliter; tsp = teaspoonful
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.
- if needed, repeat dose every 6-8 hours
- do not use more than 4 times a day
- replace original bottle cap to maintain child resistance
- wash dosage cup after each use

## Drug Facts (continued)

### Dosing Chart

Weight (lb)	Age (yr)	Dose (mL or tsp)**
under 24 lbs	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL (1 tsp)
36-47 lbs	4-5 years	7.5 mL (1½ tsp)
48-59 lbs	6-8 years	10 mL (2 tsp)
60-71 lbs	9-10 years	12.5 mL (2½ tsp)
72-95 lbs	11 years	15 mL (3 tsp)

\*\*or as directed by a doctor

## Other information

- each 5 mL (1 tsp) contains: sodium 2 mg
- do not use if printed neckband is broken or missing
- store at 20-25°C (68-77°F)
- do not freeze
- see bottom panel for lot number and expiration date

**Inactive ingredients** anhydrous citric acid, FD&C red #40, glycerin, high fructose corn syrup, hypromellose, natural and artificial bubblegum flavors, polysorbate 80, purified water, sodium benzoate, sorbitol solution, xanthan gum

## Questions or comments?

1-800-719-9260

†This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Children's Motrin®.

Distributed by: CVS Pharmacy, Inc.  
One CVS Drive, Woonsocket, RI 02895  
© 2015 CVS/pharmacy  
CVS.com® 1-800-SHOP CVS  
V-17160

CVS Quality  
Money Back Guarantee



PLEASE RECYCLE





# IBUPROFEN

ibuprofen suspension

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:59779-166
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)		IBUPROFEN	100 mg in 5 mL
Inactive Ingredients			
Ingredient Name			Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)			
FD&C RED NO. 40 (UNII: WZB9127XOA)			
GLYCERIN (UNII: PDC6A3C0OX)			
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)			
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)			
POLYSORBATE 80 (UNII: 6OZP39ZG8H)			
WATER (UNII: 059QF0KO0R)			
SODIUM BENZOATE (UNII: OJ245FE5EU)			
SORBITOL (UNII: 506T60A25R)			
XANTHAN GUM (UNII: TTV12P4NEE)			
Product Characteristics			
Color	PINK (light)	Score	
Shape		Size	

Flavor		BUBBLE GUM	Imprint Code	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:59779-166-26	1 in 1 CARTON	08/27/2002	
1		120 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:59779-166-34	1 in 1 CARTON	04/09/2008	
2		240 mL in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:59779-166-28	1 in 1 CARTON	09/08/2015	01/31/2020
3		148 mL in 1 BOTTLE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA		ANDA074937	08/27/2002	

**Labeler** - CVS Pharmacy (062312574)

Revised: 12/2022

CVS Pharmacy