

HYDROGEN PEROXIDE- hydrogen peroxide solution
Henry Schein Inc.

Hydrogen Peroxide

Drug Facts

Active Ingredient

Hydrogen Peroxide (stabilized) 3%

Purpose

First Aid Antiseptic/Oral Debriding Agent

Uses

- first aid to help prevent the risk of infection in minor cuts, scrapes and burns
- aids in removal of phlegm, mucous, or other secretions associated with occasional sore mouth

Warnings

For external use only

Do not use

- in the eyes or apply over large areas of the body
- longer than 1 week

Ask a doctor before use if you have deep puncture wounds, animal bites or serious burns.

Stop use and ask a doctor if

- the condition persists or gets worse
- sore mouth symptoms do not improve in 7 days
- irritation, pain, or redness persists or worsens
- swelling, rash, or fever develops

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

First Aid Antiseptic

**Oral Debriding Agent (oral
rinse):**

Adults and children 2 years of
age & over:

- | | |
|---|---|
| <ul style="list-style-type: none">• clean the affected area• apply a small amount of product on the area 1 to 3 times a day• may be covered with a sterile bandage• if bandaged, let dry first | <ul style="list-style-type: none">• mix with an equal amount of water• swish around in the mouth over the affected area for at least 1 minute and then spit out• use up to 4 times daily after meals and at bedtime or as directed by a dentist or doctor |
|---|---|

- children under 12 years of age should be supervised in the use of this product
- children under 2 years of age: consult a dentist or doctor

Other Information

- Keep tightly closed in a cool dark place
- Do not shake bottle
- Hold away from face when opening

Inactive Ingredient

Purified Water

Questions or Comments?

1-800-472-4346

PRINCIPAL DISPLAY PANEL - 3.79 L Bottle Label

Item # 112-7070
NDC 0404-0001-01

HENRY SCHEIN®

3 % Hydrogen Peroxide Solution

U.S.P. FIRST AID ANTISEPTIC

Tamper evident cap for your protection. If ring-band is detached from cap or missing do not use.

Distributed by (in US Only):

HENRY SCHEIN INC.

135 Duryea Road
Melville, NY 11747 USA

Rev. Date 2018/06 Made in the USA

128 FL OZ (1 Gal.) • 3.79 L

Item # 112-7070

NDC 0404-0001-01



3% Hydrogen Peroxide Solution

U.S.P. FIRST AID ANTISEPTIC
 Tamper evident cap for your protection. If ring-band is detached from cap or missing do not use.

Distributed by (in US Only):
HENRY SCHEIN INC.
 135 Duryea Road
 Melville, NY 11747 USA
 Rev Date: 2018/06 Made in the USA

128 FL OZ (1 Gal.) • 3.79 L

+H658112707009



1-07A0013N05HSC

| Drug Facts | |
|--|---|
| Active Ingredient | Purpose |
| Hydrogen Peroxide (stabilized) 3% | First Aid Antiseptic/Oral Debriding Agent |
| Uses • first aid to help prevent the risk of infection in minor cuts, scrapes and burns • aids in removal of phlegm, mucous, or other secretions associated with occasional sore mouth | |
| Warnings For external use only | |
| Do not use • in the eyes or apply over large areas of the body • longer than 1 week • on deep or puncture wounds, animal bites, or on serious burns. | |
| Ask a doctor before use if you have deep or puncture wounds, animal bites or serious burns. | |
| Stop use and ask a doctor if • the condition persists or gets worse • sore mouth symptoms do not improve in 7 days • irritation, pain, or redness persists or worsens • swelling, rash, or fever develops | |
| Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away (1-800-222-1222). | |
| Directions | |
| First Aid Antiseptic | Oral Debriding Agent (oral rinse): |
| <ul style="list-style-type: none"> • clean the affected area • apply a small amount of product on the area 1 to 3 times a day • may be covered with a sterile bandage • if bandaged, let dry first | <ul style="list-style-type: none"> • Adults and children 2 years of age & over: <ul style="list-style-type: none"> • mix with an equal amount of water • swish around in the mouth over the affected area for at least 1 minute and then spit out • use up to 4 times daily after meals and at bedtime or as directed by a dentist or doctor • children under 12 years of age should be supervised in the use of this product • children under 2 years of age: consult a dentist or doctor |
| Other Information • Keep tightly closed in a cool dark place • Do not shake bottle • Hold away from face when opening | |
| Inactive Ingredient Purified Water | |
| Questions or Comments? 1-800-472-4346 | |

HYDROGEN PEROXIDE

hydrogen peroxide solution

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:0404-0001 |
| Route of Administration | TOPICAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|-------------------|---------------|
| HYDROGEN PEROXIDE (UNII: BBX060AN9V) (HYDROGEN PEROXIDE - UNII:BBX060AN9V) | HYDROGEN PEROXIDE | 30 mg in 1 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|---------------------------------|----------|
| WATER (UNII: 059QF0KO0R) | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:0404-0001-01 | 3785 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | 06/01/2014 | |
| 2 | NDC:0404-0001-08 | 237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | 06/01/2014 | |
| 3 | NDC:0404-0001-16 | 473 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | 06/01/2014 | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M003 | 06/01/2014 | |

Labeler - Henry Schein Inc. (012430880)

Revised: 10/2023

Henry Schein Inc.