

HYDROCORTISONE ACETATE- hydrocortisone acetate suppository
Rebel Distributors Corp

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Hydrocortisone Acetate


PROCTOCORT®

Hydrocortisone Acetate

Rectal Suppositories, 25 mg

DESCRIPTION

Each Hydrocortisone Acetate 25 mg Suppository contains 25 mg hydrocortisone acetate in a hydrogenated vegetable oil base. Hydrocortisone acetate is a corticosteroid. Chemically, hydrocortisone acetate is a pregn-4-ene-3, 20-dione, 21-(acetyloxy)-11, 17-dihydroxy-(11 β) with the following structural formula:



CLINICAL PHARMACOLOGY

In normal subjects, about 26% of hydrocortisone acetate is absorbed when the suppository is applied to the rectum. Absorption of hydrocortisone acetate may vary across abraded or inflamed surfaces.

Topical steroids are primarily effective because of their anti-inflammatory, anti-pruritic and vasoconstrictive action.

INDICATIONS AND USAGE

For use in inflamed hemorrhoids, postirradiation (factual) proctitis; as an adjunct in the treatment of chronic ulcerative colitis; cryptitis; and other inflammatory conditions of anorectum and pruritus ani.

CONTRAINDICATIONS

Hydrocortisone Acetate Suppositories are contraindicated in those patients having a history of hypersensitivity to any of the components.

PRECAUTIONS

Do not use unless adequate proctologic examination is made.

If irritation develops, the product should be discontinued and appropriate therapy instituted.

In the presence of an infection, the use of an appropriate antifungal or antibacterial agent should be instituted. If a favorable response does not occur promptly, the hydrocortisone acetate should be discontinued until the infection has been adequately controlled.

Carcinogenesis: No long term studies in animals have been performed to evaluate the carcinogenic potential of corticosteroid suppositories.

Pregnancy Category C: In laboratory animals, topical steroids have been associated with an increase in the incidence of fetal abnormalities when gestating females have been exposed to rather low dosage levels. There are no adequate and well controlled studies in pregnant women. Hydrocortisone Acetate Suppositories should only be used during pregnancy if the potential benefit justifies the risk to the fetus. Drugs of this class should not be used extensively on pregnant patients, in large amounts, or for prolonged periods of time.

It is not known whether this drug is excreted in human milk and because many drugs are excreted in human milk and because of the potential for serious adverse reactions in nursing infants from Hydrocortisone Acetate Suppositories, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

ADVERSE REACTIONS

The following local adverse reactions have been reported with hydrocortisone acetate suppositories; burning, itching, irritation, dryness, folliculitis, hypopigmentation, allergic contact dermatitis, secondary infection.

DRUG ABUSE AND DEPENDENCE

Drug abuse and dependence have not been reported in patients treated with hydrocortisone acetate suppositories.

OVERDOSAGE

If signs and symptoms of systemic overdosage occur, discontinue use.

DOSAGE AND ADMINISTRATION

For rectal administration. Detach one suppository from strip of suppositories. Remove the wrapper. Avoid excessive handling of the suppository which is designed to melt at body temperature. Insert suppository into the rectum with gentle pressure, pointed end first. Insert one suppository in the rectum twice daily, morning and night for two weeks, in nonspecific proctitis. In more severe cases, one

suppository three times a day or two suppositories twice daily. In factitial proctitis, the recommended duration of therapy is six to eight weeks or less, according to the response of the individual case.

HOW SUPPLIED

Box of 12 suppositories - NDC 21695-731-12

Rx only.

Store at 20°–25°C (68°–77°F). See USP Controlled Room Temperature. Store away from heat. Protect from freezing.

For Inquiries call: 1-866-207-5636

Distributed by:

County Line Pharmaceuticals, LLC


Brookfield, WI 53005

Repackaged by:

Rebel Distributors Corp

Thousand Oaks, CA 91320

PRINCIPAL DISPLAY PANEL



HYDROCORTISONE ACETATE

hydrocortisone acetate suppository

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:21695-731(NDC:43199-021)
Route of Administration	RECTAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
HYDROCORTISONE ACETATE (UNII: 3X7931PO74) (HYDROCORTISONE - UNII:WI4X0X7BPJ)	HYDROCORTISONE ACETATE	30 mg

Inactive Ingredients

Ingredient Name	Strength
HYDROGENATED PALM OIL (UNII: 257THB963H)	
COLLOIDAL SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
BUTYLATED HYDROXYANISOLE (UNII: REK4960K2U)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:21695-731-12	12 in 1 BOX		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
Unapproved drug other		06/01/2004	

Labeler - Rebel Distributors Corp (118802834)

Establishment

Name	Address	ID/FEI	Business Operations
Rebel Distributors Corp		118802834	RELABEL, REPACK

Revised: 6/2011

Rebel Distributors Corp