

HIBICLENS FOR GEORGIA PACIFIC- chlorhexidine gluconate solution

Georgia-Pacific Consumer Products LP

Hibiclens for Georgia Pacific

Chlorhexidine gluconate solution 4% w/v

Antiseptic

Uses

- antimicrobial skin cleanser helps reduce bacteria that potentially can cause disease
- surgical hand scrub
- healthcare personnel handwash

Warnings

For external use only

Allergy alert:

This product may cause a severe allergic reaction. Symptoms may include:

- wheezing/difficulty breathing
- shock
- facial swelling
- hives
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use

- If you or the patient is allergic to chlorhexidine gluconate or any other ingredient in this product

When using this product

- keep out of eyes, ears, and mouth. May cause serious and permanent eye injury if permitted to enter and remain in the eye or may cause deafness when instilled in the middle ear through perforated eardrums.
- if contact occurs in any of these areas, rinse with cold water right away
- do not use routinely if you have wounds which involve more than the superficial layers of the skin

Stop use and ask a doctor if

irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children.

If swallowed, get medical help or contact Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- surgical hand scrub. Wet hands and forearms with water. Scrub for 3 minutes with about 5mL (8 pumps) of the product with a brush. Rinse thoroughly under running water. Repeat. Dry thoroughly.
- healthcare personnel handwash. Wet hands with water. Dispense about 5mL (8 pumps) of the product into cupped hands and wash in a vigorous manner for 15 seconds. Rinse and dry thoroughly.

Other information

- store between 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients

fragrance, gluconalactone, isopropyl alcohol 4% w/v, lauramine oxide, poloxamer 237, purified water and red 40

HIBICLENS Chlorhexidine Gluconate Solution
For Georgia-Pacific 4% w/v
Antiseptic/Antimicrobial Skin Cleanser

Drug Facts		Drug Facts (continued)	
Active ingredient Chlorhexidine gluconate solution 4% w/v.....Antiseptic	Purpose	do not use routinely if you have wounds which involve more than the superficial layers of the skin	
Uses <ul style="list-style-type: none"> antimicrobial skin cleanser helps reduce bacteria that potentially can cause disease surgical hand scrub healthcare personnel handwash 		Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition. Keep out of reach of children. If swallowed, get medical help or contact Poison Control Center right away.	
Warnings For external use only Allergy alert: This product may cause a severe allergic reaction. Symptoms may include: <ul style="list-style-type: none"> wheezing/difficulty breathing shock facial swelling hives rash If an allergic reaction occurs, stop use and seek medical help right away.		Directions <ul style="list-style-type: none"> Use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. surgical hand scrub. Wet hands and forearms with water. Scrub for 3 minutes with about 5mL (8 pumps) of the product with a brush. Rinse thoroughly under running water. Repeat. Dry thoroughly. healthcare personnel handwash. Wet hands with water. Dispense about 5mL (8 pumps) of the product into cupped hands and wash in a vigorous manner for 15 seconds. Rinse and dry thoroughly. 	
Do not use <ul style="list-style-type: none"> if you or the patient is allergic to chlorhexidine gluconate or any other ingredient in this product. 		Other information <ul style="list-style-type: none"> store between 20-25°C (68-77°F) avoid excessive heat above 40°C (104°F) 	
When using this product <ul style="list-style-type: none"> keep out of eyes, ears, and mouth. May cause serious and permanent eye injury if permitted to enter and remain in the eye or may cause deafness when instilled in the middle ear through perforated eardrums. if contact occurs in any of these areas, rinse with cold water right away 		Inactive ingredients fragrance, gluconolactone, isopropyl alcohol 4% w/v, lauramine oxide, poloxamer 237, purified water and red 40	

40.5 fl oz (1200 mL)
NDC 54 622-0575-75 5000008943/000/04

HIBICLENS FOR GEORGIA PACIFIC

chlorhexidine gluconate solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:54622-575
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
-----------------	-------------------	----------

CHLORHEXIDINE GLUCONATE (UNII: MOR84MUD8E) (CHLORHEXIDINE - UNII:R4KO0DY52L)		CHLORHEXIDINE GLUCONATE	4 g in 100 mL	
Inactive Ingredients				
Ingredient Name		Strength		
LAURAMINE OXIDE (UNII: 4F6FC4MI8W)				
GLUCONOLACTONE (UNII: WQ29KQ9POT)				
FD&C RED NO. 40 (UNII: WZB9127XOA)				
WATER (UNII: 059QF0KO0R)				
POLOXAMER 237 (UNII: JT3U72U71K)				
ISOPROPYL ALCOHOL (UNII: ND2M416302)				
Product Characteristics				
Color	red	Score		
Shape		Size		
Flavor		Imprint Code		
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:54622-575-73	1200 mL in 1 BOTTLE, DISPENSING; Type 0: Not a Combination Product	03/01/2022	
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
NDA	NDA017768	03/01/2022		

Labeler - Georgia-Pacific Consumer Products LP (806142217)

Revised: 12/2022

Georgia-Pacific Consumer Products LP