

GOOD NEIGHBOR PHARMACY ASPIRIN- aspirin tablet, chewable Amerisource Bergen

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Amerisource Bergen Aspirin Drug Facts

Active ingredient (in each chewable tablet)

Aspirin 81 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

for the temporary relief of minor aches and pains

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present
 - new symptoms occur
 - ringing in the ears or a loss of hearing occurs

These could be signs of a serious condition

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor

- children under 12 years: ask a doctor

Other information

- store at 20-25°C (68-77°F)

Inactive ingredients

corn starch, dextrose excipient, FD&C yellow no. 6 aluminum lake, flavors, saccharin sodium

Questions or comments?

1-800-719-9260

Principal Display Panel

Compare to Bayer® Chewable Aspirin active ingredient

GOOD NEIGHBOR PHARMACY®

See New Warning

Aspirin 81 mg

LOW DOSE

Pain Reliever (NSAID)

Orange Flavored Chewable Aspirin

Aspirin Regimen – Talk to your doctor or other healthcare provider before using this product for your heart.

36 Tablets | 81 mg Each

actual size



Compare to Bayer® Chewable Aspirin active ingredient™

NDC 24385-028-68

See New Warning

Aspirin 81 mg

LOW DOSE

Pain Reliever (NSAID)

Orange Flavored Chewable Aspirin

Aspirin Regimen - Talk to your doctor or other healthcare provider before using this product for your heart.

36 Tablets | 81 mg Each

actual size



46768 29 C9



DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING
ABC#: 10051308

Drug Facts

Active ingredient Purpose (in each chewable tablet)
Aspirin 81 mg (NSAID).....Pain reliever
*nonsteroidal anti-inflammatory drug

Uses for the temporary relief of minor aches and pains

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction on which may include:

- hives
- asthma (wheezing)
- facial swelling
- shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

Drug Facts (continued)

■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

■ have 3 or more alcoholic drinks every day while using this product

■ take more or for a longer time than directed

Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer

■ for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor

Ask a doctor before use if

■ stomach bleeding warning applies to you

■ you have a history of stomach problems, such as heartburn

■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

■ you are taking a diuretic ■ you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for

■ gout ■ diabetes ■ arthritis

Stop use and ask a doctor if

■ an allergic reaction occurs. Seek medical help right away.

■ you experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better

Drug Facts (continued)

■ pain gets worse or lasts more than 10 days

■ fever gets worse or lasts more than 3 days

■ redness or swelling is present

■ new symptoms occur

■ ringing in the ears or a loss of hearing occurs

These could be signs of a serious condition

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 chewable tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

Other information

■ store at 20-25°C (68-77°F)

Drug Facts (continued)
Inactive ingredients: corn starch, dextrose, excipient, FD&C yellow no. 6 aluminum lake, flavin, sodium saccharin sodium

Questions or comments?
1-800-716-0950

*Good Neighbor Pharmacy® Aspirin is not manufactured or distributed by Bayer HealthCare LLC, distributor of Bayer® brands of aspirin.
Distributed by:
Aspirin used by:
Aristo Pharmaceuticals
1 Westport Avenue
Conshohocken, PA 19048
Distributed by:
www.mingna.com

GLUTEN FREE

SUBSTITUTION

GOOD NEIGHBOR PHARMACY ASPIRIN

aspirin tablet, chewable

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:24385-028
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	

Product Characteristics

Color	ORANGE	Score	no score
Shape	ROUND	Size	8mm
Flavor	ORANGE	Imprint Code	L467
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:24385-028-68	1 in 1 CARTON	03/19/1993	
1		36 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	03/19/1993	

Labeler - Amerisource Bergen (007914906)